



Oregon

John A. Kitzhaber, MD, Governor

Board of Massage Therapists

748 Hawthorne Ave NE

Salem, OR 97301

Phone: (503) 365-8657

Fax: (503) 385-4465

www.oregon.gov/OBMT

Testimony to Senate Health Care and Human Services

SB 387 – Massage Facility Permits

To: Chair Monnes-Anderson

Vice-Chair Kruse

Senator Knopp

Senator Shields

Senator Hayward

Chair Monnes-Anderson and members of the Senate Health Care and Human Services Committee. For the record I am David Fredrickson, Chair of the Oregon Board of Massage Therapists. I reside and practice in Ashland Oregon. I have been a licensed massage therapist in Oregon for 26 years, initially licensing in 1987. I am here today to express support of SB 387.

My massage practice focuses on clinical, therapeutic massage for chronic pain conditions. I have been an educator in the field for 13 years. I co-owned the Ashland Institute of Massage, a premier small private school from 2003 to 2012. I was appointed to the Oregon Board of Massage Therapists by Governor Ted Kulongoski in February 2010 and am in my second year of service as Board Chair. I am a member of the Policy committee of the Federation of State Massage Therapy Boards, a national membership organization representing 42 State Massage Boards.

Nationwide human trafficking is the 'Number One' shared issue in the massage regulatory industry. Many states are reviewing their statutory options for regulating the business sectors known to attract traffickers. Massage is a key area of concern. Human trafficking in massage facilities affects every major municipality along the Interstate-5 corridor from San Diego to Seattle. The practice of trafficking humans, defined by the United Nations as a form of "Modern Slavery", brings together the issues of public health and safety, domestic and workplace intimidation, violence and cohesion, human rights, teen and child abuse, prostitution, immigration fraud, and national security violations.

Massage Facilities owners who traffic and prostitute their "employees" play a dangerous game with the health and safety of the citizens of Oregon. Under current Oregon law owners of massage facilities are not required to hire State Licensed Massage Therapists. A key feature of Senate Bill 387, sponsored by Sen. Alan Bates MD of Medford, would require that Oregon State Licensed Massage Therapists be employed or provided working opportunities in massage facilities that meet permit requirements.

Currently, unlicensed massage practitioners (the indentured workers) within facilities suspected of trafficking may be fined or censured individually. Before penalties can be enforced facilities owners routinely move violators between municipalities and states thereby staying beyond reach of the law knowing enforcement priorities and lack of resources favor them. This has been true in confirmed and charged cases of sexual assault within facilities as well. Facilities owners are blatantly flaunting loopholes in state regulation to continue "their business" of endangering public health and safety. Theirs is a profitable "business".

The value of Licensed Therapeutic Massage is recognized throughout the United States by being the Number One out-of-pocket complementary and alternative health care expense. Increasingly insurers, both public and private, view licensed massage therapy as a viable option physicians can

refer for in preventative, restorative and palliative circumstances for safe and effective outcomes, meeting essential health benefits standards. The passage of SB 387 in Oregon is essential in the maturation of therapeutic massage as a professional partner in state health care policy, and as a valued state licensed health service provider. (Please refer to the Study provided entitled: Evidence-Informed Massage Therapy' The research Supporting Massage Therapy as an Integral Component in the Affordable Health Care Act's Essential Health Benefits, sponsored by the State of Washington American Massage Therapy Association).

Statewide ethical massage facilities owners are ready to distinguish their profession by delineating themselves from illicit facilities through the permit process. They are joined by the coalition who have provided emails of support for SB 387 to your committee, including domestic violence and victims support service providers; community human service providers; social workers; the American Bodywork and Massage Professionals, the trade organization representing the largest number of active Oregon LMTs; the Federation of State Massage Therapy Boards; and the CHP Group, representing the Oregon Chiropractic profession.

Senators, I ask that you help the massage profession to protect the health and safety of Oregonians by supporting SB 387 and move it forward.

Respectfully submitted,

David Fredrickson
Board Chair

Evidenced-Informed Massage Therapy: The Research Supporting Massage Therapy is an Integral Component in the Affordable Care Act's Essential Health Benefits

Project funded by American Massage Therapy Association-WA

Project Coordinators: Marybeth Berney, LMP, President, AMTA-WA

Diana L Thompson, LMP, Past President, Massage Therapy Foundation

Project Authors: Marissa Brooks, MPH, LMP

Michael Hamm, LMP, CCST

Benjamin Erkan, BA

Diana L Thompson, LMP, Past President, Massage Therapy Foundation

Kenneth Pfaff, HFWLS, HPCUHGS, Head Librarian, The Grotto Library

Reviewers: Albert Moraska, PhD

Tracy Walton, MS, LMT

Ruth Werner, LMP

Introduction: Massage Therapy in Health Care

Massage therapy (MT) is defined as the intentional and systematic manipulation of the soft tissues of the body to enhance health and healing (Benjamin & Tappan 2004). MT is a profession with the intention of positively affecting the health and well-being of the client through a variety of touch techniques (American Massage Therapy Association Glossary of Terms). MT's are trained in assessment, treatment planning, application of various massage techniques/modalities, communication/reporting with both patients and members of the health care team (Goldblatt 2009).

MT use is widespread in the US, and is increasingly viewed as part of integrative health care. This is due to a rapidly expanding evidence base (Moyer & Dryden 2012, Moyer et al. 2009) and MT's popularity for treating common health complaints. A 2007 National Institute of Health (NIH) survey of US adults found that MT was the the number one out-of-pocket practitioner-based expense and the second most commonly used practitioner-based CAM therapy -- nearly equalling the volume of chiropractic/osteopathic care (Barnes 2008). A 2009 survey by the American Massage Therapy Association (AMTA Massage Therapy Industry Fact Sheet 2009) found 97% of recipients agreeing that MT should be considered a part of health care.

Studies show that patients frequently do not limit themselves to a single modality of care—they do not see conventional (allopathic) medicine as being mutually exclusive—and this pattern will probably continue and may even expand as evidence of therapies' effectiveness accumulate (IOM 2005). Chiropractic, massage, acupuncture and Asian medicine, midwifery, and naturopathic medicine were in common use before national surveys documented their sizable footprint within the US health care delivery system (Goldblatt 2009).

In the state of Washington, MT has played a role in integrative health care since 1988, with massage therapists delivering care in numerous settings, including private practice, wellness centers, physicians' offices, and hospitals, and receiving reimbursement through the Department of Labor and Industries and third party payers. Nationally, when patients discuss massage with their health care providers, 52% of physicians, 50% of chiropractors, 49% of physical therapists, and 26% of nurses strongly recommend massage (AMTA Massage Therapy Industry Fact Sheet 2012). MT is frequently used in conjunction with other health care professions, such as physical therapy and chiropractic care. (Cherkin et al 2002; Maiers et al 2010), and physicians in WA state prescribe MT for a broad variety of orthopedic, pain, and mental health conditions. Massage therapy codes typically reimbursed in WA include the CPT codes for physical medicine and rehabilitation: 97010, 97112, 97124, 97140 (AMA 2012). With the emphasis on group health care and community programs, this could expand to include: 97150. It is also important to grasp the broad scope that health care providers rely on massage therapists to treat. To best express that, attached is a list of all the diagnostic codes in rank order from one busy medical massage practice, Mayo Therapy Associates, listing the ICD-9 diagnostic codes included on physician and other health care prescriptions for massage therapy services (Appendix A).

The Affordable Care Act (ACA) supports the integration of MT into state-regulated insurance plans, both in its definitions of health care practitioners, and in its definition of Essential Health Benefits (EHBs). Two sections provide for MTs as practitioners:

Section 2706: Non-discrimination with respect to licensed or certified providers acting within their scope,

- Section 3502: Establishing community health teams that include CAM practitioners (ABMP 2010)

Of the 10 EHBs specified in the Affordable Care Act, MT has shown substantial benefit in three primary categories:

“5. Mental health and substance use disorder services, including behavioral health treatment”

“7. Rehabilitative and habilitative services and devices”

“9. Preventive and wellness services and chronic disease management”

This document will summarize the high-quality evidence for MT’s effectiveness in treating medical conditions and populations pertaining to each EHB. These summaries were compiled primarily from systematic reviews, research syntheses, meta-analyses, research texts and larger clinical trials. Excluded were small trials, small pilot studies, case reports, protocol studies, and editorials. The only exceptions are studies cited in a recent scholarly textbook “Massage Therapy: Integrating Research & Practice” (Dryden & Moyer, 2012), which was treated as a research synthesis.

(For the Ecology of Massage Therapy Research in the United States, see Appendix B. For a list of citations for each section, see Appendix C, specific subheadings therein.)

(For a list of Introduction citations see 1-8 in Appendix C).

EHB: “5. Mental health and substance use disorder services, including behavioral health treatment”

- **Anxiety & Depression**

The effect of MT on anxiety has been researched more than any other outcome. MT has been shown to reduce both “State Anxiety” (current level of anxiety) and “Trait Anxiety” (the tendency to become anxious) in both adults and children (Moyer et al 2004, Dryden & Moyer 2012).

Drawing from 21 randomized controlled trials (RCTs) and 1,026 adult subjects, Moyer et al. (2004) found a “small to medium” statistical effect on state anxiety from a single dose of MT. The same single dose appears to have a stronger effect in children, as seen across four RCTs using 81 total pediatric subjects, showing a “medium to large” statistical effect. In children, the single-dose effect also increased in strength with multiple doses, possibly due to increased comfort with the treatment environment and the practitioner administering treatment. The effect of multiples doses of MT on Trait Anxiety also appears to be strong, yielding a “medium to large” statistical effect across seven RCTs that studied 194 total participants (Moyer et al 2004, Dryden & Moyer 2012).

Depression is “consistently and substantially” reduced by multiple doses of MT, as seen in 10 RCTs totalling 249 participants, and showing that recipients of MT had a lower average post-treatment level of depression than 73% of control group participants. This was a “medium” statistical effect (Moyer et al 2004, Dryden & Moyer 2012).

The US Agency for Healthcare Research and Quality (AHRQ) found that in 2007, anxiety and mood disorders accounted for \$36.8 billion in medical spending and affected 11.8% of adults ages 18 and older (Soni 2010). A 2007 study of US health care costs found “Depression/Anxiety” to be the 6th most costly disease from 2003-2007. (Harvard University 2007). The use of MT as an adjunct to counseling and medication may improve outcomes and reduce costs for these widespread and expensive conditions.

(For a list of Anxiety and Depression citations see 22-84 in Appendix C).

- **Sexual Abuse Recovery & Post-Traumatic Stress**

Trauma from sexual abuse, and trauma disorders in general, involve symptoms of dissociation, avoidance behaviors, physical discomfort, and emotional stress (Dryden & Moyer 2012). Clinical research into trauma disorders emphasize the importance of re-connecting with the body (“dissociation reduction”) in creating positive health outcomes (Van der Kolk 2001 and 2006).

Currently only a small number of studies (Field et al. 1997, Price 2005; Price 2006; Price 2007) have investigated MT as a treatment for sexual trauma, and all with women participants. These studies show reductions in anxiety and depression, dissociation, and physical symptoms, and (from high retention rates and qualitative interviews) suggest a high level of satisfaction with MT as a treatment option (Dryden & Moyer 2012).

Briere and Eliot (2003) estimate that the childhood prevalence of sexual trauma is 14% for males and 32% for females. Kessler et al. (1995) report that approximately 1 out of every 10 women experiences sexual assault, and the rate for military women is 1 in 4 (Sadler et al., 2000). The prevalence of PTSD leads to tremendous societal costs, including increased risk of teenage childbearing, marital instability, unemployment, and dropping out of high school or college (Kessler 2000). Medical costs are also “remarkably high”. It is the most costly anxiety disorder to treat (Marciniak et al. 2005), reflecting, in part, the higher utilization of medical services by patients with PTSD (Tagay et al. 2005)” (Dryden & Moyer 2012).

Future research is needed to determine the degree of benefit from MT, and to specify the best dosing and treatment parameters for different populations of trauma sufferers. The nature of sexual abuse recovery and trauma suggest that MT should be prescribed in tandem with other therapies for the safest and most effective treatment.

(For a list of Sexual Abuse Recovery and Post-Traumatic Stress citations see 85-124 in Appendix C)

EHB: “7. Rehabilitative and Habilitative Services”

● **Scars**

Scars are the end product of certain tissue repair, caused by damage from mechanical trauma, surgical incision, burns, chemicals, or electricity. Scar tissue has different composition and behavior than surrounding tissue and can be associated with movement restriction, pain, chronic inflammation, and negative psychological state (Arabi et al. 2007; Edgar and Breeton 2004).

MT has long been used on scar tissue, with the intent of normalizing the tissue quality, mobilizing the scar, and decreasing pain/inflammation in the area. Several mechanisms for MTs effectiveness have been proposed, such as mobilizing/softening collagen, breaking down adhesions with underlying tissue layers, and displacing the fluid that caused the elevated scar. While plausible, these theories have not been rigorously tested (Dryden & Moyer 2012).

Only seven studies have directly investigated the use of MT on scars, and all have focused on burn scars. Four RCTs with 30 patients or fewer found positive effects on pain, pruritis (inflammatory itching) and anxiety/mood (Hernandez-Reif et al 2001; Palatino et al 1999; Field et al 1998; Silverberg et al. 1996). MT’s effect on tissue pliability and vascularity were inconsistent across the studies (Dryden & Moyer 2012).

The applicability of MT’s effectiveness on scars in general is limited, however, the positive effects on pain and anxiety/mood seen in other clinical domains of MT research are reinforced here, and may be an important component of scar treatment.

(For a list of Scar citations see 129-154 in Appendix C)

● **Athletic Injury and Post-exercise Recovery**

MT for several orthopedic and pain conditions is summarized in other sections. However, it is worth addressing the role of MT in athletics in general, given the high and increasing popularity of this intervention (Galloway and Watt 2004). Reasons for seeking MT include pain reduction, improved flexibility, and facilitated recovery. However, many of these claims have yet to be confirmed in this domain (Callahan 1993), and the settings, applications, and intents of MT for athletes is so varied that research has yet to focus on these individually (Dryden & Moyer 2012).

A recent literature review by Howatson and van Someren (2008) states that MT is beneficial in alleviating delayed-onset muscle soreness (DOMS). The mechanism is not confirmed, though some combination of neural, mechanical, and chemical change post-MT has been proposed. Short-term muscle length and flexibility has been shown alterable by MT, but the advantage of increased muscle length on athletic performance and injury recovery depends on the specific demands of the sport and nature of the injury (Dryden & Moyer 2012).

Some limited evidence exists for the improvement of muscle performance post-exercise and MT. In a recent RCT of 52 adults, Brooks et al. (2005) took baseline, post-exercise, and post-intervention measurements of grip strength, and found MT to outperform both placebo and nonintervention in post-fatigue grip strength. The duration of this effect is unclear, and many more activity-specific MT studies are suggested (Brummitt 2008).

(For a list of Athletic Injury and Post-exercise Recovery citations see 155-216 in Appendix C)

- **Post-Operative Recovery**

Over 48 million inpatient surgical procedures are performed annually in the US (Fast Stats: Inpatient Surgery, Centers for Disease Control and Prevention, last updated May 2012). This does not account for less invasive procedures that allow patients to recover at home. Additionally, major surgical interventions are on the rise. Knee replacement surgeries, for example, are expected to increase by nearly 700% between 2005 and 2030 (Kurtz 2007). This increase is anticipated in part due to people living 25% longer and are more physically active, yet 20% heavier than decades ago (Crowninshield 2006).

Inpatient and outpatient surgical patients often experience pain, distress, anxiety, poor sleep quality, nausea and fatigue. Routinely, patients report mild to moderate pain even with the use of pain medications. More importantly, the unpleasantness of the pain persists, even though the level of pain itself is lessened (Mitchinson 2007).

Over a decade of studies have investigated the use of post-operative MT for a range of surgical interventions, including: hematopoietic cell transplant, thoracic and cardiac surgery, mastectomies, cesarean sections and other abdominal surgeries. Subjects include children, women in childbirth, veterans, and older adults (Ackerman 2012, Dion 2011, Nerbass 2010, Cutshell 2010, Mitchinson 2011, Mehling 2007, Kshetry 2006, Chen 2005, Fletcher 2004, Wang 2004, Piotrowski 2003, Taylor 2003).

Two recent studies have used a crossover design to investigate the use of focused MT in a post-operative rat model. Bove & Chapelle (2012a) showed that a blinded massage therapist can lyse and prevent post-operative organ adhesions in a rat model. The subsequent paper (2012b) also showed a reduction in post-operative ileus (digestive sluggishness), which is a widespread and costly ailment of abdominal surgical patients.

While the surgical procedures and the patient populations in these studies vary, the post-operative symptoms are similar and consistently unrelieved through conventional treatment. Pain and its unpleasantness persist, anxiety is prevalent, nausea and poor-quality sleep are commonplace. The evidence consistently points to the positive effects of MT, regardless of the surgical procedure or the population.

(For a list of Post-Operative citations see 217-235 in Appendix C)

EHB: “9. Preventive and Wellness Services and Chronic Disease Management”

- **Headache**

According to the Centers for Disease Control, about 10% of all men and almost 22% of all women experience a migraine or severe headache within a typical three month period (CDC MMWR Weekly December 3, 2010 / 59(47);1557). Headache is an expensive disorder: up to one-third of all neurologists’ consultations are because of headaches – more than any other complaint (World Health Organization Headache disorders Fact sheet N°277 March 2004).

The research on MT as a treatment for headaches is somewhat limited; however “...the available studies show promising results for both tension-type and migraine headaches (Dryden & Moyer, 2012 pg. 120). The research has examined MT as an adjunct or multi-modal treatment option. Research texts have suggested focusing treatment on trigger points for both types of headaches, with specific application to the head, neck and shoulder anatomical regions of the body. The relaxation benefits of massage for the population that lives with headaches have also been noted as a valuable treatment approach. The evidence suggests that both immediate and “long-term benefits” have been observed from a series of MT treatments (Dryden & Moyer, 2012 pg. 122).

(For a list of Headache citations see 509-573 in Appendix C)

- **Neck and Shoulder Pain**

In a review of complementary and alternative medicine (CAM) , Hurwitz et al. (2008) determined that massage therapy, among other CAM modalities were just as effective in treating neck pain as conventional (allopathic) medicine. An evidence-based clinical guideline that reviewed ten studies (Brosseau et al. 2012) concludes: “Therapeutic massage can decrease pain, tenderness, and improve range of motion for sub-acute and chronic neck pain.” Research provides evidence for the short term relief of neck and shoulder pain symptoms. Additionally, research suggests that MT may interrupt inflammatory processes contributing to neck or shoulder pain, and the reduction of anxiety from MT treatment may help with underlying symptoms of muscle tension and pain. “The best available evidence for treatment of neck pain includes...Massage Therapy (Dryden/Moyer, 2012, pg. 133).”

Many multi-study reviews of MT for neck pain have been published recently (Rickards 2006, Harraldson et al 2006, Verhagan 2007), showing modest effects for treating neck pain. Individual studies point to MT’s effectiveness (Sherman 2009, Vassiliou et al. 2006). “Vassiliou and colleagues... found that the MT group significantly outperformed [neck pain improved] ...at the a 6-month follow-up Dryden & Moyer 2012 pg. 131-32.” These positive treatment effects are consistent with the conclusions reached in a Cochrane review by Furlan and colleagues (2009).

(For a list of Neck and Shoulder Pain citations see 574-621 in Appendix C)

- **Low Back Pain**

Americans spend at least \$50 billion each year on low back pain. It is the most common cause of job-related disabilities (NINDS Low Back Pain Fact Sheet, last updated September, 2012). The number one reason people seek CAM therapies including MT is for back pain (Barnes 2007).

A recent Cochrane review (Furlan et al. 2008) reviewed 13 trials of MT for low back pain (LBP). Two of these trials found MT to be more effective than a sham treatment on pain and function. Another had lasting effects at the one year followup. Furlan and colleagues concluded in the 2008 Cochrane review: “massage was superior for pain and function on both short- and long-term followups. Massage was...superior to joint mobilization, relaxation therapy, physical therapy, acupuncture and self care education.” In addition to the clinical effectiveness, MT for LBP has been found to be “a safe therapeutic modality with few risks or adverse effects (Dryden & Moyer 2012, pg. 140).”

The evidence shows that MT provided by trained massage therapists for chronic, subacute and nonspecific LBP will alleviate pain, reduce swelling and increase function (Dryden & Moyer 2012). The American Pain Society and the American College of Physicians (Chou & Huffman 2007) recommend MT as a nonpharmacological treatment for LBP.

(For a list of Low Back Pain citations see 622-674 in Appendix C)

- **Fibromyalgia**

“Fibromyalgia Syndrome (FMS) is characterised by widespread muscle aches and pain lasting more than 3 months.” (Dryden & Moyer, 2012). Many other symptoms and conditions commonly accompany FMS, including difficulty with cognitive function, fatigue, depression/anxiety, and headaches to name a few. Fibromyalgia affects as many as 5 million Americans ages 18 and older. About 80-90 percent of people with fibromyalgia are women (Fibromyalgia Fact Sheet, Office of Women’s Health, U. S. Department of Health and Human Services, last updated June 2010).

Lemestra and Olszynski (2005) found that the massage group had less pain and depressed mood and an improved self-perceived health status, compared to the control group after the 6-week intervention period. Both connective tissue massage and manual lymph drainage improved symptoms, and the majority of these benefits were observed after the 15-month follow-up (Ekici et al. 2009).

Much of the research demonstrating MT as an effective treatment for FMS can be found in the areas of pain (Zautra et al. 2010), and depression/anxiety and central nervous system research (Mataran-Penarrocha et al. 2009).

(For a list of Fibromyalgia citations see 675-756 in Appendix C)

- **Cancer**

Research on cancer and MT has more than 60 studies that provide evidence for the safety and feasibility of massage treatment for both adults and children across the cancer experience spectrum (Dryden & Moyer 2012). Well-known cancer symptoms and side effects of treatments include pain, anxiety, fatigue, depression, sensitivity to touch, nausea, headaches, lightheadedness, muscle tenderness, and general malaise.

The Society for Integrative Oncology recommends massage as a treatment for cancer pain and anxiety, with precautions at certain anatomical sites. Many recommendations come from the research about how to modify the treatment application, intensity and duration. One example is MacDonald (2007) who suggests that working gently, with slow, light strokes for a shorter session time would benefit cancer patients with treatment side effect symptoms.

The strongest and most consistent evidence of symptom reduction is seen in anxiety and pain from single doses of MT given to adults (see anxiety section). Several longitudinal studies also report pain reduction over multiple doses of MT, and a reduction of analgesic use after receiving massage. Similar effects are seen in children, although the pediatric studies are few, and often have low statistical power (Dryden & Moyer 2012).

A 2007 study of health care expenditures estimated that Cancer costs the US between \$62 billion in direct medical costs and up to \$269 billion annually with the inclusion of indirect costs, such as work cessation and household assistance (Dryden & Myer 2012). MT may play a vital role in the reduction of symptoms, general support and in lowering indirect costs in patients with cancer.

Below is a table with recent studies that showed a positive effects of MT on the specified cancer symptom:

Adults with Cancer	
Symptom	Positive Effect of Massage
Anxiety	Ahles et al. 1999; Campeau et al. 2007; Hernandez-Reif et al. 2004; Hernandez-Reif et al. 2005; Jane et al. 2009; Post-White et al. 2005; Quattrin et al. 2006; Stephenson et al. 2007; Tsay et al. 2008; Wilkie et al. 2000; Wilkinson et al. 2008.
Depression/Depressed Mood	Listing et al. 2009; Hernandez-Reif et al. 2004; Hernandez-Reif et al. 2005; Mehling et al. 2007; Post-White 2003; Soden et al. 2004
Pain	Cassileth and Vickers 2004; Grealish Currin & Meister 2008; Hernandez-Reif et al. 2004; Jane et al. 2009; Kutner et al. 2008; Listing et al. 2009; Mehling et al. 2009; Post-White et al. 2003; Smith et al. 2002; Stephenson et al. 2007; Tsay et al. 2008; Wilkie et al. 2000.
Nausea	Billhult, Bergbom & Sterner-Victorin 2007; Grealish, Lomasney & Witeman 2000.

Fatigue	Cassileth & Vickers 2004; Hernandez-Reif et al. 2005; Listing et al. 2009.
---------	--

(Dryden & Moyer 2012)

Below is a table with recent studies that showed a positive effects of MT on the specified symptom:

Children with Cancer	
Symptom	Positive Effect of Massage
Anxiety	Field et al. 2001; Haun, Graham-Pole & Shortely 2009; Phipps et al. 2005; Post-White et al. 2009.
Depression/Depressed Mood	Field et al. 2001; Haun, Graham-Pole & Shortely 2009.

(Dryden & Moyer 2012)

(For a list of Cancer citations see 757-834 in Appendix C)

- **Temporomandibular Joint Disorder**

Temporomandibular joint disorder (TMJD) is a group of conditions that impair the function of the the jaw and the muscles surrounding it, leading to popping, locking, and pain with chewing or speaking. TMJ disorders may affect over 10 million Americans; the conditions appear to be more common in women than men (TMJ (Temporomandibular Joint and Muscle Disorders), National Institute of Dental and Craniofacial Research, last updated January, 2012). Per Washington Administrative Code (WAC) massage practitioners with the intraoral endorsement have within their scope of practice to provide treatment to the intraoral musculature, an anatomical site often associated with TMJD.

In a 2003 survey of TMJD patients, Debar and colleagues (2003) reported MT as the most frequently utilized and helpful treatment application. Kalamir and colleagues (2007) state that “recently has manipulative [soft tissue] therapy been considered by the wider medical field as a standalone treatment, or alternative to otherwise irreversible intra-articular lavage and surgical techniques (pg. 86).” Two studies showed significant success in the treatment of TMJD through combined muscle treatments such as massage, electro-physical therapies (EPT), mobilization, biofeedback and jaw exercise (Michelotti et al., 2000; van der Glas et al., 2000). Further, Nicolakis et al. (2001) reported that TMJD patients treated with “passive and active exercises, manual therapy, postural correction and relaxation techniques resulted in statistically significant improvements in pain, impairment and range of motion within the treatment group.” This group (2002) also reported on a long-term follow-up study that demonstrated continued success.

(For a list of TMDJ citations see 835-842 in Appendix C)

- **Osteoarthritis**

By the year 2020 researchers estimate that more than 50 million Americans will have osteoarthritis (OA) (Elders 2000, Lawrence et al. 1998). Osteoarthritis will be the most prevalent chronic condition in the older adult population (Arden et al. 2006, Felson 2004). According to Kato et al. (2004), despite conventional treatment, OA often progresses and leads to chronic pain and disability. Another concern is the potential toxic effects of drugs used to treat OA (Matchaba et al. 2005, Naesdal & Brown, 2006).

The evidence for treating OA with MT can be found in many areas of research including musculoskeletal conditions (Ernst 2002, Preyde 2000). In a seminal trial specifically studying the therapeutic effects of MT on OA, Perlman and colleagues (2006) found the group that received MT demonstrated significant improvements in the mean (SD) WOMAC pain, stiffness and functional scores and in the visual analog scale of pain assessment, range of motion in degrees and time to walk 50 ft. According to Perlman and colleagues (2006) "The total cost of OA was estimated at \$60 billion in 2004" (pg. 2533) and this figure is sure to increase with the aging baby boomer population.

(For a list of Osteoarthritis citations see 843-857 in Appendix C)

Emerging Evidence with High Clinical Value

The evidence has not yet caught up with clinical experience in many areas, including but not limited to: fall prevention and balance, insomnia, constipation, and dementia in the older adult population, addiction, dosing, cost effectiveness, multiple chronic conditions, self-care compliance, therapeutic environment, and relational aspects of healing. Included in this section's corresponding references are a few smaller studies and related literature that are beginning to address conditions and situations massage therapists face in regularly clinical practice.

For example, ankle flexibility is critical to balance and massage therapy has been shown to be effective in improving range of motion. Many cancer, post operative and studies investigating the older adult population use sleep as a measure. Massage therapy has been effective in improving restful sleep. Agitated behavior is a critical issue in dementia; massage therapy has emerging evidence of calming, reducing repetitive movements and wandering.

New to MT research is the inquiry into contextual effects and the benefits of practitioner-patient relationships and the clinical setting in which care is provided. Dan Cherkin, PhD, commented on the current discrepancy between patient values and measured outcomes:

"Our analysis identified a range of positive outcomes that participants in CAM trials considered important but were not captured by standard quantitative outcome measures. Positive outcome themes included increased optimism and hope, increased ability to relax, positive changes in emotional states, increased body awareness, changes in thinking that increased the ability to cope with back pain, increased sense of well-being, improvement in physical conditions unrelated to back pain, increased energy, increased patient activation, and dramatic improvements in health or well-being" (Hsu et al 2010).

Critical to this project is an understanding of the environment within which this data collection is possible. Massage therapy research, before the national surveys documented the extent of the use of CAM services by the US consumer, was limited. Further, massage therapists were not included in the studies as investigators, consultants, or practitioners providing the MT interventions. The findings of these surveys, especially the landmark study by David Eisenberg in 1993, documented that Americans were scheduling millions of office visits to CAM providers at a cost of tens of billions of dollars, most of which was paid for out of pocket, and confirming that the majority of these patients do not disclose or discuss the CAM use with their primary care providers (Goldblatt 2009). Since then, the number of studies on massage therapy is on the rise, and the study hypotheses address issues that efficacy is practical and meaningful to the practice of MT (effectiveness) in the role of integrative medicine (See Ecology appendix B).

(For a list of Emerging Evidence citations see 988-995 in Appendix C)

Conclusion: Massage Therapy and the Affordable Care Act

Massage therapy, according to the consumer, is an essential health benefit. As stated previously, massage is the number one out-of-pocket practitioner-based expense of all CAM therapies and the second most-used practitioner-based CAM therapy (Barnes 2008). While the evidence is catching up with what the consumer knows and the practitioners address on a daily basis, the insurance industry must decide what is effective, cost effective, safe, and non-invasive. Many health issues are addressed by massage therapy that result not only in improved quality of life, satisfied patients, but in reduced hospital visits, surgeries, and missed work. Massage therapy is a necessary component of health care and an essential health benefit as defined in the Affordable Care Act.

Appendix A - Diagnoses Ranking/Rank Code Description Count* (Mayo Therapy Associates, PLLC), Accessed: 9/10/2012

1. 724.5 Bachache, NOS 2,094
2. 723.1 Cervicalgia (Neck Pain) 1,971
3. 728.85 Spasm of Muscle 1,412
4. 847.0 Strain/Sprain, neck, whiplash injury 1,342
5. 729.1 Myalgia and myositis, unspecified 1,243
6. 724.2 Low Back Pain (lumbago) 663
7. 847.2 Lumbar Strain/Sprain 638
8. 724.1 Pain in thoracic spine 329
9. 739.1 Segmental/somatic dysfunction, cervical region 304
10. 739.2 Segmental Dysfunction, Somatic Dysfunction, Thoracic 257
11. 847.1 Thoracic Strain/Sprain 248
12. 719.41 Pain in Joint - Shoulder region 241
13. 846.0 Strain/Sprain: Lumbosacral Joint/Ligament 198
14. 729.5 Pain:Arm/Leg 150
15. 719.45 HIP PAIN 137
16. 355.0 Piriformis Syndrome 120
17. 724.3 Sciatica, neuralgia or neuritis of sciatic nerve 120
18. 840.8 Trap. strain 111
19. 307.81 Psychalgia - Tension Headaches 94
20. 722.10 Lumbar intervertebral disc w/o myelopathy 70
21. 715.90 CERVICAL DJD 62
22. 724.4 THoracic or lumbosacral Neuritis or radiculitis 61
23. 840.9 ST/SP USP Shoulder/Upper Arm 59
24. 719.49 PAIN IN JOINT MULTIPLE SITES 56
25. 739.3 Lumbar Region- Segmental/Somatic Dysfunction 53
26. 784.0 Pain: Fascial, head, headache 53
27. 728.89 Other disorders of muscle, ligament, fascia (Inc ITB Synd4ro7me)
28. 723.3 Dx 43
29. 739.0 Dx 43
30. 784.92 DX 43
31. 726.10 Rotator cuff 26
32. 739.4 Segmental, Somatic Dysfunction, Sacral 26
33. 346.90 Migraine 25
34. 844.0 Knee sprain/strain 25
35. 721.0 Cervical spondylitis without myelopathy 24
36. 729.4 Laxity of ligament 22
37. 843.8 HAMSTRING STRAIN 22
38. 722.73 Disc disorder 21
39. 782.0 NUMBNESS 17
40. 726.1 Rotator Cuff Syndrome, shoulder 16
41. 847.9 BACK STRAIN 16
42. 874.2 Lumbar strain 16
43. 715.09 Dx 15
44. 719.46 Knee pain 14
45. 847.3 Sprain sacrum 14
46. 850.0 Concussion w/o loss of consciousness 14
47. 719.58 C/T/L JOINT STIFFNESS 13
48. 737.30 SCOLIOSIS 13
49. 844.9 ST/SP of Knee or Leg 13
50. 726.31 Enthesopathy of elbow - medial epicondylitis 11
51. 733.5 OSTEITIS CONDENSANS 10

- 52. 339.12 TENSION HEADACHE 9
- 53. 726.90 Tendonitis, elbow 9
- 54. 646.80 PREGNAN 8
- 55. 524.60 TMJ 7
- 56. 723.5 TORTICOLLIS 7
- 57. 724.79 COCCYDYNIA 7
- 58. 728.71 PLANTAR FACIITIS 7
- 59. 844.2 ACL TEAR TRAUMATIC 7
- 60. 648.73 Bone and joint disorders of back pelvis and lower limbs of 6mother antepartum
- 61. 719.55 Stiffness of joint - hip 6
- 62. 842.10 Sprain of unspecified site of hand 5
- 63. 719.50 Stiffness in Joint - NOS 4
- 64. 737.39 Scoliosis, Other 4
- 65. 722.0 Displacement of cervical intervertebral disc w/o myelopath2y
- 66. 723.4 Brachia neuritis or radiculitis NOS (Cervical) 2
- 67. 839.20 Lumbar Vertebra, closed 2
- 68. V22.1 Pregnancy 2
- 69. 722.93 Disc disorder of; lumbar region 1
- 70. 780.52 INSOMNIA 1
- 71. V23.81 SUPERVISION OF HIGH RISK PREG 1
- 72. XXX1 First no show 1

*(count based on charge primary diagnosis)

Ecology of Massage Therapy Research in the United States

The profession as licensed, designated health care is still relatively in its early decades, and the profession's knowledge leaders and educators have evolved into a substantially funded sector - career seekers experience greater opportunities for formalized education, and therapists can engage in higher-level training; both answering to increasing demands for evidence. This Appendix describes the most recent ecology of Massage Therapy research in the United States as provided by data extracted from PubMed for the years listed on the date listed (using KNALIJ, a visualization tool, see bottom).

The three sections are: 1) Number of Studies listed in PubMed, 2) General Categories of Studies published, and 3) Federal Funding Impact on the profession.

Number of Studies by Year

Number reported falling into the general category of Massage Therapy, 2006 to 2012:	875
Number reported by PubMed relating directly about Massage Therapy/Pain, 2012:	126
Number reported by PubMed relating directly about Massage Therapy, 2011:	233

General Categories

<u>Category, latest published year</u>	<u># of Studies*</u>
Pain Management studies, 2011	41
Physical Therapy tie-in, 2011	40
Health Knowledge, Attitudes, Practices, 2010	30
Complementary Therapies, 2011	27
Quality of Life, 2011	20
Attitude of Health Personnel, 2011	17
Patient Satisfaction, 2011	14
Patient Relations, 2011	14
Attitude to Health, 2010	13
Nurses Role, 2011	12
Exercise Therapy tie-in, 2010	12

* Number of published studies limited to category with 10 or more available via PubMed.

Federal Funding Impact

<u>Federal Department, year</u>	<u># of Studies*</u>
NCCAM, 2011	97
NIMH, 2011	77
NHLBI, 2011	53
NCI, 2011	36
NCCR, 2011	25
NICHD, 2011	22
NIAMS, 2011	20
NIDDK, 2011	19
PHS, 2011	17
AHRQ, 2011	14
NINR, 2011	12

* Number of studies limited to departments funding 12 or more available, as listed by PubMed. Note that these indicate only *funded* studies, and published findings are not necessarily available in PubMed or as yet available. For example, Dr. Perlman's Massage Therapy/Osteoarthritis study funded by NCCAM has a Primary Estimated Completion date of 2015.

Disclaimer: data extraction from such a database has language and MeSH limitations, and this data may include multi-modal studies (massage and moxibustion) or may be missing studies of physical therapists doing soft tissue work allowed in their region/scope that are also a Massage Therapists' scope.

KNALIJ "is an exploration engine for all types of Internet information. KNALIJ is the next generation of knowledge discovery. It provides a dynamic visual interface for understanding and sharing the connections and relationships in large amounts of information." KNALIJ was presented with the Most Innovation Award from the National Institutes of Health in Bethesda in November 2011.

The extraction tool groups the data based on pre-defined sets or user request, and contains such options as a timeline of published data or by sheer number or category.

Appendix C

List of Citations Displaying Evidenced-Informed Massage Therapy

Citations from Introduction

1. ABMP (2010). A comment on the prevention strategy of the national prevention, health promotion, and public health council.
2. Barnes, P., Bloom, B., & Nahin, R. (2008). *Complementary and alternative medicine use among adults and children: United states, 2007*. National Institutes of Health. *Associated Massage*
3. Benjamin, P., & Tappan, F. (2004). *Tappan's handbook of healing massage techniques: Classic, holistic and emerging methods*. (4th ed.). Pearson/Prentice Hall.
4. Cherkin DC, Deyo RA, Sherman KJ, Hart LG, Street JH, Hrbek A, Davis RB, Cramer E, Milliman B, Booker J, Mootz R, Barassi J, Kahn JR, Kaptchuk TJ, & Eisenberg DM. (2002). Characteristics of visits to license acupuncturists, chiropractors, massage therapists, and naturopathic physicians. *J Am Board Fam Pract*, 15(6):463-72
5. Institute of Medicine (IOM). (2005). *Report on complementary and alternative medicine in the us, national academy of sciences*
6. Maiers MJ, Westrom KK, Legendre CG, & Bronfort G. (2010). Integrative care for the management of low back pain: Use of a clinical care pathway. *BMC Health Serv Res*, 10:298
7. Moyer, C., Dryden, T., & Shipwright, S. (2009). Directions and dilemmas in massage therapy research: A workshop report from the 2009 North American research conference on complementary and integrative medicine. *Int J Ther Massage Bodywork*, 2, 15-27.
8. Goldblatt, E., Snider, P., Quinn, S., & Weeks, J. (2009). *Clinicians' and educators' desk reference on the licensed complementary and alternative healthcare professions*. ACCAHC.

Mental health and substance use disorder services, including behavioral health treatment

9. Blackburn, J., Price, C. (2007). Implications of presence in manual therapy. *J Bodyw Mov Thers*, 11, 68-77.
10. Diego, M., Field, T., Sanders, C., & Hernandez-Reif, M. (2004). Massage therapy of moderate and light pressure and vibrator effects on EEG and heart rate. *Int J Neurosci*, 114, 3 1 4 .
11. Ernst, E. (2003). The safety of massage therapy. *Rheumatology*, 42, 1101-6.
12. Field, T., Diego, M., & Hernandez-Reif, M. (2007). Massage therapy research. *Dev Rev*, 27, 75-89.
13. Ford, C. (1993). *Compassionate touch*. New York: Simon and Schuster.
14. Fosha, D. (2000). *The transforming power of affect: A model for accelerated change*. New York: Basic Books .
15. Gowan-Moody, D., & Baskwill, A. The Federation of Massage Therapy Regulatory Authorities of Canada (FOMTRAC), (2005). *Report on policy issues concerning the regulation of massage therapy in canada*
16. Moyer, C. (2009). Between-groups study designs demand between-groups analyses: A response to Hernandez-Reif, Shar-Posner, Baez, Soto, Mendoz, Castillo, Quintero, Perez, and Zhang. *Evid-Based Compl Alt* 6, 49-50.
17. Moyer, C., Rounds, J., & Hannum, J. (2004). A meta-analysis of massage therapy research. *Psych Bull*, 130, 3-18.
18. Moyer, C., Dryden, T., & Shipwright, S. (2009). Directions and dilemmas in massage therapy research: A workshop report from the 2009 North American research conference on complementary and integrative medicine. *Int J Ther Massage Bodywork*, 2, 15-27.
19. Palinkas, L., & Kabongo, M. (2000). The use of complementary and alternative medicine by primary care patients. A SURF*NET study. *J Fam Pract*, 49(12), 112 1-1130.
20. Scherder, E., Bouma, A., & Steen, L. (1998). Effects of peripheral tactile nerve stimulation on affective behavior of patients with probable Alzheimer's disease. *Am J Alzheimers Dis*, 13, 6149.
21. Butttagat, V., Eungpinichpong, W., Kaber, D., Chatchawan, U., & Arayawichanon, P. (2012). Acute effects of traditional thai massage on electroencephalogram in patients with scapulocostal syndrome. *Complement Ther Med*, 20(4), 167-74. PMID:22579427

Anxiety & Depression

22. Abrams, S. (1999). Attention-deficit/hyperactivity disorder children and adolescents benefit from massage therapy. *Dissertation Abstrats International*, 60, 52, 18.
23. American Psychiatric Association. (2000). *Diagnostic and statistical manual of mental disorders: DSM-IV-TR*. Washington, DC: American Psychiatric Publishing.
24. Barlow, D. (2002). *Anxiety and its disorders: The nature and treatment of anxiety and panic*. 2nd ed. New York: Guilford Press.
25. Beck, A., Epstein, N., Brown, G., & Steer, R. (1988). An inventory for measuring clinical anxiety: Psychometric properties. *J Consult Clin Psych*, 56(6), 893-7.
26. Beider, S., & Moyer, C. (2007). Randomized controlled trials of pediatric massage: A review. *Evid-Based Compl Alt*, 4, 23-34.
27. Black, S., Jacques, K., Webber, A., Spurr, K., Carey, E., Hebb, A., & Gilbert, R. (2010). Chair massage for treating anxiety in patients withdrawing from psychoactive drugs. *J Altern Complement Med*, 16(9), 979-87. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/20799900>
28. Brown, T., Campbell, L., Lehman, L., Grisham, J., & Mancill, R. (2001). Current and lifetime comorbidity of the DSM-IV anxiety and mood disorders in a large clinical sample. *J Abnorm Psychol* 110, 585-99.
29. Burton, M., & Hicks, M. (2005). Hurricane Katrina: Preliminary estimates of commercial and public sector damages. Retrieved from www.marshall.edu/cber/research/katrina/Katrina-Estimates.pdf

30. Coelho, H., Boddy, K., & Ernst, E. (2008). Massage therapy for the treatment of depression: A systematic review. *Int J Clin Pract*, 62(2), 325-33.
31. Cohen, J. (1988). *Statistical power analysis for the behavioral sciences*. 2nd ed. Hillsdale, NJ: Erlbaum.
32. Cross-National Collaborative Group. (1992). The changing rate of major depression: Cross-national comparisons. *JAMA*, 268, 3098-105.
33. Derogatis, L. (1983). *SCL-90-R administration, scoring and procedures manual*. Towson, MD: Clinical Psychometric Research.
34. Diego, M., Field, T., Sanders, C., & Hernandez-Reif, M. (2004). Massage therapy of moderate and light pressure and vibrator effects on EEG and heart rate. *Int J Neurosci*, 114, 31-44.
35. Dooley, D., Fielding, J., & Levi, L. (1996). Health and unemployment. *Annu Rev Publ Health*, 17, 449-65.
36. Dropping out of psychotherapy. (2005). *Harvard Mental Health Letter*, 22, 34.
37. Ernst, E. (2003). The safety of massage therapy. *Rheumatology*, 42, 1101-6.
38. Fava, M. (2006). Prospective studies of adverse events related to antidepressant discontinuation. *J Clin Psychiat*, 67, 14-21.
39. Fawcett, J., Scheftner, W., Fogg, L., Clark, D., Young, M., Hedeker, D., & Gibbons, R. (1990). Time related predictors of suicide in a major affective disorder. *Am J Psychiat*, 146: 1189-94
40. Ferrell-Torry, A., & Click, O. (1993). The use of therapeutic massage as a nursing intervention to modify anxiety and the perception of cancer pain. *Cancer Num*, 16, 32-5.
41. Field, T., Hernandez-Reif, M., Diego, M., Schanberg, S., & Kuhn, C. (2005). Cortisol decreases and serotonin and dopamine increase following massage therapy. *Int J Neurosci*, 115(10), 1397-413.
42. Field, T., Diego, M., Cullen, C., Hernandez-Reif, M., Sunshine, W., & Douglas, S. (2002). Fibromyalgia pain and substance p decrease and sleep improves after massage therapy. *J Clin Rheumatol*, 8, 72-6.
43. Field, T., Diego, M., & Hernandez-Reif, M. (2007). Massage therapy research. *Deu Rev*, 27, 75-89.
44. First, M., Spitzer, R., Gibbon, M., & Williams, J. (2002). *Structured clinical interview for DSM-W-KT Axis I disorders, research version, patient edition with psychotic screen (SUID/P W/ PSY SCREEN)*. New York: New York State Psychiatric Institute.
45. Greenberg, P., Sisitsky, T., Kessler, R., Finkelstein, S., Berndt, E., Davidson, J., & Fyer, A. (1999). The economic burden of anxiety disorders in the 1990s. *J Clin Psychiat*, 60, 427-35.
46. Greenberg, P., Kessler, R., Birnbaum, H., Leong, S., Lowe, S., Berglund, P., & Corey-Lisle, P. (2003). The economic burden of depression in the United States: How did it change between 1990 and 2000? *J Clin Psychiat*, 64, 1465-75.
47. Harvard University. (2007). Healthcare delivery: Deconstructing the costs. *Harvard University - Kennedy School Health Care Delivery Policy Program*.
48. Hecht, H., von Zerssen, D., & Wittchen, H. (1990). Anxiety and depression in a community sample: The influence of comorbidity on social functioning. *J Affect Disorders*, 18, 137-44.
49. Hernandez-Reif, M., Field, T., Krasgenor, J., Theakston, H., Hossain, Z., & Burman, I. (2000). High blood pressure and associated symptoms were reduced by massage therapy. *J Bodyw Moo Ther*, 4, 31-8.
50. Hernandez-Reif, M., Field, T., Krasgenor, J., & Theakston, H. (2001). Lower back pain is reduced and range of motion increased after massage therapy. *Int J Neurosci*, 106, 131-45.
51. Hernandez-Reif, M., Field, T., Krasgenor, J., Theakston, H., Hossain, & Burman, I. (2000). High blood pressure and associated symptoms were reduced by massage therapy. *J Bodyw Mov Thers*, 4, 31-8.
52. Hou, W., Chiang, P., Hsu, T., Chiu, S., & Yen, Y. (2010). Treatment effects of massage therapy in depressed people: a meta-analysis. *J Clin Psychiatry*, 71(7), 894-901. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/20361919>
53. Hulme, J., Waterman, H., & Hillier, V. (1999). The effects of foot massage on patients' perception of care following laparoscopic sterilization as day case patients. *J Adv Nun*, 30, 460-8.
54. Ingram, R., & Siegle, G. (2002). *Contemporary methodological issues in the study of depression: Not your father's Qldsmobile*. In *Handbook of depression*, eds. Gotlib, I., & Hammen, C., 86-114. New York: Guilford Press.
55. Kendler, K., Kessler, R., Walters, E., MacLean, C., Neale, M., Heath, A., & Eaves, L. (1995). Stressful life events, genetic liability, and onset of an episode of major depression in women. *Am J Psychiat*, 152, 833-42.
56. Kessler, R., Berglund, P., Demler, O., Jin, R., Merikangas, K., & Walters, E. (2005). Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the national comorbidity survey replication. *Arch Gen Psychiat*, 62, 595-602.
57. Koran, L., Thiernann, M., & Davenport, R. (1996). Quality of life for patients with obsessive-compulsive disorder. *Am J Psychiat*, 153, 783-8.
58. Marciniak, M., Lage, M., Dunayevich, E., Russell, J., Bowman, L., Landboom, R., & Levine, L. (2005). The Cost of Treating Anxiety: The Medical and Demographic Correlates that Impact Total Medical Costs. *Depression and Anxiety*, 2(1), 178-84.
59. Mitte, K., Noack, P., Steil, R., & Hautzinger, H. (2005). A meta-analytic review of the efficacy of drug treatment in generalized anxiety disorder. *J Clin Psychopharm*, 24, 141-50.
60. Moyer, C., Rounds, J., & Hannum, J. (2004). A meta-analysis of massage therapy research. *Psych Bull*, 130, 3-18.
61. Moyer, C., Dryden, T., & Shipwright, S. (2009). Directions and dilemmas in massage therapy research: A workshop report from the 2009 North American research conference on complementary and integrative medicine. *Int J Ther Massage Bodywork*, 2, 15-27.
62. Moyer, C., Seefeldt, L., Mann, E., & Jackley, L. (2011). Does massage therapy reduce cortisol? A comprehensive quantitative review. *J Bodyw Mov Ther*, 15, 3-14.
63. Radloff, L. (1977). The CESD scale: A self-report depression scale for research in the general population. *Appl Psych Meas*, 1, 385-401.

65. Rosenthal, R. (1995). Writing meta-analytic reviews. *Psych Bull*, 118, 183-92.
 66. Rush, A., Madhukar, H., Wisniewski, S., Stewart, J., Nierenberg, A., Thase, M., & Fava, M. (2006). Bupropion-SR, sertraline, or venlafaxine-XR after failure of SSRIs for depression. *New Engl J Med*, 354, 1231-42.
 67. Scherder, E., Bouma, A., & Steen, L. (1998). Effects of peripheral tactile nerve stimulation on affective behavior of patients with probable Alzheimer's disease. *Am J Alzheimers Dis*, 13, 61-9.
 68. Schleifer, S., Keller, S., Bartlett, J., Eckholdt, H., & Delaney, B. (1996). Immunity in young adults with major depressive disorder. *Am J Psychiat*, 153, 477-82.
 69. Seligman, M. (1995). The effectiveness of psychotherapy. *Am Psychol*, 50, 965-74.
 70. Soni A. (2010) Anxiety and mood disorders: Use and expenditures for adults 18 and older, US civilian noninstitutionalized population, 2007. *US Agency for Healthcare Research and Quality*. Statistical brief #303.
 71. Spielberger, C. (1972). *Conceptual and methodological issues in anxiety research*. In *Anxiety: Vol. 2. Current trends in theory and research*, ed. C.D. Spielberger, 481-93. New York: Academic Press.
 72. (1983). *Manual for the state-trait anxiety inventory*. Palo Alto, CA: Consulting Psychologists Press.
 73. Spina, E., & Scordo, M. (2002). Clinically significant drug interactions with antidepressants in the elderly. *Drugs Aging*, 19, 299-320.
 74. Taylor, S. (2006). *Health psychology*. 6th ed. New York: McGraw-Hill.
 75. Thase, M., Buysse, D., Frank, E., Cherry, C., Cornes, C., Mallinger, A., & Kupfer, D. (1997). Which depressed patients will respond to interpersonal psychotherapy? The role of abnormal EEG sleep profiles. *Am J Psychiat*, 154, 502-9.
 76. United States Department of Health and Human Services. (1997). Mental health providers in rural and isolated areas: Final report of the ad hoc rural mental health provider work group. Retrieved from www.mentalhealth.samhsa.gov/publications/allpubs/SMA9&3166/default.asp.
 77. U.S. Preventive Services Task Force. (1996). *Guide to clinical preventive services*. 2nd ed. Retrieved from <http://odphp.osophs.dhhs.gov/pubs/guidecps>.
 78. Unutzer, J., Patrick, D., Marmon, T., Simon, G., & Katon, W. (2002). Depressive symptoms and mortality in a prospective study of 2,558 older adults. *Am J Geriatr Psychiat*, 10, 521-30
 79. van der Kolk, B. (2001). The assessment and treatment of complex PTSD In *Traumatic Stress*, ed. R. Yehuda, American Psychiatric Press. 2006. Clinical implications of neuroscience research in PTSD. *Ann N Y Acad Sci*, 1071, 277-293.
 80. Walton, T. (2005). Medical conditions in massage practice: Intake forms and questions, part I. Retrieved from www.massagetoday.com/mpacms/mt/article.php?id=13221.
 81. Wampold, B. (2001). *The great psychotherapy debate*. Mahwah, NJ: Erlbaum.
 82. World Health Organization. (2007). *International statistical classification of diseases and related health problems: ICD10*. Geneva: WHO.
 83. Yerkes, R., & Dodson, J. (1908). The relation of strength of stimulus to rapidity of habit-formation. *Journal of Comprehensive Neurologic and Psychology*, 18, 459-82.
 84. Zorilla, E., McKay, J., Luborsky, L., & Schmidt, K. (1996). Relation of stressors and depressive symptoms to clinical progression of viral illness. *Am J Psychiat*, 153, 626-3
- Sexual Abuse Recovery & Post Traumatic Stress**
85. (2004). Characteristics of women seeking body-oriented therapy as an adjunct to psychotherapy during recovery from childhood sexual abuse. *J Bodyw Mov Thers*, 8, 35-42.
 86. (2005). Body-oriented therapy in recovery from child sexual abuse: An efficacy study. *Altern Ther Health Med*, 11(5): 4657.
 87. (2006). Body-oriented therapy in sexual abuse recovery: A pilot-test comparison. *J Bodyw Movu Thers*, 10, 58-64.
 88. (2006). Clinical implications of neuroscience research in PTSD. *Ann N Y AcadSci*. 1071: 277-93.
 89. (2007). Dissociation reduction in body therapy during sexual abuse recovery. *Complement Ther Clin Ract*, 13(2) 1: 16-128.
 90. American Psychiatric Association. (2000). *Diagnostic and statistical manual of mental disorders*. Revised 4th ed. Washington: American Psychiatric Association.
 91. Benjamin, B. (1996). Massage and bodywork with survivors of abuse (Part V). *Massage Therapy Journal*, 35(3).
 92. Blackburn, J., & Price, C. (2007). Implications of presence in manual therapy. *J Bodyw Mov Ther*, 11, 68-77.
 93. Briere, J., & Runtz, M. (1993). Childhood sexual abuse: Long-term sequelae and implications for psychological assessment. *J Interpers Violence*, 8(3), 312-330.
 94. Briere, J., & Elliott, I. (2003). Prevalence and psychological sequelae of self-reported childhood physical and sexual abuse in a general population sample of men and women. *Child Abuse Neglect*, 27(10), 1205-22.
 95. Breslau, N. (2002). Gender differences in trauma and post-traumatic stress disorder. *J Gend Specif Med*, 5(1), 34-40.
 96. Ciechanowski, P., Walker, E., Russo, J., Newman, E., & Katon, W. (2004). Adult health status of women HMO members with post-traumatic stress disorder symptoms. *Gen Hosp Psychiatry*, 26(4), 261-8.
 97. Dobie, D., Kivlahan, D., Maynard, C., Bush, K., Davis, T., & Bradley, K. (2004). Post-traumatic stress disorder in female VA patients: Association with self-reported health problems and functional impairment. *Archives of Internal Medicine*, 164(4), 394-400.
 98. Field, T., Hernandez-Reif, M., Hart, S., Quintin, O., Drose, L., Field, T., Kuhn, C., & Schanberg, S. (1997). Effects of sexual abuse are lessened by massage therapy. *J Bodyw Moo Ther*, 1(2), 65-9.
 99. Fitch, D., & Dryden, T. (2000). Recovering body and soul from post-traumatic stress disorder. *Massaqe Therapy Journal*, 39(1), 41-62.
 100. Ford, C. (1993). *Compassionate touch*. New York: Simon and Schuster.

101. Fosha, D. (2000). *The transforming power of affect: A model for accelerated change*. New York: Basic Books.
102. Gill, J., & Page, G. (2006). Psychiatric and physical health ramifications of traumatic events in women. *Issues Ment Health Nurs*, 27(7), 71-84.
103. Herman, J. (1997). *Trauma and recovery: The aftermath of violence-From domestic abuse to political terror*. New York: Harper Collins.
104. Kepner, J. (2003). *Psychotherapy with adult survivors of childhood abuse*. London: Routledge.
105. Kessler, R. (2000). Post-traumatic stress disorder: The burden to the individual and to society. *J Clin Psychiatry*, 61(5), S4-S14.
106. Kessler, R., Sonnega, A., Bromet, E., Hughes, M., & Nelson, C. (1995). Post-traumatic stress disorder in the National Comorbidity Survey. *Arch Gen Psychiatry*, 52(12), 1048-60.
107. Kimerling, R. (2004). An investigation of sex differences in non-psychiatric morbidity associated with post-traumatic stress disorder. *J Am Med Womens Assoc*, 59(1), 43-7.
108. Marciniak, M., Lage, M., Dunayevich, E., Russell, J., Bowman, L., Landboom, R., & Levine, L. (2005). The Cost of Treating Anxiety: The Medical and Demographic Correlates that Impact Total Medical Costs. *Depression and Anxiety*, 21, 178-84.
109. McCauley, I., Kern, E., Kolodner, K., Dill, L., Schroeder, A., DeChant, H., & Bass, E. (1997). Clinical characteristics of women with a history of childhood abuse: Unhealed wounds. *JAMA*, 277(17), 1362-8.
110. Olf, M., Langeland, W., Draijer, N., & Gersons, B. (2007). Gender differences in post-traumatic stress disorder. *Psychol Bull*, 133(2), 183-204.
111. Palinkas, L., & Kabongo, M. (2000). The use of complementary and alternative medicine by primary care patients. A SURF*NET study. *J Fam Pract*, 49(12), 1121-30.
112. Pearlman, L., & Saakvitne, K. (2005). *Trauma and the therapist: Countertransference and vicarious traumatization in psychotherapy with incest survivors*. New York: Norton.
113. Price, C. (2002). Body-oriented therapy as an adjunct to psychotherapy in recovery from childhood abuse: A case study. *J Bodyw Mov Ther*, 6(4), 228-36.
114. Price, C., Wells, E., Donovan, D., & Brooks, M. (2012). Implementation and acceptability of mindful awareness in body oriented therapy in women's substance use disorder treatment. *Journal of Alternative and Complementary Medicine*, 18(5), 1-9. doi: 10.1089/acm.2011.0126
115. Price, C., McBride, B., Hyerle, L., & Kivlahan, D. (2007). Mindful awareness in body-oriented therapy for female veterans with post-traumatic stress disorder taking prescription analgesics for chronic pain: A feasibility study. *Altern TherHealth Med*, 13(6), 32-40.
116. Sadler, A., Booth, B., Nielson, D., & Doebbeling, B. (2000). Health-related consequences of physical and sexual violence: Women in the military. *Obstetrics & Gynecology*, 96(6), 473-80.
117. Seedat, S., Stein, D., & Carey, P. (2005). Post-traumatic stress disorder in women: Epidemiological and treatment issues. *W S Drugs*, 19(5), 41-27.
118. Stein, M.B., Walker, J., & Forde, D. (2000). Gender differences in susceptibility to post-traumatic stress disorder. *Behav Res Ther* 38(6), 619-28.
119. Tagay, S., Herpertz, S., Langkafel, M., & Senf, W. (2005). Post-traumatic stress disorder in a psychosomatic outpatient clinic. Gender effects, psychosocial functioning, sense of coherence, and service utilization. *J Psychosom Res*, 58(5), 439-46.
120. Thayer, J., & Lane, R. (2000). A model of neurovisceral integration in emotion regulation and dysregulation. *J Affect Disord*, 61(3), 201-16.
121. Timms, R., & Connors, P. (1992). *Embodying healing: Integrating bodywork and psychotherapy in recovery from childhood sexual abuse*. Orwell: Safer Society Press.
122. van der Kolk, B. (2001). *The assessment and treatment of complex PTSD*. In *Traumatic Stress*, ed. R. Yehuda, American Psychiatric Press.
123. van der Kolk, B., van der Hart, O., & Marmar, C. (1996). *Dissociation and information processing in post-traumatic stress disorder*. In *Traumatic stress: The effects overwhelming experience on mind, body and society*, ed. 13. van der Kolk, C. McFarlane, and L. Weisaeth. New York: Guilford Press.
124. Weisberg, R., Bruce, S., Machan, J., Kessler, R., Culpepper, L., & Keller, M. (2002). Nonpsychiatric illness among primary care patients with trauma histories and post-traumatic stress disorder. *Psychiatr Serv*, 53(7), 848-54.

Rehabilitative and Habilitative Services

125. Arroyo-Morales, M., Olea, N., Martinez, M., HidalgoLozano, A., Ruiz-Rodriguez, C., Rodriguez, L. (2008). Psychophysiological effects of massage-myofascial release after study. *J Altern Complement Med*, 14, 1223-9.
126. Dishman, J., & Bulbulian, R. (2001). Comparison of effects of spinal manipulation and massage on motorneuron excitability. *Electromyelin C I Neurophysiol*, 41, 97-106.
127. Moyer, C. (2008). Affective massage therapy. *Int J Ther Massage and Bodyw*, 1, 35.
128. Moyer, C., Rounds, J., & Hannum, J. (2004). A meta-analysis of massage therapy research. *Psych Bull*, 130, 3-1

Scars

129. Aarabi, S., Longaker, M., & Gurtner, G. (2007). Hypertrophic scar formation following burns and trauma: New approaches to treatment. *PLoS Med*, 4(9). Retrieved from www.plosmedicine.org/article/info:doi%2F10.1371%2Fjournal.pmed.0040234.

130. Atiyeh, B. (2007). Nonsurgical management of hypertrophic scars: Evidence-based therapies, standard practices and emerging methods. *Aesthet Plast Surg*, 31, 468-92.
131. Baryza M., & Baryza, G. (1995). The Vancouver scar scale: An administrative tool and its interrater reliability. *J Burn Care Rehabil*, 6, 535-8.
132. Bell, P., & Gabriel, V. (2009). Evidence-based review for treatment of post-burn pruritus. *J Burn Care Res*, 30, 55-61.
133. Boersen, K. (2001). Treating post-burn pain and injury: Massage therapy in rehabilitation. Case report. *Rehab and Community Care Medicine Fall*, 56-7.
134. Chapman, T. (2007). Burn scar and contracture management. *J Traum*, 62(6), S8.
135. Demling, R., & DeSanti, L. (2001). Scar management strategies in wound care. *Rehab Management*. Retrieved from www.rehabpub.com/features/892001/3.asp.
136. Edgar, D., & Breerton, M. (2004). ABC of burns-Rehabilitation after burn injury. *Brit Med J*, 329, 343-5.
137. Edwards, J. (2003). Scar management. *Nursing Standard*, 17(52), 39-42.
138. Field, T., Peck, M., Krugman, S., Tuchel, T., Schanberg, S., Kuhn, C., & Burman, I. (1998). Burn injuries benefit from massage therapy. *J Burn Care Rehab*, 19, 241-4.
139. Goutos, I., Dziejewski, P., & Richardson, P. (2009). Pruritus in burns: Review article. *J Burn Care Res*, 30(2), 221-8.
140. Hernandez-Reif, M., Field, T., Largie, S., Hart, S., Redzepi, M., Nierenberg, B., & Peck, M. (2001). Children's distress during burn treatment is reduced by massage therapy. *J Burn Care Rehab*, 22, 191-5.
141. Kokoska, M., & Prendiville, S. (2007). Hypertrophic scarring and keloids. Retrieved from <http://emedicine.medscape.com/article/87621kverview>.
142. LaFrano, C. (2001). Scar tissue massage. *Massage Magazine*, 91(May/June), 151.
143. Ludwig, L. (2000). *Wounds and burns: Injuries that break the skin*. In *Clinical massage therapy: Understanding, assessing and treating over 70 conditions*, eds. F. Rattray and L. Ludwig, 249-63. Toronto, ON, Talus.
144. Melzack, R. (1975). The McGill pain questionnaire: Major properties and scoring methods. *Pain*, 1, 277-99.
145. Morien, A., Garrison, D., & Smith, N. (2008). Range of motion improves after massage in children with burns: A pilot study. *J Bodyw Mov Ther*, 12, 67-71.
146. Mustoe, T., Cotter, R., Gold, M., Hobbs, F., Ramelet, A., Shakespeare, P., & Ziegler, U. (2002). International clinical recommendations on scar management. *Plast Reconstr Surg*, 1(10), 560-71.
147. Patino, C., Novick, C., Mexlo, A., Benaim, F. (1999). Massage in hypertrophic scars. *J Burn Care Rehab*, 10, 268-71.
148. Rorno, T., Pearson, J., Yalamanchili, H., & Zoumalan, R. (2008). Wound healing, skin. Retrieved from <http://emedicine.medscape.com/article/884594-overview>.
149. Sheridan, R. (2007). Burn rehabilitation. Retrieved from <http://emedicine.medscape.com/article/318436-overview>.
150. Silverberg, R., Johnson, J., & Moffat, M. (1996). The effects of soft tissue mobilization on the immature burn scar: Results of a pilot study. *J Burn Care Rehab*, 17(3), 252-9.
151. Tsamis, R. (2005). Massage treatment of a 4 month old, second degree burn. *Journal of Soft Tissue and Manipulation*, 12(4), 12-3.
152. Wihelmi, B. (2008). Wound healing, widened and hypertrophic scars. Retrieved from <http://emedicine.medscape.com/article/1298541-overview>.
153. Wieting, M., & Cugalj, A. (2008). Massage, traction and manipulation. Retrieved from <http://emedicine.medscape.com/article/324694-overview>.
154. World Health Organization. (2010). Framework for action on interprofessional education and collaborative practice. Retrieved from http://whqlibdoc.who.int/hq/2010/WHOHRHJ-IPNN10_33eng.pdf
Athletic Injury and Post-exercise Recovery
155. Ahmaidi, S., Granier, P., Taoutaou, Z., Mercier, J., Dubouchaud, H., & Prefaut, C. (1996). Effects of active recovery on plasma lactate and anaerobic power following repeated intensive exercise. *Med Sci Sports Exerc*, 28, 450-6.
156. Armstrong, R., Warren, G., & Warren, J. (1991). Mechanisms of exercise-induced muscle fibre injury. *Sports Med*, 12, 184-207.
157. Arroyo-Morales, M., Olea, N., Martinez, M., Hidalgo-Lozano, A., Ruiz-Rodriguez, C., & Rodriguez, L. (2008). Psychophysiological effects of massage-myofascial release after exercise: A randomized sham-control study. *J Altern Complement Med*, 14: 1223-9.
158. Barnett, A. (2006). Using recovery modalities between training sessions in elite athletes: Does it help? *Sports Med*, 36, 782-96.
159. Bell, J. (2008). Massage therapy helps to increase range of motion, decrease pain and assist in healing a client with low back pain and sciatica symptoms. *J Bodyw Mov Ther*, 12, 281-9.
160. Berg, H., & Eiken, O. (1999). Muscle control in elite alpine skiing. *Med Sci Sports Exerc*, 31, 1065-7.
161. Bonen, A., Campbell, C., Kirby, R., & Belcastro, A. (1979). A multiple regression model for blood lactate removal in man. *Pflugers Arch*, 380, 205-10.
162. Brooks CP, Woodruff LD, Wright LL, & Donatelli R. (2005) The immediate effects of manual massage on power-grip performance after maximal exercise in healthy adults. *J Altern Complement Med*, 11(6):1093-101.
163. Brummitt, J. (2008). The role of massage in sports performance and rehabilitation: Current evidence and future direction. *N Am J Sports Phys Ther*. 3(1): 7-21
164. Cafarelli, E., & Flint, F. (1992). The role of massage in preparation for and recovery from exercise. An overview. *Sports Med*, 14, 1-9.
165. Callaghan, M. (1993). The role of massage in the management of the athlete: A review. *Br J Sports Med*, 27, 28-33.

166. Cheung, K., Hume, P., & Maxwell, L. (2003). Delayed onset muscle soreness: Treatment strategies and performance factors. *Sports Med*, 33, 145-64.
167. Dishman, J., & Bulbulian, R. (2001). Comparison of effects of spinal manipulation and massage on motorneuron excitability. *Electromyogr Clin Neurophysiol*, 41, 97-106.
168. Dodd, S., Powers, S., Callender, T., & Brooks, E. (1984). Blood lactate disappearance at various intensities of recovery exercise. *J Appl Physiol*, 57, 1462-5.
169. Edman, K., & Reggiani, C. (1984). Redistribution of sarcomere length during isometric contraction of frog muscle fibres and its relation to tension creep. *J Physiol*, 351, 169-98.
170. Emst, E. (1998). Does post-exercise massage treatment reduce delayed onset muscle soreness? A systematic review. *Br J Sports Med*, 32, 212-4.
171. Fowles, J., Sale, D., & MacDougall, J. (2000). Reduced strength after passive stretch of the human plantarflexors. *J Appl Physiol*, 89, 1179-88.
172. Friden, I., Sfakianos, P., & Hargens, A. (1986). Muscle soreness and intramuscular fluid pressure: Comparison between eccentric and concentric load. *J Appl Physiol*, 61, 2175-79.
173. Galloway, S., & Watt, J. (2004). Massage provision by physiotherapists at major athletics events between 1987 and 1998. *Br J Sports Med*, 38, 235-6.
174. Goats, G. (1994). Massage: The scientific basis of an ancient art: Part 2. Physiological and therapeutic effects. *Br J Sports Med*, 28, 153-6.
175. Goldbeg, J., Sullivan, S., & Seaborne, D. (1992). The effect of two intensities of massage on H-reflex amplitude. *Phys Ther*, 72, 449-57.
176. Goldberg, J., Seaborne, D., Sullivan, S., & Leduc, B. (1994). The effect of therapeutic massage on H-reflex amplitude in persons with a spinal cord injury. *Phys Ther*, 74, 728-37.
177. Gordon, A., Huxley, A., & Julian, F. (1966). The variation in isometric tension with sarcomere length in vertebrate muscle fibres. *J Physiol*, 184, 170-92.
178. Gupta, S., Goswami, A., Sadhukhan, A., & Mathur, D. (1996). Comparative study of lactate removal in short term massage of extremities, active recovery and a passive recovery period after supramaximal exercise sessions. *Int J Sports Med*, 17, 106-10.
179. Hemmings, B., Smith, M., Graydon, J., & Dyson, R. (2000). Effects of massage on physiological restoration, perceived recovery, and repeated sports performance. *Br J Sports Med*, 34, 109-14.
180. Hernandez-Reif, M., Field, T., Krasnegor, J., & Theakston, H. (2001). Lower back pain is reduced and range of motion increased after massage therapy. *Int J Neurosci*, 106, 131-45.
181. Hinds, T., McEwan, I., Perkes, J., Dawson, E., Ball, D., & George, K. (2004). Effects of massage on limb and skin blood flow after quadriceps exercise. *Med Sci Sports Exerc*, 36, 1308-13.
182. Hopper, D., Deacon, S., Das, S., Jain, A., Riddell, D., Hall, T., & Briffa, K. (2005). Dynamic soft tissue mobilisation increases hamstring flexibility in healthy male subjects. *Br J Sports Med*, 39, 594-8.
183. Howatson, G., & van Someren, K. (2008). The prevention and treatment of exercise induced muscle damage. *Sports Med*, 38, 483-503.
184. Hunter, A., Smith, L., Watt, J., Yirrell, C., & Galloway, S. (2006). The effect of massage on force production and tensiomyography. *Med Sci Sports Exerc*, 38, S27.
185. Hunter, A., Watt, J., Watt, V., & Galloway, S. (2006). Effect of lower limb massage on electromyography and force production of the knee extensors. *Br J Sports Med*, 40, 114-8.
186. Jones, G. (2007). Massaging the figures. *Cycling Weekly*, August 2. Retrieved from www.cyclingweekly.co-uk
187. Marginson, V., & Eston, R. (2001). The relationship between torque and joint angle during knee extension in boys and men. *J Sports Sci*, 19, 875-80.
188. Martin, N., Zoeller, R., Robertson, R., & Lephart, S. (1998). The comparative effects of sports massage, active recovery, and rest in promoting blood lactate clearance after supramaximal leg exercise. *J Athl Train*, 33, 30-35.
189. Maughan, R., King, D., & Lea, T. (2004). Dietary supplements. *J Sports Sci*, 22, 95-113.
190. Monedero, J., & Donne, B. (2000). Effect of recovery interventions on lactate removal and subsequent performance. *Int J Sports Med*, 21, 593-7.
191. Moraska, A. (2005). Sports massage. A comprehensive review. *J Sports Med Phys Fitness*, 45, 370-80.
192. Morelli, M., Seaborne, D., & Sullivan, S. (1991). H-reflex modulation during manual muscle massage of human triceps surae. *Arch Phys Med Rehabil*, 72, 915-9.
193. Morien, A., Garrison, D., & Smith, N. (2008). Range of motion improves after massage in children with burns: A pilot study. *J Bodyw Mov Ther*, 12, 67-71.
194. Moyer, C. (2008). Affective massage therapy. *Int J Ther Massage and Bodyw*, 1, 35.
195. Moyer, C., Rounds, J., & Hannum, J. (2004). A meta-analysis of massage therapy research. *Psych Bull*, 130, 3-18.
196. Moyer, C., Seefeldt, L., Mann, E., & Jackley, L. (2011). Does massage therapy reduce cortisol? A comprehensive quantitative review. *J Bodyw Mov Ther*, 15, 3-14.
197. NCBTMB. Consumers' massage facts. Retrieved from www.ncbtmb.org/consumers~massage~facts.php.
198. Nelson, A., Guillory, L., Cornwell, C., & Kokkonen, J. (2001). Inhibition of maximal voluntary isokinetic torque production following stretching is velocity-specific. *J Strength Cond Res*, 15, 241-6.

199. Ogai, R., Yamane, M., Matsumoto, T., & Kosaka, M. (2008). Effects of petrissage massage on fatigue and exercise performance following intensive cycle pedalling. *Br J Sports Med*, 42, 534-8.
200. Paschalis, V., Nikolaidis, M., Giakas, G., Jamurtas, A., Pappas, A., & Koutedakis, Y. (2007). The effect of eccentric exercise on position sense and joint reaction angle of the lower limbs. *Muscle Nerve*, 35, 496-503.
201. Rassier, D., Macintosh, B., Herzog, W. (1999). Length dependence of active force production in skeletal muscle. *J Appl Physiol*, 86, 1445-57.
202. Reilly, T., & Ekblom, B. (2005). The use of recovery methods post-exercise. *J Sports Sci*, 23, 619-27.
203. Robertson, A., Watt, J., & Galloway, S. (2004). Effects of leg massage on recovery from high intensity cycling exercise. *Br J Sports Med*, 38, 173-6.
204. Sherman, K., Dixon, M., Thompson, D., & Cherkin, D. (2006). Development of a taxonomy to describe massage treatments for musculoskeletal pain. *BMC Complem Alter M*, 6(24). Retrieved from www.biomedcentral.com/14724882/6/24
205. Shoemaker, J., Tiidus, P., & Mader, R. (1997). Failure of manual massage to alter limb blood flow: Measures by Doppler ultrasound. *Med Sci Sports Exerc*, 29, 610-4.
206. Sullivan, S., Williams, L., Seaborne, D., and Morelli, M. (1991). Effects of massage on alpha motoneuron excitability. *Phys Ther*, 71(8), 555-60.
207. Thomas, C., Perrey, S., Larnbert, K., Hugon, E., Mornet, D., & Mercier, J. (2005). Mono-carboxylate transporters, blood lactate removal after supramaximal exercise, and fatigue indexes in humans. *J Appl Physiol*, 98, 804-9.
208. Tiidus, P. (1999). Massage and ultrasound as therapeutic modalities in exercise induced muscle damage. *Int J Appl Physiol*, 24, 267-78.
209. Tiidus, P., & Shoemaker, J. (1995). Effleurage massage, muscle blood flow and long-term post-exercise strength recovery. *Int J Sports Med*, 16, 478-83.
210. van den Dolder, P., Ferreira, P., & Refshauge, K. (2012). Effectiveness of soft tissue massage and exercise for the treatment of non-specific shoulder pain: A systematic review with meta-analysis. *BR J Sports Med*. Retrieved from <http://www.ncbi.nlm.gov/pubmed/22844035#>
211. van den Dolder, P. A., Ferreira, P. H., & Refshauge, K. M. (2012). Effectiveness of soft tissue massage and exercise for the treatment of non-specific shoulder pain: a systematic review with meta-analysis. *British Journal of Sports Medicine*.
212. Warren, G., Ingalls, C., Lowe, D., & Armstrong, R. (2001). Excitation-contraction uncoupling: Major role in contraction-induced muscle injury. *Exerc Sport Sci Rev*, 29, 82-7.
213. Watt, J. (1999). *Massage for sport*. Wiltshire, England: Crowood Press.
214. Weerapong, P., Hume, P., & Kolt, G. (2005). The mechanisms of massage and effects on performance, muscle recovery and injury prevention. *Sports Med*, 35, 235-56.
215. Weltman, A., Stamford, B., & Fulco, C. (1979). Recovery from maximal effort exercise: Lactate disappearance and subsequent performance. *J Appl Physiol*, 47, 677-82.
216. Wiktorsson-Moller, M., Oberg, B., Ekstrand, J., & Gillquist, J. (1983). Effects of warming up, massage, and stretching on range of motion and muscle strength in the lower extremity. *Am J Sports Med*, 11, 249-52.

Post-Operative Recovery

217. Ackerman, S., Lown, E., Dvorak, C., Dunn, E., Abrams, D., Horn, & Mehling, W. (2012). Massage for Children Undergoing Hematopoietic Cell Transplantation, *A Evid Based Complement Alternat Med*, 792042.
218. Ahles, T., Tope, D., Pinkson, , Walch, S., Hann, D., Whedon, M., & Silberfarb, P. (1999). Massage therapy for patients undergoing autologous bone marrow transplantation. *J Pain Symptom Manag*, 19(3), 157-63.
219. Bauer, B., Cutshall, S., Wentworth, L., Engen, D., Messner, P., Wood C., & Sundt TM 3rd. (2010). Effect of massage therapy on pain, anxiety, and tension after cardiac surgery: A randomized study. *Complement Ther Clin Pract*, 16(2), 70-5. Retrieved from <http://www.ncbi.nlm.gov/pubmed/20347836#>
220. Bove, G & Chappelle, S. (2012a). Visceral mobilization can lyse and prevent peritoneal adhesions in a rat model. *J Bodywork Mvmt Ther*. 16(1), 78-82.
221. Bove, G & Chappelle, S. (2012b). Visceral massage reduces post-operative ileus in a rat model. *J Bodywork Mvmt Ther*. 16(3). Retrieved from <http://www.bodyworkmovementtherapies.com/article/S1360-8592%2812%2900161-1/>
222. "Patients receiving massage therapy had significantly decreased pain, anxiety, and tension. Patients were highly satisfied with the intervention, and no major barriers to implementing massage therapy were identified. Massage therapy may be an important component of the healing experience for patients after cardiovascular surgery."
223. Chen, H., Chang, F., Hsu, C. (2005). Effect of acupressure on nausea, vomiting, anxiety and pain among post-cesarean section women in Taiwan. *Kaohsiung J Med Sci*, 21(8), 341-50.
224. Cutshell, S., Wentworth, L., Engen, D., Sundt, T., Kelly, R., Bauer, B. (2010). Effect of massage therapy on pain, anxiety, and tension in cardiac surgical patients: a pilot study. *Complement Ther Clin Pract*, 16(2), 92-5.
225. Dion, L., Rogers, N., Cutshall, S., Cordes, M., Bauer, B., Cassivi, S., & Cha, S. (2012). Effect of Massage on Pain Management for Thoracic Surgery Patients. *Int J Ther Massage Bodywork*, 4(2), 2-6.
226. Hernandez-Reif, M., Ironson, G., Field, T., Hurley, J., Katz, G., Diego, M., Berman, I. (2004). Breast cancer patients have improved immune and neuroendocrine functions following massage therapy. *J Psychosom Res*, 567(1), 45-52.

227. Hulme, J., Waterman, H., & Hillier, V. (1999). The effects of foot massage on patients' perception of care following laparoscopic sterilization as day case patients. *J Adv Nur*, 30, 460-8.
228. Kshetry, V., Carole, L., Henly, S., Sendelbach, S., Kummer, B. (2006). Complementary alternative medical therapies for heart surgery patients: feasibility, safety, and impact. *Ann Thorac Surg*, 81(1), 201-5.
229. Mehling, W., Jacobs, B., Acree, M., Wilson, L., Bostrom, A., West, J., Acquah, J., Burns, B., Chapman J., Hecht, F. (2007). Symptom management with massage and acupuncture in postoperative cancer patients: a randomized controlled trial. *J Pain Symptom Manag*, 33(3), 258-66.
230. Mitchinson, A., Kim, H., Rosenberg, J., Geisser, M., Kirsh, M., Cikrit, D., Hinshaw, D. (2007). Acute postoperative pain management using massage as an adjuvant therapy: a randomized trial. *Arch Surg*, 142(12), 1158-67.
231. Nerbass, F., Feltrim, M., Souza, S., Ykeda, D., & Lorenzi-Filho, G. (2010). Effects of massage therapy on sleep quality after coronary artery bypass graft surgery. *Clinics (Sao Paulo)*, 65(11), 1105-10. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/21243280>
232. "Massage therapy is an effective technique for improving patient recovery from cardiopulmonary artery bypass graft surgery because it reduces fatigue and improves sleep."
233. Piotrowski, M., Paterson, C., Mitchinson, A., Kim, H., Kirsh, M., Hinshaw, D. (2003). Massage as adjuvant therapy in the management of acute postoperative pain: a preliminary study in men. *J Am Coll Surg*, 197(6), 1037-46.
234. Taylor, A., Galper, D., Taylor, P., Rice, L., Andersen, W., Irvin, W., & Harrell, F. (2003). Effects of adjunctive Swedish massage and vibration therapy on short-term postoperative outcomes: A randomized, controlled trial. *The Journal of Alternative and Complementary Medicine*, 9(1), 77-89.
235. Wang, H., Keck, J. (2004). Foot and hand massage as an intervention for postoperative pain. *Pain Manag Nurs*, 5(2),59-65.
Cancer
236. Ahles, T., Tope, D., Pinkson, B., Walch, S., Hann, D.,Whedon, M., & Silberfarb, P. (1999). Massage therapy for patients undergoing autologous bone marrow transplantation. *J Pain Symptom Manag*, 19(3), 157-63.
237. Anscher MS, Prosnitz LR. Postoperative radiotherapy for patients with carcinoma of the prostate undergoing radical prostatectomy with positive surgical margins, seminal vesicle involvement and/or penetration through the capsule. *J Urol* 1987;138:1407-12
238. Bertelii G, Venturini M, Fome G, et al. An analysis of prognostic factors in response in conservative treatment of post mastectomy lymphedema. *Surg Gynecol Obstet* 1992;175: 455- 460.
239. Breman M. Lymphedema following the surgical treatment of breast cancer: A review of pathophysiology and treatment. *J Pain Symptom Manage* 1992;7:110 -116.
240. Brennan MJ: Lymphedema following the surgical treatment of breast cancer: a review of pathophysiology and treatment. *J Pain Sympt Manage* 7: 110-116, 1992
241. Brennan MJ, DePompolo RW, Garden FH: Focused review: postmastectomy lymphedema. *Arch Phys Med Rehabil* 77: S74-80, 1996
242. Campeau M., Gaboriault, R., Drapeau, M., Nguyen, T., Roy, I., Fortin, B., Marois, M., & Nguyen-Tan, P. (2007). Impact of massage therapy on anxiety levels in patients undergoing radiation therapy: Randomized controlled trial. *JSac Integr Oncol*, 5(4), 133-8.
243. Çınar N, Seçin U, Keskin D, Bodur H, Bozkurt B, Cengiz O. The effectiveness of early rehabilitation in patients with modified radical mastectomy. *Cancer Nurs* 2008;31:160-5.
244. Clarke D, Khonji NI, et al. Sentinel node biopsy in breast cancer. *Ann Surg Oncol* 2004;11(3 Suppl.):208S-2010S.
245. Diego, M., Field, T., Sanders, C., & Hernandez-Reif, M. (2004). Massage therapy of moderate and light pressure and vibrator effects on EEG and heart rate. *Int J Neurosci*, 114(1), 31-44.
246. Faber LP, Jensik RJ, Kittle CF. Results of sleeve lobectomy for bronchogenic carcinoma in 101 patients. *Ann Thorac Surg* 1984;37:279-85. .
247. Fisher B, Redmond C, Fisher ER, et al. Ten year results of randomized clinical trial comparing radical mastectomy and total mastectomy with or without radiation. *N Engl J Med* 1985;312:674-81. .
248. Elder JS, Jewett HJ, Walsh PC. Radical perineal prostatectomy for clinical stage B2 carcinoma of the prostate. *J Urol* 1982; 127:704-6. .
249. Galland, C, JF Auvert, A Flahault, et al: Why and how post-mastectomy edema should be quantified in patients with breast cancer. *Breast Cancer Res. Treat.* 75 (2002), 87-89.
250. Hutzschenreuter P, Wittlinger H, Wittlinger G, Kurz I 1991 Post mastectomyarm lymphedema: treated by manual lymph drainage and compression bandage therapy. *European Journal of Physical Medicine and Rehabilitation* 1: 6
251. Johansson K, Albertsson M, Ingvar C, Ekdahl C: Effects of compression bandaging with or without manual lymph drainage treatment in patients with postoperative arm lymphedema. *Lymphology* 32: 103-10, 1999
252. Kang JC, Wanek LA, Essner R, et al. Sentinel lymphadenectomy does not increase the incidence of in-transit metastases in primary melanoma. *J Clin Oncol* 2005;23:4764-70.
253. Kilbreath SL, Refshauge KM, Beith JM, MiJoung L. Resistance and stretching shoulder exercises early following axillary surgery for breast cancer. *Rehabil Oncol* 2006;24:9-14.
254. Liu N, Olszewski W 1992 Use of tonometry to assess lower extremity lymphoedema. *Lymphology* 25: 155-158
255. Megens, AM, SR Harris, C Kim-Sing, et al: Measurement of upper extremity volume in women after axillary dissection for breast cancer. *Arch. Phys. Med. Rehabil.* 82 (2001), 1639-1644.
256. Mehling, W., Jacobs, Acree, M., Wilson, L., Bostrom, A., West, J., & Hecht, E. (2007). Symptom management with massage and acupuncture in postoperative cancer patients: A randomized controlled trial. *J Pain Sympt Manag*, 33(3), 258-66.

257. Meric, F, TA Buchholz, NQ Mirza, et al: Long-term complications associated with breast-conservation surgery and radiotherapy. *Ann. Surg. Oncol.* 9 (2002), 543-549.
258. Mirolo BR, Bunce IH, Chapman M, Olsen T, Eliadis P, Hennessy JM, Ward LC, Jones LC: Psychosocial benefits of postmastectomy lymphedema therapy. *Cancer Nurs* 18: 197–205, 1995
259. Morrow CP, Bundy BN, Kurman RJ, et al. Relationship between surgical-pathological risk factors and outcome in clinical stage I and II carcinoma of the endometrium: a Gynecologic Oncology Group study. *Gynecol Oncol* 1991;40:55–65. .
260. Moyer, A: Psychosocial outcomes of breast- conserving surgery versus mastectomy: A meta-analytic review. *Health Psychol.* 16(1997), 284-298.
261. Na Y, Lee J, Park J, Kang S, Lee H, Koo J. Early rehabilitation program in postmastectomy patients. *Yonsei Med J* 1999;40:1-8.
262. Phipps, S., Dunavant, M., Rai, S., Deng, X., & Lensing, S. (2004). The effects of massage in children undergoing bone marrow transplant. *Mssuge Ther J*, 43(3), 62-71.
263. Pruthi, S., A.C. Degnim, B.A. Bauer, R.W. DePompolo, and V. Nayar. 2009, Value of massage therapy for patients in a breast clinic. *Clin J Oncol Num* 13(4): 422-425.
264. Quattrin, R., Zanini, A., Buchini, S., Turello, D., Annunziata, M., Vidotti, C., Colambatti, A., & Brusaferrero, S. (2006). Use of reflexology foot massage to reduce anxiety in hospitalized cancer patients in chemotherapy treatment: Methodology and outcomes. *J Nurs Manag*, 14(2), 96-105.
265. Rietman JS, Dijkstra PU, Geertzen JHB, et al. Treatment-related upper limb morbidity 1 year after sentinel lymph node biopsy or axillary lymph node dissection for stage I or II breast cancer. *Ann Surg Oncol* 2004;11:1018–1024.
266. Sartori F, Binda R, Spreafic G, et al. Sleeve lobectomy in the treatment of bronchogenic carcinoma. *Int Surg* 1986;71:233–6.
267. Shin, Y., Kim, T., Shin, M., & Juon, H. (2004). Effect of acupressure on nausea and vomiting during chemotherapy cycle for Korean postoperative stomach cancer patients. *Cancer Nursing*, 27(4), 267-274.
268. Smith, M., Kemp, J., Hemphill, L., & Vojir, C. (2002). Outcomes of therapeutic massage for hospitalized cancer patients. *J Nurs Scholarship*, 34, 257-62.
269. Stillwell GK: Treatment of postmastectomy lymphedema. *Mod Treat* 6: 396–412, 1969
270. Stringer, J., Swindell, R., & Dennis, M. (2008). Massage in patients undergoing intensive chemotherapy reduces serum cortisol and prolactin. *Psych Oncoi*, 17, 1024-31.
271. Tracey GD, Reeve TS, Fitzsimons E, Rundle FF: Observations on the swollen arm after radical mastectomy. *Aust N Z J Surg* 30: 204, 1961
272. Tsay, S., Chen, H., Chen, S., En, I., & Lin, K. (2008). Effects of reflexotherapy on acute postoperative pain and anxiety among patients with digestive cancer. *Cancer Nursing*, 31(2), 109-15.
273. Walsh PC. Adjuvant radiotherapy after radical prostatectomy: is it indicated?. *J Urol* 1987;138:1427–8.
274. Zaino RJ, Kurman RJ, Diana KL, Morrow CP. Pathologic models to predict outcome for women with endometrial adenocarcinoma: the importance of the distinction between surgical stage and clinical stage—a Gynecologic Oncology Group study. *Cancer* 1996;77:1115–21. .
- Cardiac Autonomic Tone**
275. Benson, H., Beary, J. F., & Carol, M. P. (1974). The relaxation response. *Psychiatry*, 37, 37–46.
276. Carney, R. M., Saunders, R. D., Freedland, K. E., Stein, P., Rich, M. W., & Jaffe, A. S. (1995). Association of depression with reduced heart rate variability in coronary artery disease. *The American Journal of Cardiology*, 76(8), 562–564.
277. Delaney, J. P., Leong, K. S., Watkins, A., & Brodie, D. (2002). The short-term effects of myofascial trigger point massage therapy on cardiac autonomic tone in healthy subjects. *Journal of Advanced Nursing*, 37(4), 364–371.
278. Ebner, M. (1985). Connective Tissue Manipulation: Theory and Therapeutic Application. Florida: Krieger Pub Co. Field, T. M. (1998). Massage therapy effects. *The American Psychologist*, 53(12), 1270–1281.
279. Field, T., Ironson, G., Scafidi, F., Nawrocki, T., Goncalves, A., Burman, I. et al. (1996). Massage therapy reduces anxiety and enhances EEG pattern of alertness and math computations. *The International Journal of Neuroscience*, 86(3–4), 197–205.
280. Fischer, A. A. (1987). Pressure algometry over normal muscles. Standard values, validity, and reproducibility of pressure threshold. *Pain*, 30, 115–126.
281. Fischer, A. A. (1988). Documentation of myofascial trigger points. *Archives of Physical Medicine and Rehabilitation*, 69(4), 286–291.
282. Foldi, M., & Foldi, E. (1993). *Lymphoedema: Methods of Treatment and Control*. Australia: Lymphoedema Association of Victoria.
283. Goldstein, D. S., Robertson, D., Esler, M., Straus, S. E., & Eisenhofer, G. (2002). Dysautonomias: Clinical disorders of the autonomic nervous system. *Annals of Internal Medicine*, 137(9), 753–763.
284. Harker, E., Egekvist, H., & Bjerring, P. (2000). Effect of sensory stimulation (acupuncture) on sympathetic and parasympathetic activities in healthy subjects. *Journal of the Autonomic Nervous System*, 79(1), 52–59.
285. Kasseroller, R. G. (1998). The Vodder school: The Vodder method. *Cancer*, 83, 2840–2842.
286. Kay, S. M. (1987). *Modern Spectral Estimation: Theory and Application*. N.J.: Prentice-Hall Englewood Cliffs.
287. Kleiger, R. E., Miller, J. P., Bigger, J. T., Jr., & Moss, A. J. (1987). Decreased heart rate variability and its association with increased mortality after acute myocardial infarction. *The American Journal of Cardiology*, 59(4), 256–262.
288. Kuriyama, H., Watanabe, S., Nakaya, T., Shigemori, I., Kita, M., Yoshida, N. et al. (2005). Immunological and psychological benefits of aromatherapy massage. *Evidence-Based Complementary and Alternative Medicine*, 2(2), 179–184.

289. Lee, M. S., Kim, H. J., Song, J., Park, K. W., & Moon, S. R. (2004). Effects of multifunctional fabrics on cardiac autonomic tone and psychological state. *The International Journal of Neuroscience*, *114*(8), 923–931.
290. Lombardi, F., Sandrone, G., Pernpruner, S., Sala, R., Garimoldi, M., Cerutti, S. et al. (1987). Heart rate variability as an index of sympathovagal interaction after acute myocardial infarction. *The American Journal of Cardiology*, *60*(16), 1239–1245.
291. Madeleine, P., Farina, D., Merletti, R., & Arendt-Nielsen, L. (2000). Upper trapezius muscle mechanomyographic and electromyographic activity in humans during low force fatigue and non-fatiguing contractions. *European Journal of Applied Physiology*, *87*(4–5), 327–336.
292. Maeda, K. (1977). Occupational cervicobrachial disorder and its causative factors. *Journal of Human Ergology*, *6*(2), 193–202.
293. Malliani, A., Pagani, M., Lombardi, F., & Cerutti, S. (1991). Cardiovascular neural regulation explored in the frequency domain. *Circulation*, *84*(2), 482–492.
294. McCraty, R., Atkinson, M., Tiller, W. A., Rein, G., & Watkins, A. D. (1995). The effects of emotions on short-term power spectrum analysis of heart rate variability. *The American Journal of Cardiology*, *76*(14), 1089–1093.
295. Pagani, M., Lombardi, F., Guzzetti, S., Rimoldi, O., Furlan, R., Pizzinelli, P. et al. (1986). Power spectral analysis of heart rate and arterial pressure variabilities as a marker of sympatho-vagal interaction in man and conscious dog. *Circulation Research*, *59*(2), 178–193.
296. Pagani, M., & Malliani, A. (2000). Interpreting oscillations of muscle sympathetic nerve activity and heart rate variability. *Journal of Hypertension*, *18*(12), 1709–1719.
297. Pagani, M., Malfatto, G., Pierini, S., Casati, R., Masu, A. M., Poli, M. et al. (1988). Spectral analysis of heart rate variability in the assessment of autonomic diabetic neuropathy. *Journal of the Autonomic Nervous System*, *23*(2), 143–153.
298. Pomeranz, B., Macaulay, R. J., Caudill, M. A., Kutz, I., Adam, D., Gordon, D. et al. (1985). Assessment of autonomic function in humans by heart rate spectral analysis. *The American Journal of Physiology*, *248*(1 Pt 2), H151–H153.
299. Rechlin, T., Weis, M., Spitzer, A., & Kaschka, W. P. (1994). Are affective disorders associated with alterations of heart rate variability? *Journal of Affective Disorders*, *32*(4), 271–275.
300. Schnoz, M., Laubli, T., & Krueger, H. (2000). Co-activation of the trapezius and upper arm muscle with finger tapping at different rates and trunk postures. *European Journal of Applied Physiology*, *83*(2–3), 207–214.
301. Schumacher, A. (2004). Linear and nonlinear approaches to the analysis of R-R interval variability. *Biological Research for Nursing*, *5*(3), 2115–221.
302. Stein, P. K., Bosner, M. S., Kleiger, R. E., & Conger, B. M. (1994). Heart rate variability: A measure of cardiac autonomic tone. *American Heart Journal*, *127*(5), 1376–1381.
303. Selye, H. (1978). *The Stress of Life*. New York: McGraw-Hill.
304. Sztajzel, J. (2004). Heart rate variability: A noninvasive electrocardiographic method to measure the autonomic nervous system. *Swiss Medical Weekly*, *134*(35–36), 514–522.
305. Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. (1996). Heart rate variability: standards of measurement, physiological interpretation, and clinical use. *Circulation*, *93*, 1043–1065.
306. Wall, P. D., & Melzack, R. (1994). *Textbook of Pain*. Edinburgh: Churchill Livingstone.
307. Zefferino, R., L'Abbate, N., Facciorusso, A., Potenza, A., Lasalvia, M., Nuzzaco, A. et al. (2003). Assessment of heart rate variability (HRV) as a stress index in an emergency team of urban police. *Giornale italiano di medicina del lavoro ed ergonomia*, *25*, 167–169.
308. Zhong, Y., Jan, K. M., Ju, K. H., & Chon, K. H. (2006). Quantifying cardiac sympathetic and parasympathetic nervous activities using principal dynamic modes analysis of heart rate variability. *American Journal of Physiology. Heart and Circulatory Physiology*, *291*(3), H1475–H1483.
- Lymphedema**
309. Andersen L., Hojris I., Erlandsen M., & Andersen J. (2000). Treatment of breast-cancer-related lymphedema with or without manual lymphatic drainage: A randomized study. *Acta Oncol*, *39*, 399–405.
310. Anderson L., Hojris I., Mogens E., & Jorn A. (1999). Treatment of the breast cancer-related lymphedema with or without MLD: A randomized study. *Acta Oncol*, *88*, 2832–7.
311. Armer J., & Stewart B. (2005). A comparison of four diagnostic criteria for lymphedema in a post-breast cancer population. *Lymphat Res Biol*, *3*, 208–17.
312. Arsenault, K., Rielly, L., Wise, H. et al. (2011). Effects of complete decongestive therapy on the incidence rate of hospitalization for the management of recurrent cellulitis in adults with lymphedema. *Rehabilitation Oncology*, *29*(3), 14–20.
313. Badger C., Peacock J., & Mortimer P. (2000). A randomized, controlled, parallel-group clinical trial comparing multilayer bandaging followed by hosiery versus hosiery alone in the treatment of patients with lymphedema of the limb. *Cancer*, *88*:2832–7.
314. Bani H., Fasching P., Lux M., Rauh C., Willner M., Eder I., et al. (2007). Lymphedema in breast cancer survivors: assessment and information provision in a specialized breast unit. *Patient Educ Couns*, *66*:311–8.
315. Beesley V., Janda M., Eakin E., Obermair A., & Battistutta D. (2007). Lymphedema after gynecological cancer treatment: prevalence, correlates, and supportive care needs. *Cancer*, *109*(12), 2607–14.
316. Beaulac S., McNair L., Scout T., LaMorte W., & Kvanah M. (2002). Lymphedema and quality of life in survivors of early stage of breast cancer. *Arch Surg*, *137*:1253–7.
317. Berlin E., Gjores J., Ivarsson C., Palmqvist I., Thagg G., & Thulesius O. (1999). Postmastectomy lymphoedema. Treatment and a five-year follow-up study. *Int Angiol*, *18*:294–8.

318. Boris M., Weindorf S., & Lasinski B. (1997). Persistence of lymph-edema reduction after noninvasive complex lymphedema therapy. *Oncology*, 11:98–109.
319. Boris, M., Weindorf, S., & Lasinski, B. (1994). Lymphedema reduction by noninvasive complex lymphedema therapy. *Oncology*, 8, 95-106.
320. Boris, M., Weindorf, S., & Lasinski, B. (1997). Persistence of lymphedema reduction after non-invasive complex lymphedema therapy. *Oncology*, 11, 99-114.
321. Box, R., Reul-Hirche, H., Bullock-Saxton, J., & Furnival, C. (2002). Physiotherapy after breast cancer surgery: results of a randomised controlled study to minimise lymphedema. *Br Cancer Res Treat*, 75:51-64.
322. Brennan, M., Weitz, J. (1992). Lymphedema 30 years after radical mastectomy. *Am J Phys Med Rehabil*, 71, 12-4.
323. Brennan, M., & Miller, L. (1998). Overview of treatment options and review of the current role and use of compression garments, intermittent pumps, and exercise in the management of lymphedema. *Cancer*, 83, 2821–7
324. Bunce, I., Mirolo, B., Hennessy, et al. (1994). Post-mastectomy lymphoedema treatment and measurement. *Med J Aust*, 16, 125-8.
325. Burt, J., & White, G. (1999). Lymphedema: a breast cancer patient's guide to prevention and healing. Alameda (CA): Hunter House Inc.
326. Carter, B. (1997). Women's experiences of lymphedema. *Oncol. Nurs. Forum* 24, 875-82.
327. Cartier, C., Guilhem, J., & Andrieu, R. (1985). Lymphedema treatment using hydrostatic pressure of mercury. *Prog Lymphol*, 10, 173-5.
328. Casley-Smith J., & Casley-Smith J. (1997). Modern treatment for lymphedema. 5th ed. Adelaide: The Lymphedema Association of Australia, Inc.; 1997. p. 131–58.
329. Casley-Smith, J. (1995). Alterations of untreated lymphedema and its grades over time. *Lymphology*, 28, 174–85
330. Casley-Smith, J., Boris, M., Weindorf, S., & Lasinski B. (1998). Treatment for lymphedema of the arm—the Casley-Smith method. *Cancer*, 83, 2843–60
331. Casley-Smith JR: Modern treatment of lymphoedema. I. Complex physical therapy: the first 200 Australian limbs. *Australas J Dermatol* 33: 61–68, 1992
332. Casley-Smith Judith R, Casley-Smith JR. Modern treatment for lymphedema. 5th ed. Adelaide: The Lymphedema Association of Australia, Inc.; 1997. p. 228.
333. Casley-Smith JR. Alteration of untreated lymphoedema and its grades over time. *Lymphology* 1995;28:174–85.
334. Casley-Smith JR, Casley-Smith JR 1997 Lymphoedema—A Guide for Therapists and Patients. Lymphoedema Association of Australia, Adelaide
335. Chen YW, Tsai HJ, Hung HC, Tsao JY. Reliability study of measurements for lymphedema in breast cancer patients. *Am J Phys Med Rehabil* 2008;87:33-8.
336. Casley-Smith, JR, JR Casley-Smith: Modern treatment of lymphoedema. complex physical therapy: The first 200 Australian limbs. *Australas. J. Dermatol.* 33 (1992), 61-68.
337. Cha K et al. 1995 Multi frequency bioelectrical impedance estimates the distribution of body water. *Journal of Applied Physiology* 79: 1319
338. Cheville A, Tchou J. Barriers to rehabilitation following surgery for primary breast cancer. *J Surg Oncol* 2007;95:409-18.
339. Cheville AL, McGarvey C, Petreck JA, Russo SA, Hiadens SRJ, Taylor ME. The grading of lymphedema in oncology clinical trials *Semin Radiat Oncol* 2003;13:214-25.
340. Chikly, B. (2001). *Theory of Practice of Lymph Drainage Therapy*. Arizona: International Health & Healing.
341. Clark B, J Sitzia, W Harlow: Incidence and risk of arm oedema following treatment for breast cancer: a three-year follow-up study. *Q. J. Med.* 98 (2005), 343-348.
342. Coen J, Taghian A, Kachnic L, Assaad S, Powel S. Risk of lymphedema after regional nodal irradiation with breast conservation therapy. *Int J Radiat Oncol Biol Phys* 2003;55:1209-15.
343. Cornish BH, Chapman M, Hirst C et al. 2001 Early diagnosis of lymphoedema using multifrequency bioimpedance. *Lymphology* 34: 2–11
344. Cox NH. Oedema as a risk factor for multiple episodes of cellulitis/erysipelas of the lower leg: a series with community follow-up. *Br J Dermatol.* 2006;155:947-950.
345. Department of Physical Therapy (S-JK), Yongdong University, Chungbuk and Department of Physical Therapy (C-HY,O-YK), College of Health Science, Yonsei University, Wonju, Kangwon-do, Republic of Korea. , , , & , (2007). Effect of complex decongestive therapy on edema and the quality of life in breast cancer patients with unilateral lymphedema. *Lymphology*, 40, 143-51.
346. Devoogdt, N., Van Kampen, M., Geraerts, I., Coremans, T., & Christiaens, M. (2010). Different physical treatment modalities for lymphoedema developing after axillary lymph node dissection for breast cancer: A review. *European Journal of Obstetrics & Gynecology and Reproductive Biology*, 149, 3-9. doi: 10.1016/j.ejogrb.2009.11.016
347. Didem K, Ufuk YS, Serdar S, Zumre A. The comparison of two different physiotherapy methods in treatment of lymphedema after breast surgery. *Breast Cancer Res Treat* 2005;93:49–54.
348. Dini D, Del Mastro L, Gozza A, et al. The role of pneumatic compression in the treatment of postmastectomy lymphedema. A randomized phase III study. *Ann Oncol* 1998;9:187–90.
349. Ekici, G., Bakar, Y., Akbayrak, T., & Yuksel, I. (2009). Comparison of manual lymph drainage therapy and connective tissue massage in women with fibromyalgia: A randomized controlled trial. *Journal of Manipulative and Physiological Therapeutics*, 32(2), 127-33. doi: 10.1016/j.jmpt.2008.12.001

350. Engler HS, Sweat RD: Volumetric arm measurements: technique and results. *Am Surg* 28: 465–468, 1962
351. Erickson VS, Pearson ML, Ganz PA, Adams J, Kahn KL. Arm edema in breast cancer patients. *J Natl Cancer Inst* 2001;93:96–111.
352. Fernandez JC, Laroche JP, Serin D, Felix-Faure C, Vinot JM. Lymphoscintigraphic aspects of the effects of manual lymphatic drainage. *J Mal Vasc* 1996;21:283–9.
353. Foeldi M. (1994). Treatment of lymphedema. *Lymphology*, 23:1–5.
354. Földi E, Földi M. The lymphoedema chaos: a lancet. *Ann Plast Surg* 1989;22:505-15.
355. Foldi, E: Treatment of lymphedema and patient rehabilitation. *Anticancer Res.* 18 (1998), 2211-2212.
356. Foldi M, Foldi E, Dubik S. Textbook of lymphology for physicians and lymphedema therapists. Munchen: Urban & Fischer; 2003. .
357. Foldi E. The treatment of lymphedema. *Cancer* 1998;83:2833–4.
358. Forner-Cordero I, Munoz-Langa J, Forner-Cordero A, DeMiguel-Jimeno JM. (2010). Predictive factors of response to decongestive therapy in patients with breast-cancer-related lymphedema. *Ann Surg Oncol*, 17(3):744–51
359. Godal R, Swedborg I: A correction for the natural asymmetry of the arms in the determination of the volume of oedema. *Scand J Rehabil Med* 14: 193–195, 1982
360. Godette, K., Mondy, T., & Johnstone, P. (2006). Can manual treatment of lymphedema promote metastasis?. *Journal of the Society for Integrative Oncology*, 4(1), 8-12.
361. Hack TF, Coen L, Katz J, Robson P, Goss P. Physical and psychological morbidity after axillary lymph node dissection for breast cancer. *J Clin Oncol* 1999;17:143-9.
362. Hamner JB, Fleming MD. Lymphedema therapy reduces the volume of edema and pain in patients with breast cancer. *Ann Surg Oncol.* 2007;14:1904-1908.
363. Hampton SE. Ivarsen compression garments in the management of lymphedema. *Br J Nurs* 2003;12:925–929.
364. Harris R 1994 Edema and its treatment in massage therapy. *Journal of Soft Tissue Manipulation* 1: 4
365. Harris R 1992 The Vodder method, an introduction to manual lymph drainage. *Massage Therapy Journal* 31: 1
366. Harris S, Hugi M, Olivotto I, Levine M. Clinical practice guidelines for the care and treatment of breast cancer: Lymphedema. *Can Med Assoc J* 2001;164:191-9.
367. Harris, R., & Piller, N. (2003). Three case studies indicating the effectiveness of manual lymph drainage on patients with primary and secondary lymphedema using objective measuring tools. *J Bodyw Mov Ther*, 7(4), 213-21. doi: 10.1016/S1360-8592(03)00036-6
368. Harris SR, Hugi MR, Olivotto RA, Levine M. Clinical practice guidelines for the care and treatment of breast cancer: lymphedema. *Can Med Assoc J.* 2001;164(2):191-199.
369. Hayes S, Cornish B, Newman B. Comparison of methods to diagnose lymphoedema among breast cancer survivors: 6-month follow-up. *Breast Cancer Res Treat* 2005;89:221-6.
370. Herd-Smith A, Russo A, Muraca MG, et al. Prognostic factors for lymphedema after primary treatment of breast carcinoma 2001. *Cancer* 1992;7:1788 –1797.
371. Hinrichs CS, Gibbs JF, Driscoll D, Kepner JL, Wilkinson NW, Edge SB, et al. The effectiveness of complete decongestive physiotherapy for the treatment of lymphedema following groin dissection for melanoma. *J Surg Oncol* 2004;85(4):187–92.
372. Hornsby R. The use of compression to treat lymphoedema. *Prof Nurse* 1995; 11:127–8.
373. Hutzschenreuter P, Herpertz U 1993 Primary and secondary lymphedema in children treated with manual lymph drainage and compression therapy. *European Journal of Lymphology* 4: 14
374. Hutzschenreuter P, Brummer H, Ebberfeld K, et al. Experimental and clinical studies of the mechanism of effect of MLD therapy. *J Lymphol* 1989;13:62– 64.
375. International Society of Lymphology, author The diagnosis and treatment of peripheral lymphedema. Consensus Document of the International Society of Lymphology. *Lymphology* 2003;36:84–91.
376. International Society of Lymphology. (2009)The diagnosis and treatment of peripheral lymphedema. 2009 consensus document of the International Society of Lymphology. *Lymphology*, 42:51–60.
377. Jager, G, W Doller, R Roth: Quality-of-life and body image impairments in patients with lymphedema. *Lymphology* 39 (2006), 193-200.
378. Jimenez Cossio JA, Farrajota A, Samaniego E, et al. editors. Proceedings of the 16th International Congress of Lymphology. *Lymphology* 1998;31 Suppl:1–621.
379. Johansson SK 2002 Lymphoedema and Breast Cancer. A Physiotherapeutic Approach. Studentlitteratur, Lund University, Lund. ISBN: 91-628-5117-9.
380. Johansson K, Albertsson M, Ingvar C, Ekdahl C. Effects of compression bandaging with or without manual lymph drainage treatment in patients with postoperative arm lymphedema. *Lymphology* 1999;32:103–10.
381. Johansson K, Lie E, Ekdahl C, Lindfeldt J. A randomized study comparing manual lymph drainage with sequential pneumatic compression for treatment of postoperative arm lymphedema. *Lymphology* 1998;31:56–64.
382. Johansson K, Ohlsson K. Factors associated with the development of arm lymphedema following breast cancer treatment: a match pair case-control study. *Lymphology* 2002;35:59-71.
383. Jungi WF: The prevention and management of lymphoedema after treatment for breast cancer. *Int Rehabil Med* 3:129–134, 1981
384. Karadibak D, Yavuzsen T, Saydam S. Prospective trial of intensive decongestive physiotherapy for upper extremity lymphedema. *J Surg Oncol.* 2008;97:572-577.

385. Karges JR: Assessing the relationship between water displacement and circumferential measurements in determining upper extremity volume in women with lymphedema. Krannert School of Physical Therapy, University of Indianapolis, Indianapolis, USA, 1996
386. Karges J, Mark B, Stikeleather S, Worrell T. Concurrent validity of upper extremity volume estimates: comparison of calculated volume derived from girth measurements and water displacement. *Phys Ther*, 2003;83:134-45.
387. Kasseroller, R 1998 Compendium of Dr Vodder's Manual Lymph Drainage, Haug, Heidelberg, 1st Ed.
388. Kasseroller, R. (1998). The Vodder School: The Vodder method. *Cancer* 83, 2840-2
389. Kelly DG. A primer on lymphedema. Upper Saddle River (NJ): Pearson Education, Inc.; 2002.
390. Kim, SJ: Lymphedema. Jungdam Media Publishing Co, Seoul, 2002.
391. Kim, S., & Yi, O. (2009). Effects of manual lymph drainage on cardiac autonomic tone in healthy subjects. *International Journal of Neuroscience*, 119, 1105-17. doi: 10.1080/00207450902834884
392. Kim SJ, Park YD. Effects of complex decongestive physiotherapy on the edema and the quality of life of lower unilateral lymphoedema following treatment for gynecological cancer. *Eur J Cancer Care* 2008; 17(5):463-8.
393. Kitamura Y, Ohno Y, Kasahara S, Murata K, Sugiyama H, Oshima A, et al. Statistical estimation of the number of breast cancer patients with disabilities resulting from surgery. *Breast Cancer* 2005;12:130-4.
394. Kligman L, Wong RK, Johnston M, Laetsch NS. (2004). The treatment of lymphedema related to breast cancer: a systematic review and evidence summary. *Support Care Cancer*, 12(6):421-31.
395. Klose G: Lymphedema Bandaging. Lohmann & Rauscher, Germany, 1998, pp 14-23
396. Ko DS, Lerner R, Klose G, Cosimi AB. Effective treatment of lymphedema of the extremities. *Arch Surg*. 1998;133:452-458.
397. Koak Z, Overgaard J. Risk factors of arm lymphedema in breast cancer patients. *Acta Oncol* 2000;39:389-92.
398. Kosir MA, Rymal C, Koppolu C, Hryniuk L, Draga L, Du W. Surgical outcomes after breast cancer surgery: measuring acute lymphedema. *J Surg Res* 2001;95:147-51.
399. Koul R, Dufan T, Russell C, et al. Efficacy of complete decongestive therapy and manual lymphatic drainage on treatment-related lymphedema in breast cancer. *Int J Radiat Oncol Biol Phys*. 2007;67:841-6.
400. Koul, R., Dufan, T., Russell, C., Guenther, W., Nugent, Z., Sun, X., & Cooke, A. (2006). Efficacy of complete decongestive therapy and manual lymphatic drainage on treatment-related lymphedema in breast cancer. *Int J Radiation Oncology Biol Phys*, doi: 10.1016/j.ijrobp.2006.09.024
401. Kurz I., 1996. Textbook of Dr Vodder's Manual Lymph Drainage (Treatment Manual), Vol. 3, 3rd Ed. Haug, Heidelberg
402. Lacomba, M., Sanchez, M., Goni, A., Merino, D., Mayoral de Moral, O., Tellez, E., & Minayo, E. (2009). Effectiveness of early physiotherapy to prevent lymphoedema after surgery for breast cancer: randomised, single blinded, clinical trial. *BMJ*,340(b5396), 1-8. doi: 10.1136/bmj.b5396
403. Lane K, Worsley D, McKenzie D. Exercise and the lymphatic system: implications for breast-cancer survivors. *Sports Med* 2005;35:461-71.
404. Lawenda BD, Mondry TE, Johnstone PAS. Lymphedema: a primer on the identification and management of a chronic condition in oncologic treatment. *CA Cancer J Clin*. 2009;59:8-24.
405. Leduc O, Leduc A, Bourgeois P, Belgrado JP: The physical treatment of upper limb edema. *Cancer* 83: 2835-2839, 1998
406. Liao SF, Huang MS, Li SH, Chen IR, Wei TS, Kuo SJ, et al. Complex decongestive physiotherapy for patients with chronic cancer-associated lymphedema. *J Formos Med Assoc* 2004;103(5):344-8.
407. Liao SF, Huang MS, et al. Successful complex decongestive physiotherapy for lymphedema and lymphocutaneous reflux of the female external genitalia after RT. *J Formos Med Assoc* 2003;102:404-406.
408. Liao, S., Li, S., & Huang, H. (2012). The efficacy of complex decongestive physiotherapy (cdp) and predictive factors of response to cdp in lower limb lymphedema (III) after pelvic cancer treatment. *Gynecologic Oncology*, 125, 712-15. doi: 10.1016/j.ygyno.2012.03.017
409. Little AG, Ferguson MK, Golomb HM, et al. (1986). Pleuroperitoneal shunting for malignant pleural effusions. *Cancer*, 58, 2740-3.
410. Liu N. Trafficking of hyaluronan in the interstitium and its possible implications. *Lymphology* 2004;37:6-14.
411. Lymphology Executive Committee: The diagnosis and treatment of peripheral lymphedema. Consensus document of the International Society of Lymphology Executive Committee. *Lymphology*. 28 (1995), 113-117.
412. MacLaren JA. Skin changes in lymphoedema: pathophysiology and management options. *Int J Palliat Nurs* 2001;7:381-8.
413. MacWayne, J, SP Heiney: Psychologic and social sequelae of secondary lymphedema: A review. *Cancer* 104 (2005), 457-466.
414. Masmoudi A, Maaloul I, Turki K, Elloumi Y, Marrekchi S, Bouassida S, et al. Erysipelas after breast cancer treatment (26 cases). *Dermatology Online Journal* 2004:12.
415. McKenzie DC, Kalda AL. Effect of upper extremity exercise on secondary lymphedema in breast cancer patients: a pilot study. *J Clin Oncol* 2003;21:463-6.
416. McNeely, M., Magee, D., Lees, A., Bagnall, K., Haykowsky, M., & Hanson, J. (2004). The addition of manual lymph drainage to compression therapy for breast cancer related lymphedema: a randomized controlled trial. *Breast Cancer Research and Treatment*, 86, 95-106.
417. McTiernan A. Obesity and cancer: the risks, science, and potential management strategies. *Oncology* 2005;19:871-81.
418. Meek AG. Breast radiotherapy and lymphedema. *Cancer* 1998;83:2788-2797.
419. Meeske K, Sullivan-Halley J, Smith A, McTiernan A, Baumgartner K, Harlan L, et al. Risk factors for arm lymphedema following breast cancer diagnosis in black women and white women. *Breast Cancer Res Treat* 2008;14:982-91.

420. Megens A., & Harris S. (1998). Physical therapist management of lymphedema following treatment for breast cancer: A critical review of its effectiveness. *Phys Ther*, 78, 1302--11.
421. Mondry T, Riffenburgh RH, Johnstone PA. Complete decon-gestive therapy for upper extremity lymphedema after breast cancer therapy. *Cancer J*. 2004;10:42-48.
422. Morgan RG, Casley-Smith JR, Mason MR. (1992). Complex physical therapy for the lymphoedematous arm. *J Hand Surg – British*, 17, 437–41.
423. Mortimer, P. S. (1998). The pathophysiology of lymphedema. *Cancer*, 83, 2798–2802.
424. Moseley AL, Piller NB, Carati CJ. The effect of gentle arm exercise and deep breathing on secondary arm lymphedema. *Lymphology* 2005;38:136–45.
425. Moseley A, Piller N 2002 Combined opto- electronic perometry and bio-impedance to measure objectively the effectiveness of a new treatment intervention for chronic secondary leg lymphoedema. *Lymphology* 35: 136–143
426. Moskovitz A, Anderson B, Yeung R, Byrd D, Lawton T, Moe R. Axillary web syndrome after axillary dissection. *Am J Surg* 2001;181:434-9.
427. Nardone L, Palazzoni G. Impact of dose and volume on lymphedema. *Rays* 2005;30:149-55.
428. Nesvold I, Dahl A, Løkkevik E, Marit Mengshoel A, Fosså S. Arm and shoulder morbidity in breast cancer patients after breast-conserving therapy versus mastectomy. *Acta Oncol* 2008;47:835-42.
429. Ocana A, Delgado C. Case 3. Upper limb lymphangiosarcoma following breast cancer therapy. *J Clin Oncol* 2006;24:1477-8.
430. Olszewski, W. L. (1991). *Lymph Stasis: Pathophysiology, Diagnosis and Treatment*. Florida: CRC Press.
431. Ozaslan C, Kuru M. Lymphedema after treatment of breast cancer. *Am J Surg* 2004;187:69-72.
432. Pain, SJ, SL Vowler, AD Purushotham: Is physical function a more appropriate measure than volume excess in the assessment of breast cancer-related lymphoedema (BCRL)? *Eur. J. Cancer* 39 (2003), 2168-2172.
433. Passik SD, McDonald MV: Psychosocial aspects of upper extremity lymphedema in women treated for breast carcinoma. *Cancer* 83: 2817–2820, 1998
434. Pereira de Godoy, JM, DM Braile, M de Fatima Godoy, et al: Quality of life and peripheral lymphedema. *Lymphology* 35(2002), 72-35.
435. Passik, SD, MV McDonald: Psychosocial aspects of upper extremity lymphedema in women treated for breast carcinoma. *Cancer* 83 (12 Suppl Am) (1998), 2817-2820.
436. Pecking A, Cluzan R, Despret-Curly JP 1983 Indirect lymphoscintigraphy in patients with limb edema. In *Progress in Lymphology, Proceedings of the 9th International Congress of Lymphology*, Tel Aviv.
437. Petlund CF. Volumetry of limbs. In: Olszewski WI, editor. *Lymph stasis: Pathophysiology, diagnosis and treatment*. Boston: CRC Press; 1991. p. 444 – 451.
438. Petrek, J. A., Pressman, P. I., & Smith, R. A. (2000). Lymphedema: Current issues in research and management. *CA: A Cancer Journal for Clinicians*, 50(5), 292–307.
439. Petrek, J. A., & Heelan, M. C. (1998). Incidence of breast carcinoma-related lymphedema. *Cancer*, 83, 2776–2781.
440. Petrek JA, Lerner R: Lymphedema. In: Harris JR, Lippman ME, Morrow M, Helman S, (eds) *Diseases of the Breast*. Lippincott-Raven, Philadelphia, 1996, pp 896–900.
441. Piller NB, Harris R 2002 Objective measurement of the effectiveness of a single session of manual lymphatic drainage on primary and secondary lymphoedema of the lower leg. *Lymphology* 35(Suppl): 289–292
442. Piller NB, Goodear M, Peter D 1998 Lymphoscintigraphic evidence supports the evidence of axillo-anastomotic pathways in a patient with chronic secondary leg lymphoedema subsequent to inguinal node clearance and radiotherapy. *European Journal of Lymphology* 6: 97–100
443. Pinell XA, Kirkpatrick SH, Hawkins K, et al. Manipulative therapy of secondary lymphedema in the presence of locore-gional tumors. *Cancer*. 2008;112:950-954.
444. Preston NJ, Seers K, Mortimer PS. Physical therapies for reducing and controlling lymphoedema of the limbs. *Cochrane Database of Systematic Reviews*. 2004; Issue 4. Art. No.: CD003141. DOI: 10.1002/14651858.CD003141.pub2.
445. Ramos SM, D'Donnell LS, Knight Galen. Edema volume, not timing, is the key to success in lymphedema treatment. *Am J Surg* 1991;178:311–5.
446. Ramos SM, O'Donnell LS, Knight G: Edema volume, not timing, is the key to success in lymphedema treatment. *Am J Surg* 178: 311–315, 1999
447. Ridner S. Pretreatment lymphedema education and identified educational resources in breast cancer patients. *Patient Educ Couns* 2006;61:72-9.
448. Rockson SG, Miller LT, Senie R, Brennan MJ, Casley-Smith JR, Foldi E, Walder AL: American Cancer Society Lymphedema Workshop. Workgroup III: Diagnosis and management of lymphedema. *Cancer* 83: 2882– 2885, 1998
449. Ruckson S. Precipitating factors in lymphedema: Myth and realities. *Cancer* 1998;83:2814 –2816.
450. Runowick C. Lymphedema: patient and provider education. In: American Cancer Society lymphedema workshop. American Cancer Society, 1998:2874-6.
451. Sander AP, Hajer NM, Hemenway K. Incidence and risk of arm oedema following treatment for breast cancer: a three-year follow-up study. *Q J Med* 2005;98:343-8.
452. Schillinger A, Koenig D, Haefele C, Vogt S, Heinrich L, Aust A, & Schmid A. (2006). Effect of manual lymph drainage on the course of serum levels of muscle enzymes after an extended treadmill exercise. *Am J Phys Med Rehabil*, 85, 516-20.

453. Scuba A, Achalu R, Rockson SG. (2002). Decongestive lymphatic therapy for patients with breast carcinoma associated lymphedema. A randomized, prospective study of a role for adjunctive intermittent pneumatic compression. *Cancer*, 95, 2260–7.
454. Shih YCT, Xu Y, Cormier JN, Giodarno S, Ridner SH, Buchholz TA. Incidence, treatment costs, and complications of lymphedema after breast cancer: a 2-year follow-up study. *J Clin Oncol* 2009;12:2001-14.
455. Simon MS, Cody RL: Cellulitis after axillary lymph node dissection for carcinoma of the breast. *Am J Med* 93: 543– 548, 1992
456. Sitzia, J, AW Stanton, C Badger: A review of outcome indicators in the treatment of chronic limb edema. *Clin. Rehabil.* 11 (1997), 181-191.
457. Sitzia, J, L Sobrido: Measurement of health-related quality of life of patients receiving conservative treatment for limb lymphoedema using Nottingham Health Profile. *Qual. Life. Res.* 6 (1997), 373-384.
458. Sitzja J. Volume measurement in lymphoedema treatment: examination of formula. *Eur J Cancer Care (Engl)* 1995;4:11–6.
459. Smeltzer, D. M., Stickler, G. B., & Schirger, A. (1985). Primary lymphedema in children and adolescents: A follow-up study and review. *Pediatrics*, 76(2), 206–218.
460. Stanton AWB, Badger C, Sitzia J 2000 Non invasive assessment of the lymphoedematous limb. *Lymphology* 33: 122–135
461. Stanton A, Northfield J, Holyroyd BP 1997 Validation of an optoelectronic limb volumeter (perometer). *Lymphology* 30: 77–97
462. Starling EH. Physiological factors involved in the causation of dropsy. *Lancet* 1896;Vol. 1:1267–1270.
463. Stout Gergich N, Pfalzer L, McGarvey C, Springer B, Gerber L, Soballe P. Preoperative assessment enables the early diagnosis and successful treatment of lymphedema. *Cancer* 2008;112:2809-18.
464. Stranden E. A comparison between surface measurements and water displacements volumetry for the quantification of edema. *J Oslo City Hosp* 1981;31:153–155.
465. Swedborg I. Effectiveness of combined methods of physiotherapy for post-mastectomy lymphoedema. *Scand J Rehabil Med* 1980;12:77–85.
466. Szuba A, Cooke J, Shuja Y, Rockson S. Decongestive lymphatic therapy for patients with cancer-related or primary lymphedema. *Am J Med.* 2000;109:296-300.
467. Szuba A, Cooke JP, Yousuf S, Rockson SG. (2000). Decongestive lymphatic therapy for patients with cancer-related or primary lymphedema. *Am J Med*, 109(4):296–300.
468. Szuba A, Achalu R, Rockson SG. Decongestive lymphatic therapy for patients with breast carcinoma-associated lymphedema. A randomized, prospective study of a role for adjunctive intermittent pneumatic compression. *Cancer* 2002;95:2260–7.
469. Tada H, Teramukai S, Fukushima M, Sasaki H. Risk factors for lower limb lymphedema after lymph node dissection in patients with ovarian and uterine carcinoma. *BMC Cancer* 2009;9:47.
470. Tan, I., Maus, E., Rasmussen, J., Marshall, M., Adams, K., Fife, C., & Sevick-Muraca, E. (2011). Assessment of lymphatic contractile function after manual lymphatic drainage using near- infrared fluorescence imaging. *Arch Phys Med Rehabil*, 92(5), 756-64. PMID: 21530723
471. Taylor R, Jayasingue U, Koelmeyer L, Ung O, Boyages J. Reliability and validity of arm volume measurements for assessment of lymphoedema. *Phys Ther* 2006;86:205-14.
472. Thomas RC, Hawkins K, Kirkpatrick SH, Mondry TE, Gabram-Mendola S, Johnstone PA. Reduction of lymph-edema using complete decongestive therapy: roles of prior radiation therapy and extent of axillary dissection. *J Soc Integr Oncol.* 2007;5:87-91.
473. The diagnosis and treatment of peripheral lymphedema. Consensus document of the International Society of Lymphology. *Lymphology* 2003;36:84–91.
474. Tobbia D, Semple J, Baker A, Dumont A, Semple A, Johnston M. Lymphedema development and lymphatic function following lymph node excision in sheep. *J Vasc Res* 2009;46:426-34.
475. Tobin MB, Lacey HJ, Meyer L, Mortimer PS: The psychological morbidity of breast cancer-related arm swelling. *Cancer* 72: 3248–3252, 1993
476. Todd J, Scally A, Dodwell D, Horgaan K, Topping A. A randomised controlled trial of two programmes of shoulder exercise following axillary lymph node dissection for invasive breast cancer. *Physiotherapy* 2008;94:265-73.
477. Velanovich, V, W Szymanski: Quality of life of breast cancer patients with lymphedema. *Am. J. Surg.* 177 (1999), 184-187.
478. Vignes S, Porcher R, Champagne A, Dupuy A. Predictive factors of response to intensive decongestive physiotherapy in upper limb lymphedema after breast cancer treatment: a cohort study. *Breast Cancer Res Treat* 2006;98(1):1–6.
479. Voogd AC, Ververs JM, Vingerhoets AJ, Roumen RM, Coebergh JW, Crommelin MA: Lymphoedema and reduced shoulder function as indicators of quality of life after axillary lymph node dissection for invasive breast cancer. *Br J Surg* 90: 76–81, 2003
480. Wantanabe RK, Miura K, Inoue K 1989 Evaluation of leg oedema using a multi-frequency impedance meter on patients with lymphatic obstruction. *Lymphology* 22: 85–92
481. Weiss, JM, BJ Spray: The effect of complete decongestive therapy on the quality of life of patients with peripheral lymphedema. *Lymphology* 35 (2002), 46-58.
482. Weissleder, H., & Schuchhardt, C. (2001). *Lymphedem: Diagnosis and Therapy*. Koln: Viavital Verlag GmbH.
483. Werner RS, McCormick B, Petrek J, et al. Arm edema in conservative management of breast cancer: Obesity is a major predictive factor. *Therapeut Radiol* 1991;180:177–184.
484. Williams AF, Vadgama A, Franks PJ, Mortimer PS. A randomized controlled crossover study of manual lymphatic drainage therapy in women with breast cancer-related lymphoedema. *Eur J Cancer Care (Engl)* 2002;11:254–61.
485. Wittlinger, H., & Wittlinger, G. (1998). *Textbook of Dr. Vodder's Manual Lymph Drainage*. Vol. I, 6th Ed. Heidelberg: Haug.

486. Wittlinger G, Wittlinger H. Textbook of Dr.Vodder's manual lymph drainage: Basic course. 6th ed. New York: Thieme Medical Publishers; 1998. p. 72–74.
487. Yamamoto R, Yamamoto T. (2007). Effectiveness of the treatment-phase of two-phase complex decongestive physiotherapy for the treatment of extremity lymphedema. *Int J Clin Oncol*, 12(6):463–8.
488. Zanolli R, Monzeglio C, Balzarini A, Martino G. Evaluation of the results of three different methods of postmastectomy lymphedema treatment. *J Surg Oncol* 1984;26:210–3.

Preventive and Wellness Services and Chronic Disease Management

489. Alraek, T., Lee, M., Choi, T., Cao, H., & Liu, J. (2011). Complementary and alternative medicine for patients with chronic fatigue syndrome: a systematic review. *BMC Complement Altern Med*, 11, 87. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/21982120>
490. "Studies of qigong, massage and tuina were demonstrated to have positive effects..."
491. Arroyo-Morales, M., Olea, N., Martinez, M., HidalgoLozano, A., Ruiz-Rodriguez, C., Rodriguez, L. (2008). Psychophysiological effects of massage-myofascial release after study. *J Altern Complement Med*, 14, 1223-9.
492. Barnes, P., Bloom, B., & Nahin, R. (2008). *Complementary and alternative medicine use among adults and children: United states, 2007. Nat Health Stat Rep*, 12, 1-23.
493. Benjamin, P., & Tappan, F. (2004). *Tappan's handbook of healing massage techniques: Classic, holistic and emerging methods*. (4th ed.). Pearson/Prentice Hall.
494. Denneson, L., Corson, K., & Dobscha, S. (2011). Complementary and alternative medicine use among veterans with chronic noncancer pain. *J Rehabil Res Dev*, 48(9), 1119-28. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/22234716>
495. "Chiropractic care was the least preferred option, whereas massage therapy was the most preferred (75% and 96%, respectively)."
496. Dishman, J., & Bulbulian, R. (2001). Comparison of effects of spinal manipulation and massage on motorneuron excitability. *Electromyelin C I Neurophysiol*, 41, 97-106.
497. Fattah, M., & Hamdy, B. (2011). Pulmonary functions of children with asthma improve following massage therapy. *J Altern Complement Med*, 17(11), 1065-8. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/22087616>
498. "A beneficial role for massage therapy in pediatric asthma is suggested. It improved the key pulmonary functions of the children, namely, FEV1 and FEV1/FVC ratio."
499. Field, T., Diego, M., Delgado, J., Garcia, D., & Funk, C. (2011). Hand pain is reduced by massage therapy. *Complementary Therapies in Clinical Practice*.
500. Harden VA. Koch's postulates and the etiology of AIDS: an historical perspective. *Hist Phil Life Sci* 1992;14:249–69. .
501. Hawk, C., Ndetan, H., & Willard, M. (2011). Potential role of complementary and alternative health care providers in chronic disease prevention and health promotion: An analysis of national health interview survey data. *Preventive Medicine*, doi: :10.1016/j.jpmed.2011.07.002
502. Hillier, S., Louw, Q., Morris, L., Uwimana, J., & Statham, S. (2010). Massage therapy for people with hiv/aids. *Cochrane Database Syst Rev*, (1), CD007502. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/20091636>
503. "There is some evidence to support the use of massage therapy to improve quality of life for people living with HIV/AIDS (PLWHA), particularly in combination with other stress-management modalities, and that massage therapy may have a positive effect on immunological function. "
504. *Institute of Medicine (IOM), Report on Complementary and Alternative Medicine in the US, National Academy of Sciences*. (2005).
505. Jane, S., Chen, S., Wilkie, D., Lin, Y., Foreman, S., Beaton, R., & Liao, M. (2011). Effects of massage on pain, mood status, relaxation, and sleep in taiwanese patients with metastatic bone pain: A randomized clinical trial. *Pain*, 152(10), 2432-42. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/21802850>
506. "... the reduction in pain with massage was both statistically and clinically significant, and the massage-related effects on relaxation were sustained for at least 16-18 hours postintervention. ... Overall, results from this study support employing MT as an adjuvant to other therapies in improving bone pain management."
507. Moyer, C. (2008). Affective massage therapy. *Int J Ther Massage and Bodyw*, 1, 35.
508. Moyer, C., Rounds, J., & Hannum, J. (2004). A meta-analysis of massage therapy research. *Psych Bull*, 130, 3-1 8.

Headache

509. (2006). Myofascial trigger points and their relationship to headache clinical parameters in chronic tension-type headache. *Headache*, 46(8), 1264-72.
510. Andrasik, F., Lipchik, G., McCrory, D., & Wittrock, D. (2005). Outcome measurement in behavioral headache research: Headache parameters and psychosocial outcomes. *Headache*, 45(5), 429-37.
511. Barna, S., & Hashmi, A. (2004). Occipital neuralgia. *Pain Manag Rounds* 1(7), 1-6.
512. Bendtsen, L. (2000). Central sensitization in tension-type headach: Possible patho-physiological mechanisms. *Cephalalgia*, 20(5), 486-508.
513. Calandre, E., Hidalgo, J., Garcia-Leiva, J., & RiceVillademoros, F. (2006). Trigger point evaluation in migraine patients: An indication of peripheral sensitization linked to migraine predisposition? *Eur J Neurol*, 13(3), 244-9.
514. Chakravarty, A., & Sen, A. (2010). Migraine, neuropathic pain and nociceptive pain: Towards a unifying concept. *Med Hypotheses*, 74(2), 225-31.

515. Chaibi, A., Tuchin, P., & Russell, M. (2011). Manual therapies for migraine: A systematic review. *J Headache Pain*, 12(2), 127-33. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/21298314>
516. "The RCTs suggest that massage therapy, physiotherapy, relaxation and chiropractic spinal manipulative therapy might be equally effective as propranolol and topiramate in the prophylactic management of migraine."
517. Chou, C., Chao, A., Lu, S., Hu, H., & Wang, S. (2004). Cephalic venous congestion aggravates only migraine-type headaches. *Cephalalgia*, 24(11), 975-9.
518. De Benedittis, G., & Lorenzetti, A. (1992). The role of stressful life events in the persistence of primary headache: Major events vs. daily hassles. *Pain*, 51(1), 35-42.
519. De Benedittis, G., Lorenzetti, A., & Pieri, A. (1990). The role of stressful life events in the onset of chronic primary headache. *Pain*, 40(1), 65-75.
520. Doepp, F., Schreiber, S., Dreier, J., Einhaupl, K., & Valdueza, J. (2003). Migraine aggravation caused by cephalic venous congestion. *Headache*, 43(2), 96-8.
521. Drummond, P., & Lance, J. (1983). Extracranial vascular changes and the source of pain in migraine headache. *Ann Neurol*, 13(1), 32-7.
522. Fernandez-de-las-Penas, C., Alonso-Blanco, C., Cuadrado, M., Gerwin, R., & Pareja, J. (2006). Trigger points in the suboccipital muscles and forward head posture in tension-type headache. *Headache*, 46(3), 454-60.
523. Fernandez-de-las-Penas, C., Alonso-Blanco, C., Fernandez-Carnero, J., & Miangolarra-Page, J. (2006). The immediate effect of ischemic compression technique and transverse friction massage on tenderness of active and latent myofascial trigger points: A pilot study. *J Bodyw Mov Ther*, 10(1), 3-9.
524. Fernindez-de-las-Peias, C., M.L. Cuadrado, L. Arendt-Nielsen, D.G. Simons, and J.A. Pareja. 2007. Myofascia1 trigger points and sensitization: An updated pain model for tension-type headache. *Cephalalgia* 27(5): 383493.
525. Fernandez-de-las-Penas, C., Ge, H., Arendt-Nielsen, L., Cuadrado, M., & Pareja, J. (2007). The local and referred pain from myofascial trigger points in the temporalis muscle contributes to pain profile in chronic tension-type headache. *CJP*, 23(9), 786-92.
526. Fernandez-de-las-Penas, C., Cleland, J., Cuadrado, M., & Pareja, J. (2008). Predictor variables for identifying patients with chronic tension-type headache who are likely to achieve short-term success with muscle trigger point therapy. *Cephalalgia*, 28(3), 264-75.
527. Fernandez-de-las-Penas, C., Madeleine, P., Cuadrado, M., Ge, H., Arendt-Nielsen, L., & Pareja, J. (2009). Pressure pain sensitivity mapping of the temporalis muscle revealed bilateral pressure hyperalgesia in patients with strictly unilateral migraine. *Cephalalgia*, 29(6), 670-6.
528. Fryer, G., & Hodgson, L. (2005). The effect of manual pressure release on myofascial trigger points in the upper trapezius muscle. *J Bodyw Mov Ther*, 9, 248-55.
529. Fukui, P., Goncalves, T., Strabelli, C., Lucchino, N., Matos, F., Santos, J., & Peres, M. (2008). Trigger factors in migraine patients. *Arq Neuro-psiquiatr*66(3A): 494499.
530. Gaul, C., Eismann, R., Schmidt, T., May, A., Leinisch, E., Wieser, T., Evers, S., Henkel, K., Franz, G., & Zierz, S. (2009). Use of complementary and alternative medicine in patients suffering from primary headache disorders. *Cephalalgia*, 29(10), 1069-78
531. Hammill, J., Cook, T., & Rosecrance, J. (1996). Effectiveness of a physical therapy regimen in the treatment of tension-type headache. *Headache*, 36(3), 149-53.
532. Hauge, A., Kirchrnann, M., & Olesen, J. (2010). Trigger factors in migraine with aura. *Cephalalgia*, 30(3), 346-53.
533. Heneghan, N., Adab, P., Balanos, G., & Jordan, R. (2012). Manual therapy for chronic obstructive airways disease: A systematic review of current evidence. *Man Ther*, Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/22703901>
534. "...patient reported measures for 'improved health' and 'breathing difficulty' however did improve following OMT compared to the control. Evidence for MT as an adjunctive management approach for COPD is lacking...."
535. Hernandez-Reif, M., Dieter, I., Field, T., Swerdlow, B., & Diego, M. (1998). Migraine headaches are reduced by massage therapy. *Int J Neurosci*, 96, 1-11.
536. Herren-Gerber, R., Weiss, S., Arendt-Nielsen, L., Petersen-Felix, S., Di Stefano, G., Radanov, B., & Curatolo, M. (2004). Modulation of central hypersensitivity by nociceptive input in chronic pain after whiplash injury. *Pain Med*, 5(4), 366-76.
537. Holroyd, K. (2002). Behavioral and psychologic aspects of the pathophysiology and management of tension-type headache. *Curr Pain Headache Rep*, 6(5), 401-7.
538. Holroyd, K., Malinoski, P., Davis, M., & Lipchik, G. (1999). The three dimensions of headache impact: Pain, disability and affective distress. *Pain*, 83(3), 571-8.
539. Holroyd, K., Stensland, M., Lipchik, G., Hill, K., O'Donnell, F., & Cordingley, G. (2000). Psychosocial correlates and impact of chronic tension-type headaches. *Headache*, 40(1), 3-16.
540. Houle, T., & Nash, J. (2008). Stress and headache chronification. *Headache*, 48(1), 40-4.
541. Jensen, R., & Stovner, L. (2008). Epidemiology and comorbidity of headache. *Lancet Neurol*, 7(4), 354-61.
542. Juang, K., Wang, S., Fuh, J., Lu, S., & Su, T. (2000). Comorbidity of depressive and anxiety disorders in chronic daily headache and its subtypes. *Headache*, 40(10), 818-23.
543. Kuan, T. (2009). Current studies on myofascial pain syndrome. *Curr Pain Headache Rep*, 13(5), 365-9.
544. Lawler, S., & Cameron, L. (2006). A randomized, controlled trial of massage therapy as a treatment for migraine. *Ann Behav Med*, 32(1): 50-9.
545. Leistad, R., Sand, T., Westgaard, R., Nilsen, K., & Stovner, L. (2006). Stress-induced pain and muscle activity in patients with migraine and tension-type headache. *Cephalalgia*, 26(1), 64-73.

546. Linde, K., Allais, G., Brinkhaus, B., Manheimer, E., Vickers, A., & White, A. (2009). Acupuncture for migraine prophylaxis. *Cochrane Db Syst Rev* 1: CD001218.
547. Lipchik, G., Holroyd, K., France, C., Kvaal, S., Segal, D., Cordingley, G., Rokicki, L., & McCool, H. (1996). Central and peripheral mechanisms in chronic tension-type headache. *Pain*, 64(3), 467-75.
548. Lipton, R., Diamond, S., Reed, M., Diamond, M., & Stewart, W. (2001). Migraine diagnosis and treatment: Results from the American Migraine Study II. *Headache*, 41(7), 638-45.
549. Lipton, R., Stewart, W., Diamond, S., Diamond, M., & Reed, M. (2001). Prevalence and burden of migraine in the United States: Data from the American Migraine Study II. *Headache*, 41(7), 646-57.
550. Lipton, S. (1986). Prevention of classic migraine headache by digital massage of the superficial temporal arteries during visual aura. *Ann Neurol*, 19(5), 515-6.
551. Melzack, R. (1999). From the gate to the neuromatrix. *Pain*, 6: S121-6.
552. Mongini, F., Ciccone, G., Deregibus, A., Ferrero, L., Mongini, T. (2004). Muscle tenderness in different headache types and its relation to anxiety and depression. *Pain*, 112(1-2), 59-64.
553. Moraska, R., & Chandler, C. (2009). Changes in psychological parameters in patients with tension-type headache following massage therapy: A pilot study. *J Man Manip Ther*, 17(2), 86-94.
554. Moraska, A., Chandler, C. (2008). Changes in clinical parameters in patients with tension-type headache following massage therapy: A pilot study. *J Man Manip Ther*, 16(2), 106-12.
555. Olesen, J., Boussier, M., Diener, D., Dodick, D., First, M., Goadsby, P., & Steiner, T. (2004). The international classification of headache disorders: 2nd edition. *Cephalalgia*, 24(1), S9-S160.
556. Olesen, J., Burstein, R., Ashina, M., & Tfelt-Hansen, P. (2009). Origin of pain in migraine: Evidence for peripheral sensitisation. *Lancet Neurol*, 8(7), 679-90.
557. Pascual, L., Colas, R., & Castillo, J. (2001). Epidemiology of chronic daily headache. *Curr Pain Headache Rep*, 5(6), 529-36.
558. Piovesan, E., Di Stani, F., Kowacs, P., Mulinaxi, R., Radunz, V., Utiumi, M., Muranka, E., Giublin, M., & Werneck, L. (2007). Massaging over the greater occipital nerve reduces the intensity of migraine attacks: Evidence for inhibitory trigemint, cervical convergence mechanisms. *Arq Neuro-psiquiah*, 65(3A), 599-604.
559. Puustjarvi, K., Airaksinen, O., & Pontinen, P. (1990). The effects of massage in patients with chronic tension headache. *Acupunct Electrother Res*, 15(2), 159-62,
560. Quinn, C., Chandler, C., & Moraska, A. (2002). Massage therapy and frequency of chronic tension headaches. *Am J Public Health*, 92(10), 1657-61.
561. Rasmussen, B. (1993). Migraine and tension-type headache in a general population: Precipitating factors, female hormones, sleep pattern and relation to lifestyle. *Pain*, 53(1), 65-72.
562. (1999). Epidemiology and socio-economic impact of headache. *Cephalalgia*, 19(25), S20-3.
563. Rasmussen, B., Jensen, R., & Olesen, J. (1992). Impact of headache on sickness absence and utilisation of medical services: A Danish population study. *J Epidemiol Community Health*, 46(4): 443-6 .
564. Rossi, P., Di Lorenzo, G., Faroni, J., Malpezzi, M., Cesarino, F., & Nappi, G. (2006). Use of complementary and alternative medicine by patients with chronic tension-type headache: Results of a headache clinic survey. *Headache*, 46(4), 622-31.
565. Schwartz, B., Stewart, W., & Lipton, R. (1997). Lost workdays and decreased work effectiveness associated with headache in the workplace. *J Occup Environ Med*, 39(4), 320-7.
566. Schwartz, B., Stewart, W., Simon, D., & Lipton, R. (1998). Epidemiology of tension-type headache. *JAMA*, 279(5), 381-3.
567. Stovner, L., Hagen, K., Jensen, R., Katsarava, Z., Lipton, R., Scher, A., Steiner, T., & Zwart, J. (2007). The global burden of headache: A documentation of headache prevalence and disability worldwide. *Cephalalgia*, 27(3), 193-210.
568. Torelli, P., Jensen, R., & Olesen, J. (2004). Physiotherapy for tension-type headache: A controlled study. *Cephalalgia*, 24(1), 29-36.
569. Toro-Velasco, C., Arroyo-Morales, M., Fernandez-de-las-Penas, C., Cleland, J., & Barrero-Hernandez, E. (2009). Short-term effects of manual therapy on heart rate variability, mood state, and pressure pain sensitivity in patients with chronic tension-type headache: A pilot study. *J Manipulative Physiol Ther*, 32(7), 527-35.
570. Treaster, D., Marras, W., Burr, D., Sheedy, J., & Hart, D. (2006). Myofascial trigger point development from visual and postural stressors during computer work. *J Electromyogr Kinesiol*, 16(2), 115-24.
571. Turkdogan, D., Cagirici, S., Soylemez, D., Sur, H., Bilge, C., & Turk, U. (2006). Characteristic and overlapping features of migraine and tension-type headache. *Headache*, 46(3), 461-8.
572. Zanchin, G., Maggioni, F., Granella, F., Rossi, P., Falco, L., & Manzoni, G. (2001). Self-administered pain-relieving manoeuvres in primary headaches. *Cephalalgia*, 21(7), 718-26.
573. Zwart, J., Dyb, G., Hagen, K., Odegard, K., Dahl, A., Bovim, G., & Stovner, L. (2003). Depression and anxiety disorders associated with headache frequency. The Nord-Trondelag Health Study. *Eur J Neurol*, 10(2), 147-52
- Neck and Shoulder Pain**
574. Beaudreuil, J., Nizard, R., Thomas, T., Peyre, M., Liotard, J., Boileau, P., & Walch, G. (2009). Contribution of clinical tests to the diagnosis of rotator cuff disease: A systematic literature review. *Joint, Bone, Spine: Reuue du Rhumatisme*, 76(1), 15-9.
575. Borghouts, J., Koes, B., & Bouter, L. (1998). The clinical course and prognostic factors of nonspecific neck pain: A systematic review. *Pain*, 77(1), 1-13.
576. Boudreau, L., & Pinto, A. (2001). Acute lymphangitis mimicking mechanical neck pain. *J Manip Physiol Ther*, 24(7), 474-76.

577. Brosseau, L., Tugwell, P., Wells, G., & Panel P., (2001). Philadelphia Panel evidence based clinical practice guidelines on selected rehabilitation interventions for neck pain. *Phys Ther* 81(10), 1701-17.
578. Brosseau, L., Wells, G., Tugwell, P., Casimiro, L., Novikov, M., Loew, L., & School of Rehabilitation Sciences, Faculty of Health Sciences, University of Ottawa, Ottawa, Ontario, Canada, (2012). Ottawa panel evidence-based clinical practice guidelines on therapeutic massage for neck pain. *J Bodyw Mov Ther*, 16(3), 300-25.
579. Brosseau, L., Wells, G., Tugwell, P., & Casimiro, L. (2012). Ottawa panel evidence-based clinical practice guidelines on therapeutic massage for neck pain. *J Bodyw Mov Ther*, 16(3), 300-25. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/22703740>
580. "The Ottawa Panel was able to demonstrate that the massage interventions are effective for relieving immediate post-treatment neck pain symptoms, but data is insufficient for long-term effects."
581. Carroll, L., Holm, L., Hogg-Johnson, S., Cote, P., Cassidy, J., Haldeman, S., Nordin, M., & Guzman, J. (2008). Course and prognostic factors for neck pain in whiplash-associated disorders (WAD): Results of the bone and joint decade 2000-2010 task force on neck pain and its associated disorders. *Spine*, 33(4), S83-92.
582. Carroll, L., Hogg-Johnson, S., Cote, P., Cassidy, D., Haldeman, S., Nordin, M., & Guzman J. (2008). Course and prognostic factors for neck pain in workers: Results of the bone and joint decade 2000-2010 task force on neck pain and its associated disorders. *Spine*, 33(4), S93-100.
583. Carroll, L., Hogg-Johnson, S., van der Velde, G., Haldeman, S., Holm, L., Carragee, E., & Cassidy, J. (2008). Course and prognostic factors for neck pain in the general population: Results of the bone and joint decade 2000-2010 task force on neck pain and its associated disorders. *Spine*, 33(4), S75-82.
584. Cote, P., Cassidy, J., & Carroll, L. (2000). The factors associated with neck pain and its related disability in the Saskatchewan population. *Spine*, 25(9), 1109-17.
585. Day, L., Stecco, C. & Stecco A., (2009). Application of fascial manipulation technique in chronic shoulder pain: Anatomical basis and clinical implications. *J Bodyw Mov Ther*, 13(2), 128-35.
586. Dinnes, J., Loveman, E., McIntyre, L., & Waugh, N. (2003). The effectiveness of diagnostic tests for the assessment of shoulder pain due to soft tissue disorders: A systematic review. *Health Technol Asses*, 7(29), iii, 1-166.
587. Eisenberg, D., Davis, R., Ettner, S., Appel, S., Wilkey, S., Van Rompay, M., & Kessler, R. (1998). Trends in alternative medicine use in the United States, 1990-97: Results of a follow-up national survey. *JAMA*, 280(18), 1569-75.
588. Ezzo, J. (2007). What can be learned from Cochrane systematic reviews of massage that can guide future research? *J Altern Complem Med*, 13(2), 291-95.
589. Furlan, A., Imamura, M., Dryden, T., & Irvin, E. (2009). Massage for low back pain. *Spine*, 34(16), 1669-84.
590. Gross, A., Haines, T., Goldsmith, C., Santaguida, L., McLaughlin, L., Peloso, P., & Cervical Overview Group (COG). (2009). Knowledge to action: A challenge for neck pain treatment. *J Orthop Sport Phys*, 39(5), 351-63.
591. Guzman, J., Haldeman, S., Carroll, L., Carragee, E., Hurwitz, E., Peloso, P., & Hogg-Johnson, S. (2008). Clinical practice implications of the bone and joint decade 2000-2010 task force on neck pain and its associated disorders: From concepts and findings to recommendations. *Spine*, 33(4), S199-213.
592. Guzman, J., Hurwitz, E., Carroll, L., Haldeman, S., Cote, P., Carragee, E., P.M. Peloso, & J. D. Cassidy. (2008). A new conceptual model of neck pain: Linking onset, course, and care: The bone and joint decade 2000-2010 task force on neck pain and its associated disorders. *Spine*, 33 (4), S14-23.
593. Guzman, J., E.L.Hurwitz, L.J. Carroll, S. Haldeman, P. Cbte, E J Carragee, P.M. Peloso, G. van der Velde, L.W. Holm, S. Hogg-Johnson, M. Nordin, and J. D.Cassidy. 2008. A new conceptual model of neck pain: Linking onset, course,and care: The bone and joint decade 2000-2010 task force on neck pain and its associated disorders. *Spine*, 33(4): S14-S23.
594. Haas, M., Group, E., Panzer, E., Partna, L., Lumsden, S., & M, Aickin, M. (2003). Efficacy of cervical endplay assessment as an indicator for spinal manipulation. *Spine*, 28(11), 1091-96.
595. Haines, T., Gross, A., Goldsmith, C., Perry, L. (2008). Patient education for neck pain with or without radiculopathy. *Cochrane Db Syst Rev*, 4, CD005106.
596. Haldeman, S., Carroll, L., & Cassidy, L. (2008). The empowerment of people with neck pain: Introduction: The bone and joint decade 2000-2010 task force on neck pain and its associated disorders. *Spine*, 33(4), S8-13.
597. Haraldsson, B., Gross, A., Myers, C., Ezzo, J., Morien, A., Goldsmith, C., & Cervical Overview Group. (2006). Massage for mechanical neck disorders. *Cochrane Db Syst Rev*, 3, CD004871.
598. Hilbert, J., Sforzo, G., & Swensen, T. (2003). The effects of massage on delayed onset muscle soreness. *Brit J Sport Med*, 37, 72-5.
599. Hort, C., Sole, G., Munn, J. (2009). The effectiveness of manual therapy in the management of musculoskeletal disorders of the shoulder: A systematic review. *Manual Ther*, 14(5), 463-74.
600. Hughes, P., Taylor, N., & Green, R. (2008). Most clinical tests cannot accurately diagnose rotator cuff pathology: A systematic review. *Aust J Physiother*, 54, 159- 70.
601. Hurwitz, E., Carragee, E., van der Velde, G., Carroll, L., Nordin, M., Guzman, J., & Haldeman, S. (2008). Treatment of neck pain: Noninvasive interventions: Results of the bone and joint decade 2000-2010 task force on neck pain and its associated disorders. *Spine*, 33(4), S123-152.
602. Jull, G., Sterling, M., Falla, D., Treleaven, J., & O'Leary, S. (2008). *Whiplash, headache, and neck pain: Research-based directions for physical therapies*. 1st ed. Edinburgh: Churchill Livingstone.
603. Martin, B., Deya, R., Mirza, S., Turner, J., Cornstock, B., Hollingworth, W., & Sullivan, S. (2008). Expenditures and Health Status Among Adults With Back and Neck Problems. *JAMA*, 299(6), 656-64.

604. Meislin, R., Sperling, J., & Stitik, T. (2005). Persistent shoulder pain: Epidemiology, pathophysiology, and diagnosis. *Am J Orthopedics*, 34(12), S5-9.
605. Moyer, C., Rounds, J., & Hannum, J. (2004). A meta-analysis of massage therapy research. *Psychol Bull*, 130, 3-18.
606. Nordin, M., Carragee, E., Hogg-Johnson, S., Weiner, S., Hurwitz, E., Peloso, P., & Haldeman, S. (2008). Assessment of neck pain and its associated disorders: Results of the bone and joint decade 2000-2010 task force on neck pain and its associated disorders. *Spine*, 33(4), S101-22.
607. Olaya-Contreras, P., & Sty, J. (2009). Illness behavior in patients on long-term sick leave due to chronic musculoskeletal pain. *Acta Orthopaedica*, 80(3), 380-85.
608. Patedex, D., Berg, J., & R Thal. (2009). Neck and shoulder pain: differentiating cervical spine pathology from shoulder pathology. *J Surg Orthop Adv* 18(4), 170-74.
609. Quinlan, K., Annest, J., Myers, B., Ryan, G., & Hill, H. (2004). Neck strains and sprains among motor vehicle occupants: United States, 2000. *Accident Anal Prev*, 36(1), 21-27.
610. Rickards, L. (2006). The effectiveness of non-invasive treatments for active myofascial trigger point pain: A systematic review of the literature. *Int J Osteopath Med*, 9(4), 120-36.
611. Rubinstein, S., Pool, J., van Tulder, M., Riphagen, I., & de Vet, H. (2007). A systematic review of the diagnostic accuracy of provocative tests of the neck for diagnosing cervical radiculopathy. *Eur Spine J*, 16(3), 307-19.
612. Sherman, K., Cherkin, D., Hawkes, R., Miglioretti, D., & Deyo, R. (2009). Randomized trial of therapeutic massage for chronic neck pain. *Clin J Pain*, 25(3), 233-38.
613. Sherman, K., Cherkin, D., Kahn, J., Erro, J., Hrbek, A., Deyo, R., & Eisenberg, D. (2005). A survey of training and practice patterns of massage therapists in two U.S. states. *BMC Complement Altern M*, 5, 13.
614. Sherman, K., Cherkin, D., Hawkes, R., Miglioretti, D., & Deyo, R. (2009). Randomized trial of therapeutic massage for chronic neck pain. *Clin J Pain*, 25, 233-238.
615. Stevenson, J., & Trojian, T. (2002). Evaluation of shoulder pain. *J Fam Practice*, 51(7), 605-11.
616. Spitzer, W., Skovron, M., Salmi, L., Cassidy, J., Duranceau, I., Suissa, & Zeiss, E. (1995). Scientific monograph of the Quebec task force on whiplash-associated disorder: Redefining 'whiplash' and its management. *Spine*, 20(8), S1-73.
617. Vassiliou, T., Kaluza, G., Putzke, C., Wulf, H., & Schnabel, M. (2006). Physical therapy and active exercises: An adequate treatment for prevention of late whiplash syndrome? Randomized controlled trial in 200 patients. *Pain*, 124(1), 69-76.
618. Verhagen, A., Scholten-Peeters, G., van Wijngaard, S., de Bie, R., & Bierma-Zeinstra, S. (2007). Conservative treatments for whiplash. *Cochrane DbSyst Rev*, 2: CD003338.
619. Walton, D., Pretty, J., Macdermid, J., & Teasell, R. (2009). Risk factors for persistent problems following whiplash injury: Results of a systematic review and meta-analysis. *J Orthop Sport Phys*, 39(5), 334-50.
620. Weerapong, P., Hume, P., & Kolt, G. (2005). The mechanisms of massage and effects on performance, muscle recovery and injury prevention. *Sports Med*, 35, 235-56.
621. Williams, M., Williamson, E., Gates, S., Lamb, S., & Cooke, M. (2007). A systematic literature review of physical prognostic factors for the development of late whiplash syndrome. *Spine*, 32(25), E764-80.
- Low Back Pain**
622. (2012). Stud health technol inform. 176, 411-4. Furlan, A., Imamura, M., Dryden, T., Irvin, E., & Institute for Work & Health, Toronto, ON, Canada, (2009). Massage for low back pain: An updated systematic review within the framework of the cochrane back review group. *Spine*, 34(16), 1669-84.
623. Anderson, G. (1999). Epidemiological features of chronic low-back pain. *Lancet*, 354, 581-5.
624. Bell, J. (2008). Massage therapy helps to increase range of motion, decrease pain and assist in healing a client with low back pain and sciatica symptoms. *J Bodyw Mov Ther*, 12, 281-9.
625. Buttagat, V., Eungpinichpong, W., Chatchawan, U., & Kharmwan, S. (2011). The immediate effects of traditional Thai massage on heart rate variability and stress-related parameters in patients with back pain associated with myofascial trigger points. *Journal of Bodywork & Movement Therapies*, 15, 15-23. Retrieved from 10.1016/j.jbmt.2009.06.005
626. Chatchawan, U., Thinkhamrop, B., & Kharmwan, S. 2005. Effectiveness of traditional Thai massage versus Swedish massage among patients with back pain associated with myofascial trigger points. *J Bodyw Moo Ther*, 9(4), 298-309.
627. Chaitow, L. (2004). *Palpation and assessment skills*. 2nd ed. Edinburgh: Churchill Livingstone.
628. Chenot, J., Becker, A., Leonhardt, C., Keller, S., Donner-Banzho, N., Baum, E., Pflingsten, M., Hildebrandt, J., Basler, H., & Kochen, M. (2007). Use of complementary alternative medicine for low back pain consulting in general practice: A cohort study. *BMC Complement Altern Med*, 7, 42.
629. Cherkin, D., Eisenberg, D., Sherman, K., Barlow, W., Kaptchuk, T., Street, J., & Deyo, R. (2001). Randomized trial comparing traditional Chinese medical acupuncture, therapeutic massage, and self-care education for chronic low back pain. *Arch Intern Med*, 161(8), 1081-8.
630. Cherkin, D., Sherman, K., Kahn, J., Erro, J., Deyo, R., Haneuse, S., & Cook, A. (2009). Effectiveness of focused structural massage and relaxation massage for chronic low back pain: Protocol for a randomized controlled trial. *Trials*, 10(1), 96.
631. Cherkin, D., Sherman, K., Kahn, J., Wellman, R., Cook, A., Johnson, E., Erro, J., Delaney, K., & Deyo, R. (2011). A comparison of the effects of 2 types of massage and usual care on chronic low back pain. *Ann of Intern Med*, 155, 1-9.

632. Chou, R., Huffman, L., & American Pain Society; American College of Physicians, (2007). Nonpharmacologic therapies for acute and chronic low back pain: A review of the evidence for an american pain society/american college of physicians clinical practice guideline. *Ann Intern Med*, 147(7), 492-504.
633. Deyo, R., Battie M., Beurskens, A., Bombardier, C., Croft, P., Koes, B., Malmivaara, A., Roland, M., Von Korff, M., & Waddell G. (1998). Outcome measures for low back pain research: A proposal for standardized use. *Spine*, 23(18), 2003-13.
634. Eisenberg, D., Davis, R., Ettner, S., Appel, S., Wilkey, S., Van Rompay, M., & Kessler, R. (1998). Trends in alternative medicine use in the United States, 1990-1997: Results of a follow-up national survey. *JAMA*, 280(18), 1569-75.
635. Ernst, E. (1999). Massage therapy for low back pain: a systematic review. *J Pain Symptom Manage*, 17(1), 65-9.
636. Ernst, E. (2003). The safety of massage therapy. *Rheumatology*, 42, 1101-6.
637. Ezzo, J. (2007). What can be learned from Cochrane systematic reviews of massage that can guide future research? *J Altern Complem Med*, 13(2), 291-6.
638. Farasyn, A., Meeusen, R., & Nijs, J. (2006). A pilot randomized placebo-controlled trial of roptrotherapy in patients with subacute non-specific low back pain. *J Back Musculoskelet*, 19, 111-7.
639. Field, T., Hernandez-Reif, M., Diego, M., & Fraser, M. (2007). Lower back pain and sleep disturbance are reduced following massage therapy. *J Bodyw Moo Ther*, 11, 141-5.
640. Franke, A., Gebauer, Franke, S., & Brockow, T. (2000). Acupuncture massage vs. Swedish massage and individual exercise vs. group exercise in low back pain sufferers: A randomized controlled clinical trial in a 2-by-2 factorial design. *Forsch Komp Klass Nat*, 7(6), 286-93.
641. Fregni, F., Imamura, M., Chien, H., Lew, H., Boggio, P., Kaptchuk, T., Riberto, M., Hsing, W., Battistella, L., & Furlan, L. (2010). International placebo symposium working group. Challenges and recommendations for placebo controls in randomized trials in physical and rehabilitation medicine: A report of the international placebo symposium working group. *Am J Phys Med Rehabil*, 89(2), 160-72.
642. Furlan, A., Imamura, M., Dryden, T., & Irvin, E. (2008). Massage for low-back pain. *Cochrane Db Syst Rev*, 8(4), CD001929.
643. Furlan, A., Imamura, M., Dryden, T., & Irvin, E. (2009). Massage for low back pain: An updated systematic review within the framework of the Cochrane Back Review Group. *Spine*, 34(16), 1669-84.
644. Geisser, M., Wiggert, E., Haig, A., Colwell, M. (2005). A randomized, controlled trial of manual therapy and specific adjuvant exercise for chronic low back pain. *Clin J Pain*, 21(6), 463-70.
645. Goats, G. (1994). Massage: The scientific basis of an ancient art: Part 2. Physiological and therapeutic effects. *Br J Sports Med*, 28(3), 153-6.
646. Haraldsson, B., Gross, A., Myers, C., Ezzo, J., Morien, A., Goldsmith, C., & Branfort, G. (2006). Cervical overview group. Massage for mechanical neck disorders. *Cochrane Db Syst Rev*, 3, CD004871.
647. Hernandez-Reif, M., Field, T., Krasnegor, J., & Theakston, H. (2001). Lower back pain is reduced and range of motion increased after massage therapy. *Int J Neurosci*, 106, 131-45.
648. Hernandez-Reif, M., Field, T., Krasnegor J., & Theakston, H. (2001). Lower back pain is reduced and range of motion increased after massage therapy. *Int J Neurosci*, 106, 131-45.
649. Hsieh, L., Kuo, C., Yen, M., & Chen, T. (2004). A randomized controlled clinical trial for low back pain treated by acupressure and physical therapy. *Prev Med*, 39(1), 168-76.
650. Hsieh, L., Kuo, C., Lee, L., Yen, A., Chien, K., & Chen, T. (2006). Treatment of low back pain by acupressure and physical therapy: Randomised controlled trial. *BMJ*, 332(7543), 696-700.
651. Kania, A., Porcino, A., & Verhoef, M. (2008). Value of qualitative research in the study of massage therapy. *Int J Ther Massage Bodywork*, 1(2), 6-11.
652. Lee, M., Itoh, K., & Yang, G. (1990). *Physical therapy and rehabilitation medicine: Massage*. In The management of pain, ed. J.J. Bonica, 1777-8. Philadelphia: Lea and Febiger.
653. Little, P., Lewith, G., Webley, F., Evans, M., Beattie, A., Middleton, K., & Sharp, D. (2008). Randomised controlled trial of Alexander technique lessons, exercise, and massage (ATEAM) for chronic and recurrent back pain. *BMJ*, 19(337), a884.
654. Liu, J., & Zhang, S. (2000). Treatment of protrusion of lumbar intervertebral disc by pulling and turning manipulations. *J Tradit Chin Med*, 20, 195-7.
655. Mackawan, S., Eungpinichpong, W., Panturnethakul, R., Chatchawan, U., & Hunsawong, T. (2007). Effects of traditional Thai massage versus joint mobilization on substance P and pain perception in patients with non-specific low back pain. *J Bodyw Mov Ther*, 11(1), 416.
656. Melzack, R., & Wall, P. (1996). *The challenge of pain*. 2nd ed. London: Penguin Books.
657. Moher, D., Schulz, K., & Altman, D. (2001). The CONSORT statement: Revised recommendations for improving the quality of reports of parallel-group randomized trials. *Ann Intern Med*, 134(8), 657-62.
658. Morhenn, V., Park, J., Piper, E. & Zak, P. (2008). Monetary sacrifice among strangers is mediated by endogenous oxytocin release after physical contact. *Evol Hum Behav*, 29, 375-83.
659. Moyer, C., Rounds, J., & Hannum, J. (2004). A meta-analysis of massage therapy research. *Psychol Bull*, 130, 318.
660. Moyer, C., Dryden, T., & Shipwright, S. (2009). Directions and dilemmas in massage therapy research: A workshop report from the 2009 North American Research Conference on Complementary and Integrative Medicine. *Int J Ther Massage Bodywork*, 2(2), 15-27.
661. Poole, H., Glenn, S., & Murphy, P. (2007). A randomised controlled study of reflexology for the management of chronic low back pain. *Eur J Pain*, 11(8), 878-87.

662. Preyde, M. (2000). Effectiveness of massage therapy for subacute low-back pain: A randomized controlled trial. *CMAJ*, 162(13), 1815-20.
663. Quinn, F., Hughes, C., & Baxter, G. (2008). Reflexology in the management of low back pain: A pilot randomised controlled trial. *Complement Ther Med*, 16(1), 3-8.
664. Rachlin, I. (2002). *Physical therapy treatment approaches for myofascial pain syndromes and fibromyalgia: Therapeutic massage in the treatment of myofascial pain syndromes and fibromyalgia*. In Myofascial pain and fibromyalgia. Trigger point managemen4 ed. E.S. Rachlin and I. Rachlin, 467-87. St. Louis: Mosby.
665. Roland, M., & Fairbank, J. (2000). The Roland-Morris Disability Questionnaire and the Oswestry Disability Questionnaire. *Spine*, 25(24), 3115-24.
666. Sagar, S., Dryden, T., & Wong, K. (2007). Massage therapy for cancer patients: A reciprocal relationship between body and mind. *Current Oncology*, 14(2), 45-56.
667. Scott, J., & Huskisson, E. (1976). Graphic representation of pain. *Pain*, 2, 175-84.
668. Sefton, J., Yarar, C., Berry, J., & Pascoe, D. (2010). Therapeutic massage of the neck and shoulders produces changes in peripheral blood flow when assessed with dynamic infrared thermography. *J Altern Complement Med*, 16(7), 725-32.
669. Simons, D., Travell, J., & Simons, L. (1999). *Apropos of all muscles: Trigger point release*. In Travell & Simons ' myofascial pain and dysfunction: The trigger point manual. Upper half of body. 2nd ed., ed. D.G. Simons, 94-177. Baltimore: Williams & Wilkins.
670. Vickers, A., & Zollman, C. (1999). ABC of complementary medicine. Massage therapies. *BMJ*, 319(7219), 1254-7.
671. Waddell, G. (1998). *The back pain revolution*. Edinburgh: Churchill Livingstone.
672. Yip, Y., & Tse, S. (2004). The effectiveness of relaxation acupoint stimulation and acupressure with aromatic lavender essential oil for non-specific low back pain in Hong Kong: A randomised controlled trial. *Complement Ther Med*, 12(1), 28-37.
673. Zaproudina, N., Hietikko, T., Hinninen, O., & Airaksinen, O. (2009). Effectiveness of traditional bone setting in treating chronic low back pain: A randomised pilot trial. *Complement Ther Med*, 17(1), 23-8.
674. Zhang, J., & Chen, W. (2004). Curative effect of nonoperative therapy for the lumbar disc herniation. *Chin J Clin Rehabil*, 8, 2006-7.
- Fibromyalgia**
675. Akbayrak T, Akarcali I, Karabudak R, Demirtük F. (2002). The results of connective tissue manipulation in the treatment of tension type headache. *Pain Clin*, 13, 343-7.
676. Aksu S, Asli K, Tülin D, et al. (2003). Are results of the SF-36 health survey and the Nottingham Health Profile similar in migraine patients? *Pain Clin*, 1-5.
677. Al-Allaf W, Khan F, Moreland J, Belch JF, Pullar T. (2001). Investigation of cutaneous microvascular activity and flare response in patients with fibromyalgia syndrome. *Rheumatology*, 40, 1097-101.
678. Alnigenis, M., Bradley, J., Wallick, J., & Emsley, C. (2001). Massage therapy in the management of fibromyalgia: A pilot study. *J Musculoskelet Pain*, 9(2), 55-67.
679. Asplund R. (2003). Manual lymph drainage therapy using light massage for fibromyalgia sufferers: a pilot study. *Orthop Nurs*, 7, 192-6.
680. Baldry, P., Yunus, M., & Inanici, F. (2001). *Myofascial pain and fibromyalgia syndromes*. Edinburgh: Churchill Livingstone.
681. Barker, K. (2005). *The fibromyalgia story*. Philadelphia: Temple University Press.
682. Bennett, R. (2005). Fibromyalgia: Present to future. *Current Rheumatology Reports*, 7(5), 371-6.
683. Bennett R. (2005). The Fibromyalgia Impact Questionnaire (FIQ): a review of its development, current version, operating characteristics and uses. *Clin Exp Rheumatol*, 5, 154-62.
684. Bieber C, Müller KG, Blumenstiel K, et al. (2006). Long-term effects of a shared decision-making intervention on physician-patient interaction and outcome in fibromyalgia: a qualitative and quantitative 1 year follow-up of a randomized controlled trial. *Patient Educ Couns*, 63, 357-66.
685. Bliddal H, Danneskiold-Samsøe B. (2007). Chronic widespread pain in the spectrum of rheumatological diseases. *Best Pract Res Clin Rheumatol*, 21, 391-402.
686. Bradley, L. (2008). Family and genetic influences on fibromyalgia syndrome. *J Musculoskelet Pain*, 16(1-2), 49-57.
687. Brattberg, G. (1999). Connective tissue massage in the treatment of fibromyalgia. *Eur J Pain*, 3(3), 235-44.
688. Burckhardt, C., Clark, S., & Bennett, R. (1991). The fibromyalgia impact questionnaire: Development and validation. *J Rheumatol*, 18(5), 728-33.
689. Busch, A., Thille, P., Barber, K., Schachter, C., Bidonde, J., & Collacott, B. (2008). Best-practice: E-model - prescribing physical activity and exercise for individuals with fibromyalgia. *PTP* 24(3), 151-66.
690. Buskila, D., Neumann, L., Vaisberg, G., Alkalay, D., & Wolfe, F. (2005). Increased rates of fibromyalgia following cervical spine injury: A controlled study of 161 cases of traumatic injury. *J Arthritis Rheum*, 40(3), 446-52.
691. Castro-Sanchez, A., Mataran-Penarrocha, G., Granero-Molina, J., Aguilera-Manrique, G., Quesada-Rubio, J., & Moreno-Lorenzo, C. (2011). Benefits of massage-myofascial release therapy on pain, anxiety, quality of sleep, depression, and quality of life in patients with fibromyalgia. *Evid Based Complement Alternat Med*, Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/21234327>
692. "Myofascial release techniques improved pain and quality of life in patients with fibromyalgia."
693. Castro-Sanchez, A., Mataran-Penarrocha, G., Granero-Molina, J., Aguilera-Manrique, G., Quesada-Rubio, J., & Moreno-Lorenzo, C. (2011). Effects of myofascial release techniques on pain, physical function, and postural stability in patients with fibromyalgia: a randomized controlled trial. *Clin Rehabil*, 25(9), 800-13. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/21673013>

694. "The results suggest that myofascial release techniques can be a complementary therapy for pain symptoms, physical function and clinical severity but do not improve postural stability in patients with fibromyalgia syndrome."
695. Çıtak-Karakaya İ, Akbayrak T, Demirtürk F, Ekici G, Bakar Y. (2006). Short- and long-term results of connective tissue manipulation and combined ultrasound therapy in patients with fibromyalgia. *J Manipulative Physiol Ther*, 29, 524-8.
696. Desmeules, J., Cedraschi, C., Rapiti, E., Baumgartner, E., Finckh, A., Cohen, P., Dayer, P., & Vischer, T. (2003). Neurophysiologic evidence for a central sensitization in patients with fibromyalgia. *Arthritis Rheum*, 48, 1420-9
697. Ebner M. (1978). Connective tissue massage. *Physiotherapy*, 64, 208-10.
698. Ekici, G., Bakar, Y., Akbayrak, T., & Yuksel, I. (2009). Comparison of manual lymph drainage and connective tissue massage in women with fibromyalgia: A randomized controlled trial. *J Manipulative Physiol met*, 32(2), 127-33.
699. Field, T., Diego, M., Cullen, C., Hernandez-Reif, M., Sunshine, W., & Douglas, S. (2002). Fibromyalgia pain and substance p decrease and sleep improves after massage therapy. *J Clin Rheumatol*, 8, 72-76.
700. Fischer AA. (1987). Pressure algometry over normal muscles. Standard values, validity and reproducibility of pressure threshold. *Pain*, 30(1), 115-26.
701. Földi M, Kubik S. (2001). *Anatomie des Lymphgefäßsystems, 'Lehrbuch der Lymphologie'*. München: Urban and Fischer, p. 4-37.
702. Fontaine, K., & Haaz, S. (2007). Effects of lifestyle physical activity on health status, pain and function, in adults with fibromyalgia syndrome. *J Musculoskeletal Pain*, 15(1), 3-9.
703. Fryer G, Hodgson L. (2005). The effect of manual pressure release on myofascial trigger points in the upper trapezius muscle. *J Bodywork Mov Ther*, 9, 248-55.
704. Geisser, M., Glass, J., Rajcevska, L., Clauw, D., Williams, D., Kileny, P., & Gracely, R. (2008). A psychophysiological study of auditory and pressure sensitivity in patients with fibromyalgia and healthy controls. *J Pain*, 9(5), 417-22.
705. Giovengo, S., Russell, J., & Larson, A. (1999). Increased concentration of nerve growth factor in cerebrospinal fluid of patients with fibromyalgia syndrome. *J Rheumatol*, 26, 1564-9.
706. Goats GC, Keir KA. (1991). Connective tissue massage. *Br J Sports Med*, 25, 131-3.
707. Goldenberg, D., Burckhardt, C., & Crafford, L. (2004). Management of fibromyalgia syndrome. *JAMA* 292, (19), 2388-95.
708. Guedj, E. (2008). Clinical correlate of brain SPECT perfusion abnormalities in fibromyalgia. *J Nucl Med* 49, (11), 1798-803.
709. Harris, R., Clauw, D., Scott, D., McLean, S., Gracely, R., & Zubieta, J. (2007). Decreased central m-opioid receptor availability in fibromyalgia. *J Neurosci*, 27(31), 10000-6.
710. Henriksson KG, Sörensen J. (2002). The promise of N-methyl-Laspartate receptor antagonists in fibromyalgia. *Rheum Dis Clin North Am*, 28, 343-51.
711. Hidalgo J, Rico-Villademoros F, Calandre EP. (2007). An open-label study of quetiapine in the treatment of fibromyalgia. *Prog Neuropsychopharmacol Biol Psychiatry*, 31, 71-7.
712. Hinds, T., McEwan, I., Perkes, J., Dawson, E., Ball, D., & George, K. (2004). Effects of massage on limb and skin blood flow after quadriceps exercise. *Med Sci Sports Exerc*, 36(8), 1308-13.
713. Holey EA. (2000).)Connective tissue massage: a bridge between complementary and orthodox approaches. *J Bodywork Mov Ther*, 4, 72-80.
714. Hurtig IM, Raak RI, Kendall SA, Gerdle B, Wahren LK. (2001). Quantitative sensory testing in fibromyalgia patients and in healthy subjects: identification of subgroups. *Clin J Pain*, 17, 316-22.
715. Kaada B, Torsteinbo O. (1989). Increase of plasma beta-endorphins in connective tissue massage. *Gen Pharmacol*, 20, 487-9.
716. Kosek, E., Eckholm, J., & Hansson, P. (1996). Modulation of pressure pain thresholds during and following isometric contraction in patients with fibromyalgia and in healthy normal controls. *Pain*, 64, 415-23.
717. Küçükdeveci AA, McKenna SP, Kutlay S. (2000). The development and psychometric assessment of the Turkish version of the Nottingham Health Profile. *Clin Rehab*, 23, 31-8.
718. Kurz I. (1989).)Textbook of Dr. Vodder's manual lymph drainage, Vol. 2. *Therapy, 2nd ed.*
719. Ledingham, J., Doherty, S., & Doherty, M. (1993). Primary fibromyalgia syndrome: An outcome study. *Br SOC Rheumatology*, 32(2), 139-42.
720. Lemstra, M., & Olszynski, W. (2005). The effectiveness of multidisciplinary rehabilitation in the treatment of fibromyalgia: A randomized controlled trial. *Clin J Pain*, 21(2), 166-74.
721. Mannerkorpi K, Henriksson C. (2007). Non-pharmacological treatment of chronic widespread musculoskeletal pain. *Best Pract Res Clin Rheumatol*, 21, 513-34.
722. Martinez-Lavin M, Hermosillo AG. (2000).Autonomic nervous system dysfunction may explain the multisystem features of fibromyalgia. *Arthritis Rheum*, 29, 197-9.
723. Mataran-Penarrocha, G., Castro-Sanchez, A., Garcla, G., Moreno-Lorenzo, C., Carrefio, T., & Zafra, M. (2009). Influence of craniosacral therapy on anxiety, depression and quality of life in patients with fibromyalgia. Retrieved from <http://ecam.oxfordjournals.org/cgi/reprint/nep125yl>.
724. McKechnie AA, Wilson F, Watson N, Scott D. (1983). Anxiety states: a preliminary report on the value of connective tissue massage. *J Psychom Res*, 27, 125-9.
725. Moldofsky, H. (2008). The assessment and significance of the sleep/waking brain in patients with chronic widespread musculoskeletal pain and fatigue syndromes. *J Musculoskeletal Pain*, 16(1-2), 37-48.
726. Mori, H., Ohsawa, H., Tanaka, T., Taniwaki, E., Leisman, G., & Nishijo, K. (2004). Effect of massage on blood flow and muscle fatigue following isometric lumbar exercise. *Med Sci Monit*, 10(5), 173-8.

727. Moyer, C. (2007). Between-groups study designs demand between-groups analyses: A response to Hernandez-Reif, Shor-Posner, Baez, Soto, Mendoza, Castillo, Quintero, Perez, and Zhang. *Evid Based Complement Alternat Med*, 6(1), 49-50.
728. Neumann, L., & Buskila, D. (2003). Epidemiology of fibromyalgia. *Curr Pain Headache Rep*, 7, 362-8.
729. Offenbacher M, Stuck G. (2000). Physical therapy in the treatment of fibromyalgia. *Scand J Rheumatol*, 113, 78-85.
730. Persson AL, Hansson GA, Kalliomaki A, Moritz U, Sjolund BH. (2000). Pressure pain thresholds and electromyographically defined muscular fatigue induced by a muscular endurance test in normal women. *Clin J Pain*, 16, 155-63.
731. Price DD, McGrath PA, Rafii A, Buckingham B. (1983). The validation of visual analogue scales as ratio scale measures for chronic and experimental pain. *Pain*, 17, 45-56.
732. Reed B, Held J. (1988). Effects of sequential connective tissue massage on autonomic nervous system of middle aged and elderly adults. *Phys Ther*, 68:1231-4.
733. Rose KE, Taylor HM, Twycross R. (1991). Long-term compliance with treatment in obstructive arm lymphoedema in cancer. *Palliat Med*, 5, 52-5.
734. Russell, J. (2001). Fibromyalgia syndrome. *Bonica's Management of Pain, 3rd ed., ed. J.D. Loeser, S.H. Butler, C.R. Chapman, and D.C. Turk*. Philadelphia: Lippincott Williams and Wilkins.
735. Sarmer S, Ergin S, Yavuzer G. (2000). The validity and reliability of the Turkish version of the Fibromyalgia Impact Questionnaire. *Rheumatol Int*, 20, 9-12.
736. Sandberg M, Larsson B, Lindberg L, Gerdle B. (2005). Different patterns of blood flow response in the trapezius muscle following needle stimulation (acupuncture) between healthy subjects and patients with fibromyalgia and work-related trapezius myalgia. *Eur J Pain*, 9, 497-510.
737. Schliac H. (1978). *Theoretical basis of working mechanism of connective tissue massage*. In: Dicke E, Shliak H, editors. A manual reflexive therapy of the connective tissue. New York: Sidney Simon Publishers; p. 14-33.
738. Sim J, Adams N. (1999). Physical and other non-pharmacological interventions for fibromyalgia. *Baillieres Best Pract Res Clin Rheumatol*, 13, 507-23.
739. Sim, J., & Adams, N. (2002). Systematic review of randomized controlled trials of nonpharmacological interventions for fibromyalgia. *Clin J Pain*, 18(5), 324-6.
740. Soden, K., Vincent, K., Craske, S., Lucas, C., & Ashley, S. (2004). A randomized controlled trial of aromatherapy massage in a hospice setting. *Palliat Med*, 18(2), 87-92.
741. Staud R. (2006). Biology and therapy of fibromyalgia: pain in fibromyalgia syndrome. *Arthritis Res Ther*, 8, 208.
742. Staud, R. (2008). The role of peripheral input for chronic pain syndromes like fibromyalgia syndrome. *J Musculoskeletal Pain*, 16(1-2), 74-84.
743. Staud R, Domingo M. (2001). Evidence for abnormal pain processing in fibromyalgia syndrome. *Pain Med*, 2, 208-15.
744. Staud, R., Cannon, R., Mauderli, A., Robinson, M., Price, D., & Vierck, C., Jr. (2003). Temporal summation of pain from mechanical stimulation of muscle tissue in normal controls and subjects with fibromyalgia syndrome. *Pain*, 102, 87-95.
745. Staud R, Price DD, Robinson ME, Vierck CJ. (2004). Bodily pain area and pain-related negative affect predict clinical pain intensity in patients with fibromyalgia. *J Pain*, 5, 338-43.
746. Strobel, E., Krapf, M., Suckfull, M., Bruckle, W., Fleckenstein, W., & Muller, W. (1997). Tissue oxygen measurement and 31P magnetic resonance spectroscopy in patients with muscle tension and fibromyalgia. *Rheumatol Int*, 16, 175-80.
747. Sunshine, W., Field, T., Quintino, O., Fierro, K., Kuhn, C., Burman, I., & Schanberg, S. (1996). Fibromyalgia benefits from massage therapy and transcutaneous electrical stimulation. *J Clin Rheumatol*, 2(1), 18-22.
748. Terry, R., Perry, R., & Ernst, E. (2012). An overview of systematic reviews of complementary and alternative medicine for fibromyalgia. *Clin Rheumatol*, 32(1), 55-66. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/21614472>
749. "The reviews found some evidence of beneficial effects arising from acupuncture, homoeopathy, hydrotherapy and massage, whilst no evidence for therapeutic effects from chiropractic interventions for the treatment of FM symptoms was found."
750. Theadom A, Cropley M, Humphrey KL. (2007). Exploring the role of sleep and coping in quality of life in fibromyalgia. *J Psychosom Res*, 62, 145-51.
751. Thieme, K., Turk, D., & Flor, H. (2004). Comorbid depression and anxiety in fibromyalgia syndrome: Relationship to somatic and psychosocial variables. *Psychosom Med*, 66, 837-44.
752. Vernon H, Kim Humphreys K, Hagino C. (2007). Chronic mechanical neck pain in adults treated by manual therapy: a systematic review of change scores in randomized clinical trials. *J Manipulative Physiol Ther*, 30, 215-27.
753. Wickelgren, I. (2009). I do not feel your pain. *Scientific American Mind*, 20(5);50-7.
754. Wolfe F, Smythe H, Yunus M, Bennett R, Bombardier C, Goldenberg D, et al. (1990) The American College of Rheumatology 1990 criteria for the classification of fibromyalgia. Report of the multicenter criteria committee. *Arthritis Rheum*, 33, 160-72.
755. Wolfe F, Anderson J, Harkness D, et al. (1997). Health status and disease severity in fibromyalgia: results of a six-center longitudinal study. *Arthritis Rheum*, 40, 1571-9.
756. Zautra, A., Fasman, R., Davis, M., & Craig, A. (2010). The effects of slow breathing on affective responses to pain stimuli: An experimental study. *Pain*, 149(1), 12-8.

Cancer

757. 2009. Massage therapy for cancer palliation and supportive care: A systematic review of randomised clinical trials. *Support Care Cancer* 17(4): 333 -337.

758. Billhult, A., Bergborn, I., & Stener-Victorin, E. (2007). Massage relieves nausea in women with breast cancer who are undergoing chemotherapy. *J Altern Complem Med*, 13(1), 53-7.
759. Billhult, A., Lindholm, C., Gunnarsson, R., & Stener-Victorin, E. (2009). The effect of massage on immune function and stress in women with breast cancer: A randomized controlled trial. *Auton INeurosci-Basic*, 150(1-2), 111-5.
760. Boon, H., Olatunde, F., & Zick, S. (2007). Trends in complementary/alternative medicine use by breast cancer survivors: Comparing survey data from 1998 and 2005. *BMC Women 's Health*, 7(4). Retrieved from www.biomedcentral.com/ 1472-6874174.
761. Brennan, M. J., DePompolo, R. W., & Garden, F. H. (1996). Focused review: Postmastectomy lymphedema. *Archives of Physical Medicine and Rehabilitation*, 77, s74-s80.
762. Campbell, S., Louie-Gao, Q., Hession, M., Bailey, E., Geller, A., & Cummins, D. (2012). Skin cancer education among massage therapists: A survey at the 2010 meeting of the American Massage Therapy Association. *J Canc Educ*, doi: 10.1007/s13187-012-0403-7
763. "...massage therapists routinely have the opportunity to examine high risk anatomic sites, such as the back or legs, on many patients....massage therapists can play an important role in the early detection of skin cancer."
764. Canadian Cancer Society/National Cancer Institute of Canada: Canadian Cancer Statistics 2006, Toronto, Canada, 2006.
765. Cassileth B., & Vickers, A. (2004). Massage therapy for symptom control: Outcome study at a major cancer center. *J Pain Symptom Manag*, 28(0), 244-9.
766. Corbin, L. (2005). Safety and efficacy of massage therapy for patients with cancer. *Cancer Control*, 12, 158-64.
767. 2009. *Massage therapy*. In Integrative Oncology, ed. Donald I. Abrams and Andrew T. Weil, 232-43. New York: Oxford University Press.
768. Corner, J., Cawley, N., & Hildebrand, S. (1995). An evaluation of the use of massage and essential oils on the well being of cancer patients. *Int J Palliat Nur I*, (2), 67-73.
769. Currin, J., & Meister, E. (2008). A hospital-based intervention using massage to reduce distress among oncology patients. *Cancer Nursing*, 31(3), 214-22 1.
770. Datta YH, Adams PT, Drobyski WR, et al. Sensitive detection of occult breast cancer by the reverse-transcriptase polymerase chain reaction. *J Clin Oncol* 1994;12:475-82. .
771. Deng, G., Frenkel, M., Cohen, L., Cassileth, B., Abrams, D., Capodice, J., & Sagar, S. (2009). Evidence-based clinical practice guidelines for integrative oncology: Complementary therapies and botanicals. *J Soc Integr Oncol*, 7(Q), 85120.
772. Dibble, S., Luce, J., Cooper, B., Israel, J., Cohen, M., Nussey, & Rugo, H., (2007). Acupressure for chemotherapy-induced nausea and vomiting: A randomized clinical trial. *Oncol Nur Forum*, 34(4), 813-20.
773. Eillhult, A., Lindholm, C., Gunnarsson, R., & Stener-Victorin, E. (2008). The effect of massage on cellular immunity, endocrine and psychological factors in women with breast cancer: A randomized controlled clinical trial. *Aufon Neumsci-Basic*, 140:8895.
774. Ernst, E. (2003). The safety of massage therapy. *Rheumatology* 42(9): 1101-1 106.
775. Fellowes, D., Barnes, K., & Wilkinson, S. (2008). Aromatherapy and massage for symptom relief in patients with cancer. *Cochrane Db Syst Rev* (4). CD002287.pub3.
776. Ferrell-Torry, A., & Glick, O. (1993). The use of therapeutic massage as a nursing intervention to modify anxiety and the perception of cancer pain. *Cancer Nursing*, 16(2), 93-101.
777. Fidler IJ, Krippe ML. Metastasis results from preexisting variant cells within a malignant tumor. *Science* 1977;197:893-5. .
778. Field, T., Cullen, M., Diego, M., Hernandez-Reif, P., Sprinz, K., Beebe, B., & Bango-Sanchez V. (2001). Leukemia immune changes following massage therapy. *J Bodyw Mov Thers*, 271-4.
779. Field, T., M. Hernandez-Reif, M. Diego, S. Schanberg, and C. Kuhn. 2005, Cortisol decreases and serotonin and dopamine increase following massage therapy. *Int J Neurosci* 115(10): 1397-1413.
780. Ganz, PA: Sexual functioning after breast cancer: A conceptual framework for future studies. *Ann. Oncol.* 8 (1997), 105-107.
781. Gecsed, R. (2002). Massage therapy for patients with cancer. *Clin J Oncol Nurs*, 6(1), 52-54.
782. Gelmon K, Trevisan C, Olivotto IA, et al. Primary care for survivors of breast cancer. *N Engl J Med* 2000;344:309-310.
783. Ghossein RA, Rosai J, Scher HI, et al. Prognostic significance of detection of prostate-specific antigen transcripts in the peripheral blood of patients with androgen-independent prostatic carcinoma. *Urology* 1997;50:100-5. .
784. Giuliano AE, Jones RC, Brennan M, Statman R. Sentinel lymphadenopathy in breast cancer. *J Clin Oncol* 1997;15:2345-50. .
785. Goodfellow, L. (2003). The effects of therapeutic back massage on psychophysiologic variables and immune function in spouses of patients with cancer. *Nurs Res*, 52(5), 318-28.
786. Grealish, L.O., A. Lomansey, and B. Whiteman. 2000. Foot massage: A nursing intervention. *Cancer Nursing* 23: 237-243.
787. Gunn J, McCall JL, Yun K, Wright PA. Detection of micrometastases in colorectal cancer patients by K19 and K20 reverse-transcription polymerase chain reaction. *Lab Invest* 1996;75:611-6. .
788. Harris SR, Hugi MR, Olivotto IA, Levine M: Clinical practice guidelines for the care and treatment of breast cancer: 11. Lymphedema. *Can Med Assoc J* 164: 191-199, 2001
789. Haun, J., Graham-Pole, J., & Shortley, B. (2009). Children with cancer and blood diseases experience positive physical and psychological effects from massage therapy. *Int . Ther Massage Bodywork*, 2(Q), 7-14.
790. Hayes S, Battistutta D, Newman B. Objective and subjective upper body function six months following diagnosis of breast cancer. *Breast Cancer Res Treat* 2005;94:1-10.
791. Hernandez-Reif, M., Ironson, G., Field, T., Hurley, J., K t z, G., Diego, M., & Burman, I. (2004). Breast cancer patients have improved immune and neuroendocrine functions following massage therapy. *J Psychosom Res*, 5, 45-52.

792. Hernandez-Reif, M., Field, T., Ironson, G., Beutler, J., Vera, Y., Hurley, J., & Fraser, M. (2005). Natural killer cells and lymphocytes increase in women with breast cancer following massage therapy. *Int J Neurosci*, *115*(4), 495-510.
793. Hughes, D., Ladas, E., Rooney, D., & Kelly, K. (2008). Massage therapy as a supportive care intervention for children with cancer. *Oncol Nurs Forum*, *35*(3), 431-42.
794. Jane, S., Wilkie, D., Gallucci, B., & Beaton, R. (2008). Systematic review of massage intervention for adult patients with cancer: A methodological perspective. *Cancer Nursing*, *31*(6), E24-35.
795. Karki A, Simonen R, Malkia E, Selve J: Efficacy of physical therapy methods and exercise after a breast cancer option: a systematic review. *Crit Rev Phys Rehabil Med* 13: 159–190, 2001
796. Kutner, J., Smith, M., Corbin, L., Hemphill, L., Bentan, K., Mellis, B., & Fairclough, D. (2008). Massage therapy versus simple touch to improve pain and mood in patients with advanced cancer. *Ann Intern Med*, *149*, 369-79.
797. Kwan W, Jackson J, Weir L, Dingee C, McGregor G, Olivotto I. Chronic arm morbidity after curative breast cancer treatment: prevalence and impact on quality of life. *J Clin Oncol* 2002;20:4242-8.
798. Lee, CO: Quality of life and breast cancer survivors. Psychosocial and treatment issues. *Cancer Pract*. 5 (1997), 309-316.
799. Leidenius M, Leppanen E, Krogerus L, von-Smitten K. Motion restriction and axillary web syndrome after sentinel node biopsy and axillary clearance in breast cancer. *Am J Surg* 2003;185:127-30.
800. Liotta LA, Stetler-Stevenson WG. Principles of molecular cell biology of cancer: cancer metastasis. In: DeVita VT, Hellman S. Rosenberg SA, editors. Principles and practice of oncology. 4th ed. Philadelphia: J.B. Lippincott Co; 1993. p. 134–49. .
801. Listing, M., Ueiphauer, M., Krohn, M., Voigt, B., Tjahono, G., Becker, J., Klapp, B., & Rauchfup, M. (2009). Massage therapy reduces physical discomfort and improves mood disturbances in women with breast cancer. *Psychooncology*, *18*(12), 1290-9.
802. Liu, Y., & Fawcett, T. (2008). The role of massage therapy in the relief of cancer pain. *Nursing Standard*, *22*(21), 35-40.
803. Lund, J, Y Ge, L. Yu, K. Uvnas-Moberg, J. Wang, C. Yu, M. Kurosawa, G. Agren, A. Rosen, M. Lekrnan, arid T Lundeberg. 2002. Repeated massage like stimulation induces long-term effects on nociception: Contribution of oxytocinergic mechanisms. *Eur J Neurosci* 16(2): 330-338. MacDonald, G.2007. *Medicine hands: Massage therapy for people with cancer* Foorres, UK: Findhorn Press.
804. MacDonald, G. (2007). *Medicine hands: Massage therapy for people with cancer*. Foorres, UK: Findhorn Press.
805. Myers, C., Walton, T., & Small, B. (2008). The value of massage therapy in cancer care. *Hematol Oncol Clin N*, *22*, 649-60.
806. Paget S. The distribution of secondary growths in cancer of the breast. *Lancet* 1889;571–3. .
807. Pelkey TJ, Frierson HF, Bruns DE. Molecular and immunological detection of circulating tumor cells and micrometastases from solid tumors. *Clin Chem* 1996;42:1369–82.
808. Petrek JA, Heelan MC. Incidence of breast carcinoma-related lymphedema. *Cancer* 1998;83 Suppl:2776–81. .
809. Post-White, J., Kinney, M., Savik, K., Gau, J., Wilcox, C., Lerner, I. (2003). Therapeutic massage and healing touch improve symptoms in cancer. *Integr Cancer Ther*, *2*(4): 332-44.
810. Post-White, J., Fitzgerald, M., Hageness, S., & Sencer, S. (2009). Complementary and alternative medicine use in children with cancer and general and specialty pediatrics. *J Pediatr Oncol Nurs*, *26*(1), 7-15.
811. Post-White, J., Fitzgerald, M., Savik, K., Hooke, M., Hannahan, A., & Sencer, S. (2009). Massage therapy for children with cancer. *J Pediatr Oncol*, *26*, 16-28.
812. Purushotham AD, Upponi S, Klevesath MB, et al. Morbidity after sentinel lymph node biopsy in primary breast cancer: results from a randomized controlled trial. *J Clin Oncol* 2005;23:4312–21. .
813. Rietman J, Dijkstra P, Debreczeni R, Geertzen J, Robinson D, Vries J. Impairments, disabilities and health related quality of life after treatment of breast cancer: a follow-up study 2.7 after surgery. *Disabil Rehabil* 2004;26:78-84.
814. Ruitler DJ, van Krieken JH, van Muijzen GN, de Waal RM. Tumour metastasis: is tissue an issue?. *Lancet Oncol* 2001;2:109–12. .
815. Russell, N., Surnler, S., Beinhorn, C., & Frenkel, M. (2008). Role of massage therapy in cancer care. *J Altern Complem Med*, *14*(2), 209-14.
816. Sagar, S., Dryden, T., & Myers, C. (2007). Research on therapeutic massage for cancer patients: Potential biologic mechanisms. *J Soc Integr Oncol*, *5*(4), 155-62.
817. Schoenfeld A, Kruger KH, Gomm J, et al. The detection of micrometastases in the peripheral blood and bone marrow of patients with breast cancer using immunohistochemistry and reverse transcription polymerase chain reaction for keratin 19. *Eur J Cancer* 1997;33:854–61. .
818. Shamley DR, Barker K, Simonite V, Beardshaw A. Delayed versus immediate exercises following surgery for breast cancer: a systematic review. *Br Cancer Res Treat* 2005;90:263-71.
819. Smith, M., Yamashita, T., Bryant, L., Hemphill, L., & Kutner, J. (2009). Providing massage therapy for people with advanced cancer: What to expect. *J Altern Complem Med*, *15*(4), 367-71.
820. Soden, K., K. Vincent, S. Craske, C. Lucas, and S. Ashley. 2004. A randomized controlled trial of aromatherapy massage in a hospice setting. *Palliat Med* 18(2): 87-92.
821. Soerjomataram I, Louwman W, Ribot J, Roukema J, Coebergh J. An overview of prognostic factors for long-term survivors of breast cancer. *Breast Cancer Res Treat* 2008;107:309-30.
822. Stephenson, N., Swanson, M., Dalton, J., Keefe, F., & Engelke, M. (2007). Partner-delivered reflexology: Effects on cancer pain and anxiety. *Oncol Nurs Forum*, *34*(1), 127-32.
823. Sturgeon, M., Wetta-Hall, R., Hart, T., Good, M., & Dakhil, S. (2009). Effects of therapeutic massage on the quality of life among patients with breast cancer during treatment. *J Altern Complem Med*, *15*(4), 373-80.

824. Tarin D, Price JE, Kettlewell MGW, et al. Mechanisms of human tumor metastasis studied in patients with peritoneovenous shunts. *Cancer Res* 1984;44:3584–92. .
825. Tengrup I, Tennvall-Nittby L, Christiansson I, Laurin M. Arm morbidity after breast-conserving therapy for breast cancer *Acta Oncol* 2000;39:393-7.
826. Torres Lacomba M, Mayoral del Moral O, Coperias Zazo JL, Ferrandez JC, Zapico Goñi A. Axillary web syndrome after axillary dissection in breast cancer: a prospective study. *Breast Cancer Res Treat* 2009;doi:10.1007/s10549-009-0371-8.
827. Van Poll D, Thompson JF, Colman MH, et al. A sentinel node biopsy does not increase the incidence of in-transit metastasis in patients with primary cutaneous melanoma. *Ann Surg Oncol* 2005;12:5597–608. .
828. Weinich, S., & Weinrich, M. (1990). The effect of massage on pain in cancer patients. *Appi Nurs Res*, 3, 140-45.
829. Weir H, Thun M, Hankey B, Ries L. Annual report to the nation on the status of cancer 1975-2000, featuring the uses of surveillance data for cancer prevention and control. *J Natl Cancer Inst* 2003;95:1276-98.
830. Wilkie, D., Kampbell, J., Cutdshall, S., Halabisky, H., Harmon, H., Johnson, L., Weinacht, L., & RakeMarona, M. (2000). Effects of massage on pain intensity, analgesics, and quality of life in patients with cancer pain. *Hospice J*, 15, 31-53.
831. Wilkinson, S., Love, S., Westcornbe, A., Gambles, M., Burgess, C., Cargill, A., Young, T., Maher, E., & Ramirez, A. (2007). Effectiveness of aromatherapy massage in the management of anxiety and depression in patients with cancer: A multicenter randomized controlled trial. *J Clin Oncol*, 25(5), 532-39.
832. Wilkinson, S., Barnes, K., & Storey L. (2008). Massage for symptom relief in patients with cancer: Systematic review. *J Adv Nurs*, 63(5), 430-39.
833. Williams, A., Vadgama, A., Franks, P., & Mortimer, P. (2002). A randomized controlled crossover study of manual lymphatic drainage therapy in women with breast cancer related lymphoedema. *Eur J Cancer Care*, 1(4), 254-61.
834. Woods, M, M Tobin, P Mortimer: The psychosocial morbidity of breast cancer with lymphoedema. *Cancer Nurs*. 18 (1995), 467-471.

TMJD

835. Capellini, V., Souza, G., & Faria, C. (2006). Massage therapy in the management of myogenic tmd: A pilot study. *J Appl Oral Sci*, 14(1), 21-6. Retrieved from www.fob.usp.br/revista or www.scielo.br/jaos
836. DeBar, L., Vuckovic, N., Schneider, J., et al. (2003). Use of complementary and alternative medicine for temporomandibular disorders. *Journal of Orofacial Pain*, 17(3), 224–36.
837. Eisensmith, L. (2007). Massage therapy decreases frequency and intensity of symptoms related to temporomandibular joint syndrome in one case study. *Journal of Bodywork and Movement Therapies*, 11, 223-30. Retrieved from www.intl.elsevierhealth.com/journals/jbmt
838. Kalamir, A., Chiro, M., Pollard, H., Vitiello, A., Chiro, M., & Bonello, R. (2007). Manual therapy for temporomandibular disorders: A review of the literature. *Journal of Bodywork and Movement Therapies*, 11, 84-90.
839. Michelotti, A., Parisini, F., Farella, M., et al. (2000). Muscular physiotherapy in patients with temporomandibular disorders. Controlled clinical trial. *Minerva Stomatologica*, 49(11–12), 541–8.
840. Nicolakis, P., Burak, E., Kollmitzer, J., et al. (2001). An investigation of the effectiveness of exercise and manual therapy in treating symptoms of TMJ osteoarthritis. *Journal of Craniomandibular Practice*, 19(1), 26–32.
841. Nicolakis, P., Erdogmus, C., Kollmitzer, J., et al. (2002). Longterm outcome after treatment of temporomandibular joint osteoarthritis with exercise and manual therapy. *Journal of Craniomandibular Practice*, 20(1), 23–7.
842. van der Glas, H., Buchner, R., van Grootel, R., (2000). Comparison of treatment options for myogenous temporomandibular dysfunction. *Nederlands Tijdschrift voor Tandheelkunde*, 107(12), 505–12.

Osteoarthritis

843. Arden N., & Nevitt M. (2006). Osteoarthritis: epidemiology. *Best Pract Res Clin Rheumatol*, 20, 3-25.
844. Buckwalter J., Saltzman C., & Brown T. (2004). The impact of osteoarthritis: implications for research. *Clin Orthop*, 427(suppl), S6-S15.
845. Butler, R., Barrios, J., Royer, T., & Davis, I. (2011). Frontal-plane gait mechanics in people with medial knee osteoarthritis are different from those in people with lateral knee osteoarthritis. *Physical Therapy*, 91(8), 1235-43. doi: 10.2522/ptj.20100324
846. Elders M. (2000). The increasing impact of arthritis on public health. *J Rheumatol*, 60(suppl), 6-8.
847. Ernst E. (2002). Complementary and alternative medicine for pain management in rheumatic disease. *Curr Opin Rheumatol*, 14, 58-62.
848. Felson D. (2004). An update on the pathogenesis and epidemiology of osteoarthritis. *Radiol Clin North Am*, 42, 1-9.
849. Grotle, M., Garratt, A., Klokkerud, M., Lochting, I., Uhlig, T., & Hagen, K. (2010). What's in team rehabilitation care after arthroplasty for osteoarthritis? Results from a multicenter, longitudinal study assessing structure, process and outcome. *Physical Therapy*, 90, 121-31. doi: 10.2522/ptj.20080295
850. Kato T., Xiang Y., Nakamura H., & Nishioka K. (2004). Neoantigens in osteoarthritic cartilage. *Curr Opin Rheumatol*, 16, 604-8.
851. Katz, W. (2007). Themed review: Nonpharmacologic approaches to osteoarthritis. *American Journal of Lifestyle Medicine*, 1(249), doi: 10.1177/1559827607301573.
852. Lawrence R., Helmick C., Arnett F., et al. (1998). Estimates of the prevalence of arthritis and selected musculoskeletal disorders in the United States. *Arthritis Rheum*, 41, 778-99.

853. Lund, H., Henriksen, M., Bartels, E., Danneskiold-Samsøe, B., & Bliddal, H. (n.d.). Can stimulating massage improve joint repositioning error in patients with knee osteoarthritis?. *Journal of Geriatric Physical Therapy*, 32(3).
854. Matchaba P., Gitton X., Krammer G., et al. (2005). Cardiovascular safety of lumiracoxib: a meta-analysis of all randomized controlled trials. 1 week and up to 1 year in duration of patients with osteoarthritis and rheumatoid arthritis. *Clin Ther*, 27, 1196-214.
855. Naesdal J., & Brown K. (2006). NSAID-associated adverse effects and acid control aids to prevent them: A review of current treatment options. *Drug Saf*, 29, 119-32.
856. Perlman, A., Sabina, A., Williams, A., Yanchou, V., & Katz, D. (2006). Massage therapy for osteoarthritis of the knee. *Arch Intern Med*, 166.
857. Preyde M. (2000). Effectiveness of massage therapy for subacute low-back pain: A randomized controlled trial. *CMAJ*, 162, 1815-20.

Pregnancy

858. Agren, A., & Berg, M. (2006). Tactile massage and severe nausea and vomiting during pregnancy: Women's experiences. *Scand J Caring Sci*, 20, 169-76.
859. Barnes, P., Bloom, B., & Nahin, R. (2008). Complementary and alternative medicine use among adults and children: United States, 2007. *Natl Health Stat Report*, 12, 1-24.
860. Billhult, A., Bergborn, I., & Stener-Victorin, E. (2004). Massage relieves nausea in women with breast cancer who are undergoing chemotherapy. *JA Altern Complem Med*, 13, 53-7.
861. Chang, M., Wang, S., & Chen, C. (2002). Effects of massage on pain and anxiety during labor: A randomized controlled trial in Taiwan. *J Adv Nurs*, 38, 68-73.
862. (2006). A comparison of massage effects on labor pain using the McGill pain questionnaire. *J Nun Res*, 14, 190-6.
863. Cookson, H., Granell, R., Joinson, C., BenShlomo, Y., & Henderson, A. (2009). Mother's anxiety during pregnancy is associated with asthma in their children. *J Allergy Clin Immun*, 123, 847- 53.
864. Diego, M., Field, T., Hernandez-Reif, M., Schanberg, S., Kuhn, C., & Gonzalez-Quintero, V. (2009). Prenatal depression restricts fetal growth. *Early Hum Dev*, 85, 65-70.
865. Ernst, E. (2003). The safety of massage therapy. *Rheumatol*, 42, 1101-6.
866. Field, T., Hernandez-Reif, M., Taylor, S., Quintino, O., & Burman, I. (1997). Labor pain is reduced by massage therapy. *J Psychosom Obst Gyn*, 18, 286-91.
867. Field, T., Hernandez-Reif, M., Hart, S., Theakston, H., Schanberg, S., & Kuhn, C. (1999). Pregnant women benefit from massage therapy. *J Psychosom Obst Gyn*, 20, 31-8.
868. Field, T., Diego, M., Hernandez-Reif, M., Schanberg, S., & Kuhn, C. (2004). Massage therapy effects on depressed pregnant women. *J Psychosom Obst Gyn*, 15, 115-22.
869. Field, T., Figueiredo, B., Hernandez-Reif, M., Diego, M., Deeds, O., & Ascencio, A. (2008). Massage therapy reduces pain in pregnant women, alleviates prenatal depression in both parents and improves their relationship. *J Bodyw Mov Ther*, 12, 146-50.
870. Field, T., Diego, M., Hernandez-Reif, M., Deeds, O., & Figueiredo, B. (2009). Pregnancy massage reduces prematurity, low birthweight and postpartum depression. *Infant Behav Dev*: Epublication.
871. Huntley, A., Thompson Coon, J., & Ernst, E. (2004). Complementary and alternative medicine for labor pain: A systematic review. *Am J Obstet Gynecol*, 191, 36-44.
872. Kosfeld, M., Heinrichs, M., Zak, P., Fischbacher, U., & Fehr, E. (2005). Oxytocin increases trust in humans. *Nature*, 435, 673-6
873. McCain, G., & Deatrick, J. (1994). The experience of high-risk pregnancy. *JOGNN*, 23, 421-7.
874. Melzack, R., & Wall, P. (1965). Pain mechanisms: A new theory. *Science*, 150, 971-9.
875. Mens, J., Vleeming, A., Stoeckart, R., Stam, H., & Snijders, C. (1996). Understanding peripartum pelvic pain: Implications of a patient survey. *Spine*, 21, 1363-9.
876. Moyer, C., Rounds, J., & Hannum, J. (2004). A meta-analysis of massage therapy research. *Psych Bull*, 130, 3-18.
877. Moyer, C., Seefeldt, L., Mann, E., & Jackley, L. (2011). Does massage therapy reduce cortisol? A comprehensive quantitative review. *J Bodyw Mov Ther*, 15, 3-14.
878. Pettigrew, A., O'Brien King, M., McGee, K., & Rudolph, C. (2004). Complementary therapy use by women's health clinic clients. *Altern Ther*, 10, 50-5.
879. Ven den Bergh, B., Mulder, E., Mennes, M., & Glover, V. (2005). Antenatal maternal anxiety and stress and the neurobehavioural development of the fetus and child: Links and possible mechanisms: A review. *Neurosci Biobehav R*, 29, 237-58.
880. Wang, S., DeZinno, P., Fermo, L., William, K., Caldwell-Andrews, A., Bravemen, F., & Kain, Z. (2002). Complementary and alternative medicine for low-back pain in pregnancy: A cross-sectional survey. *J Altern Complem Med*, 11, 459- 64.

Pediatrics

881. Adams, D., midden, A., Smith, K., Sikora, S., Dryden, T., & Vohra, D. (2009). Safety of pediatric massage: A systematic review. *Altern Ther Med*, 15(3), s135.
882. Barnes, P., Bloom, B., & Nahin, R. (2008). Complementary and alternative medicine use among adults and children: United States, 2007. *Natl Health Stat Report*, 12, 1-23.d
883. Beider, S., & Moyer, C. (2007). Randomized controlled trials of pediatric massage: A review. *Evid-Based Compl Alt*, 4(1), 23-34
884. Beider, S., Mahrer, N., & Gold, J. (2007). Pediatric massage therapy: An overview for clinicians. *Pediatr Clin N Am*, 54(6), 1025-41.

885. Canadian Cancer Society's Steering Committee. (2009). *Canadian cancer statistics 2009*. Toronto, ON: Canadian Cancer Society.
886. Centers for Disease Control and Prevention. (2010). *Autism spectrum disorders (ASDs)*. Retrieved from www.cdc.gov/ncbddd/autism/research.html.
887. Cullen-Powell, L., Barlow, J., & Cushway, D. (2005). Exploring a massage intervention for parents and their children with autism: The implications for bonding and attachment. *J Child Health Care, 9*(4), 245-55.
888. Diego, M., Field, T., Hernandez-Reif, M., Shaw, J., Rothe, E., Castellanos, D., & Mesner, L. (2002). Aggressive adolescents benefit from massage therapy. *Adolescence, 37*(147), 597-607.
889. Diego, M., Field, T., Hernandez-Reif, M., Deeds, O., Ascencio, A., & Begert, G. (2007). Preterm infant massage elicits consistent increases in vagal activity and gastric motility that are associated with greater weight gain. *Acta Paediatr, 96*(11), 1588-91.
890. Diego, M., Field, T., & Hernandez-Reif, M. (2009). Procedural pain heart rate responses in massaged preterm infants. *Infant Behav Dev, 32*(2), 226-9.
891. Escalona, A., Field, T., Singer-Strunck, R., Cullen, C., & Hartshorn, K. (2001). Brief report: Improvements in the behavior of children with autism following massage therapy. *J Autism Dev Disord, 31*(5): 513-6.
892. Esmail, N. (2007). Complementary and alternative medicine in Canada: Trends in use and public attitudes, 1997-2006. In *Public Policy Sources*, ed. K. McCahon. Vancouver, BC: Fraser Institute.
893. Field, T., Schanberg, A., Scafidi, F., Bauer, C., Vega-Lahr, N., Garcia, R., Nystrom, J., & Kuhn, C. (1986). Tactile/kinesthetic stimulation effects on preterm neonates. *Pediatrics, 77*(5), 654-8.
894. Field, T., Morrow, C., Valdeon, C., Larson, S., Kuhn, C., & Schanberg, S. (1992). Massage reduces anxiety in child and adolescent psychiatric patients. *J Am Acad Child Adolesc Psychiatry, 31*(1), 125-31.
895. Field, T., Quintino, O., Hernandez-Reif, M., & Koslovsky, G. (1998). Adolescents with attention deficit hyperactivity disorder benefit from massage therapy. *Adolescence, 33*(129), 103-8.
896. Field, T., Henteleff, T., Hernandez-Reif, M., Martinez, E., Mavunda, K., Kuhn, C., & Schanberg, S. (1998). Children with asthma have improved pulmonary functions after massage therapy. *J Pediatr, 132*(5), 854-8.
897. Field, T., Hernandez-Reif, M., Diego, M., Schanberg, S., & Kuhn, C. (2005). Cortisol decreases and serotonin and dopamine increase following massage therapy. *Int J Neurosci, 115*(10), 1397-413.
898. Garner, B., Phillips, L., Schmidt, H., Markulev, C., O'Connor, J., Wood, S., Berger, G., Bumett, P., & McGorry, P. (2008). Pilot study evaluating the effect of massage therapy on stress, anxiety and aggression in a young adult psychiatric inpatient unit. *Aust N Z J Psychiatry, 42*(5), 414-22.
899. Gonzalez, A., Vasquez-Mendoza, G., Garcia-Vela, A., Guzman-Ramirez, A., Salazar-Torres, M., & Rornerffutierrez, G. (2009). Weight gain in preterm infants following parent administered Vimala massage: A randomized controlled trial. *Am J Perinatal, 26*(4), 247-52.
900. Gray, L., Watt, L., & Blass, E. (2000). Skin-to-skin contact is analgesic in healthy newborns. *Pediatrics, 105*(1), e14.
901. Guzzetta, A., Baldini, S., Bancalè, A., Baroncelli, L., Ciucci, F., Ghirri, P., & Maffei, L. (2009). Massage accelerates brain development and the maturation of visual function. *J Neurosci, 29*(18), 6042-51.
902. Harlow, H. (1958). The nature of love. *Am Psychol, 13*(12), 673-85.
903. Harlow, H., & Suomi, S. (1970). Nature of love--Simplified. *Am Psychol, 25*(2), 161-8.
904. Haun, J., Graham-Pole, J., & Shortley, B. (2009). Children with cancer and blood diseases experience positive physical and psychological effects from massage therapy. *Int J Ther Massage Bodywork, 2*(2), 7-14.
905. Hernandez-Reif, M., Aeld, T., Krasnegor, J., Martinez, E., Schwartzman, M., & Mavunda, K. (1999). Children with cystic fibrosis benefit from massage therapy. *J Pediatr Psychol, 24*(2), 175-81.
906. Hernandez-Reif, M., Field, T., Lergie, S., Hart, S., Redzepi, M., Nierenberg, B., & Peck, T. (2001). Children's distress during burn treatment is reduced by massage therapy. *J Burn Care Rehabil, 22*(2), 191-5; discussion 190.
907. Hernandez-Reif, M., Diego, M., & Field, T. (2007). Preterm infants show reduced stress behaviors and activity after 5 days of massage therapy. *Infant Behav Dev, 30*(4), 557-61.
908. Hertenstein, M. (2002). Touch: Its communicative functions in infancy. *Hum Dev, 45*(2), 70-94.
909. Hertenstein, M., & Campos, J. (2001). Emotion regulation via maternal touch. *Infancy, 2*(4), 549-66.
910. Hughes, D., Ladas, E., Rooney, D., & Kelly, K. (2008). Massage therapy as a supportive care intervention for children with cancer. *Oncol Nurs Forum, 35*(3), 431-42.
911. Inui, K., Tsuji, T., & Kakigi, R. (2006). Temporal analysis of cortical mechanisms for pain relief by tactile stimuli in humans. *Cereb Cortex, 16*(3), 355-65.
912. Iwasaki, M. (2005). Interventional study on fatigue relief in mothers caring for hospitalized children: Effect of massage incorporating techniques from oriental medicine. *Kurume Med J, 52*, 19-27.
913. Kelmanson, I., & Adulas, E. (2006). Massage therapy and sleep behaviour in infants born with low birth weight. *Complement Ther Clin Pract, 12*(3), 200-5.
914. Khilnani, S., Field, T., Hernandez-Reif, M., & Schanberg, S. (2003). Massage therapy improves mood and behavior of students with attention deficit/hyperactivity disorder. *Adolescence, 38*(152), 623-38.
915. Maddigan, B., Hodgson, P., Heath, P., Dick, B., St. John, K., McWilliam-Burton, T., & White, H. (2003). The effects of massage therapy and exercise therapy on children/adolescents with attention deficit hyperactivity disorder. *Can Child Adolesc Psychiatr Rev, 12*(2), 40-3.
916. Massaro, A., Hammad, T., Jazzo, B., & Aly, H. (2009). Massage with kinesthetic stimulation improves weight gain in preterm infants. *J Perinatal, 29*(5), 352-7.

917. Melzack, R., & Wall, P. (1965). Pain mechanisms: A new theory. *Science*, 150(699), 971-9.
918. Moore, I. (2004). Advancing biobehavioral research in childhood cancer. *JPON*, 21(3), 12&131.
919. Moyer, C., Seefeldt, L., Mann, E., & Jackley, L. (2011). Does massage therapy reduce cortisol? A comprehensive quantitative review. *J Bodyw Mov Ther*, 15, 3-14.
920. National Cancer Institute. (2008). *Childhood cancers*. Retrieved from www.cancer.gov/cancertopics/factsheet/Sites-Types/childhood.
921. Norris, S., Pare, J., & Starky, S. (2006). *Childhood autism in Canada: Some issues relating to behavioural intervention*. Ottawa, ON: Library of Parliament.
922. Owens, J., & Jones, C. (2011). Parental knowledge of healthy sleep in young children: Results of a primary care clinic survey. *J Dev Behav Pediatr*, 32(6), 447-53.
923. Post-White, J., Fitzgerald, M., Savik, K., Hooke, M., Hannahan, A., & Sencer, S. (2009). Massage therapy for children with cancer. *J Pediatr Oncol Nurs*, 26(1), 16-28.
924. Procianny, R., Mendes, E., & Silveira, R. (2010). Massage therapy improves neuro-development outcome at two years corrected age for very low birth weight infants. *Early Hum Dev*, 86(1), 7-11.
925. Sagar, S., Dryden, T., & Wong, R. (2007). Massage therapy for cancer patients: A reciprocal relationship between body and mind. *Curr Oncol*, 14(2), 45-56.
926. Sefton, J., Yarar, C., Berry, J., & Pascoe, D. (2010). Therapeutic massage of the neck and shoulders produces changes in peripheral blood flow when assessed with dynamic infrared thermography. *J Altern Complement Med*, 16(7), 723-32.
927. Snyder, H., & Sickmund, M. (2006). *Juvenile offenders and victims: 2006 National Report*, ed. Office of Juvenile Justice and Delinquency Prevention. Washington, DC: Office of Juvenile Justice and Delinquency Prevention.
928. Suomi, S., Collins, M., & Harlow, H. (1973). Effects of permanent separation from mother on infant monkeys. *Dev Psychol*, 9(3), 376-84.
929. Taylor-Butts, A., & Bressan, A. (2008). *Youth crime in Canada, 2006*, ed. Statistics Canada. Ottawa, ON: Minister of Industry.
930. Temple, R. (2004). *Antidepressant drug use in pediatric populations*. Statement before the Senate Subcommittee on Oversight and Investigations, Committee on Energy and Commerce. Retrieved from www.fda.gov/NewsEvents/Testimony/ucrn113265.htm.
931. Underdown, A., Barlow, J., Chung, V., & Stewart-Brown, S. (2006). Massage intervention for promoting mental and physical health in infants aged under six months. *Cochrane Lib* 4 CD005038.
932. Vannorsdall, T., Dahlquist, L., Pendley, J., & Power, T. (2004). The relation between nonessential touch and children's distress during lumbar punctures. *Child Health Care*, 33(4), 299-315.
933. Vickers, A., Ohlsson, A., Lacy, J., & Horsley, A. (2004). Massage for promoting growth and development of preterm and/or low birth-weight infants. *Cochrane Db Syst Rev*, (2), CD000390.
934. von Knarring, A., Soderberg, A., Austin, L., & Uvnas-Moberg, K. (2008). Massage decreases aggression in preschool children: A long-term study. *Acta Paediatr*, 97(9), 1265-9.
935. Watson, J. & Watson, R. (1928). *Psychological care of infant and child*. New York: W. W. Norton.
936. Weber, W., & Newmark, S. (2007). Complementary and alternative medical therapies for attention-deficit/hyperactivity disorder and autism. *Pediatr Clin North Am*, 54(6), 983-1006; xii.
937. Young, K. (2005). Pediatric Procedural Pain. *Ann Emerg Med*, 45, 160-71.
- Older Adults**
938. American Association of Retired Persons (AARP), & National Centre for Complementary and Alternative Medicine (UCCAM). (2007). Complementary and alternative medicine: What people 50 and older are using and discussing with their physicians. Retrieved from <http://assets.aarp.org/rgcenter/health/cam~2007.pdf>.
939. Barnes, P., Bloom, B., & Nahin, R. (2008). Complementary and alternative medicine use among adults and children: United States, 2007. *Nat Health Stat Rep*, 12, 1-23.
940. Berger, L., Klein, C., & Commandeur, M. (2008). Evaluation of the immediate and midterm effects of mobilization in hot spa water on static and dynamic balance in elderly subjects. *Ann Readapt Med Phys*, 51, 90-5.
941. Beider, S. (2005). An ethical argument for integrated palliative care. Retrieved from <http://ecam.oxfordjournals.org/cgi/reprint/neh089vl.pdf>.
942. Billhult, A., Bergborn, I., & Stener-Victorin, E. (2007). Massage relieves nausea in women with breast cancer who are undergoing chemotherapy. *J Altern Complement Med* 13, (1), 55-7.
943. Centers for Disease Control and Prevention (CDC). (2006). Prevalence of doctor diagnosed arthritis and arthritis-attributable activity limitation: United States, 2005. *MWR Weekly*, 55(40), 1089-92.
944. Cerrone, R., Giani, L., Galbiati, B., Messina, G., Castraghi, M., Proserpio, E., & Gardani, G. (2008). Efficacy of ht 7 point acupressure stimulation in the treatment of insomnia in cancer patients and in patients suffering from disorders other than cancer. *Minerva Medical*, 99, 535-7.
945. Cherkin, D., Eisenberg, D., Sherman, K., Barlow, W., Kaptchuk, T., Street, J., & Deyo, R. (2001). Randomized trial comparing traditional Chinese medical acupuncture, therapeutic massage, and selfcare education for chronic low back pain. *Arch intern Med*, 161, 1081-8.
946. Cronfalk, B., Strang, P., Texnstedt, B., & Friedrichsen, M. (2009). The existential experiences of receiving soft tissue massage in palliative home care—An intervention. *Support Care Cancer*, 17, 1203-11.

947. Currin, J., & Meister, E. (2008). A hospital-based intervention using massage to reduce distress among oncology patients. *Cancer Nurs*, 31(3), 214-21.
948. Davis, M., & Srivastava, M. (2003). Demographics, assessment and management of pain in elderly. *Drug Aging*, 20(1), 23-57.
949. Dryden, T., Baskwill, A., & Preyde, M. (2004). Massage therapy for the orthopaedic patient: A review. *Orthop Nurs*, 23(5), 327-32.
950. Federal Interagency Forum on Aging-Related Statistics. (2008). *Older Americans 2008: Key indicators of well-being*. Federal Interagency Forum on Aging-Related Statistics. Washington, DC: U. S. Government Printing Office.
951. Flaherty, J. (2008). Insomnia among hospitalized older persons. *Clin Geriatr Med*, 24, 51-67. doi: 10.1016/j.cger.2007.08.012
952. Forestier, R., Desfour, H., Tessier, J., Francon, A., Foote, A., Genty, C., & Bosson, J. (2009). Spa therapy in the treatment of knee osteoarthritis, a large randomized multicentre trial. *Ann Rheum Dis*, 69(4), 660-5.
953. Gerdner, L., Hart, L., & Zimmerman, M. (2008). Craniosacral still point technique: Exploring its effect in individuals with dementia. *J Gerontol Nurs*, 34(3), 36-45.
954. Greenberg, S. (2008). A profile of older Americans: 2008 administration on aging. *U.S. Department of Health and Human Services*. Retrieved from www.aoa.gov/aoaroot/agingstatistics/Profile/index.aspx.
955. Gooneratne, N. (2008). Complementary and alternative medicine for sleep disturbances in older adults. *Clin Geriatr Med*, 24, 121-38. doi: 10.1016/j.cger.2007.08.002
956. Hanley, J., Stirling, O., & Brown, C. (2003). Randomized controlled trial of therapeutic massage in the management of stress. *Br J Gen Pract*, 53(486), 20-5.
957. Hasson, D., Arnetz, B., Jelveus, L., & Edelstam, B. (2004). A randomized clinical trial of the treatment effects of massage compared to relaxation tape recordings on diffuse long-term pain. *Psychother Psychosom*, 73, 17-24.
958. Hernandez-Reif, M., Field, T., Krasnegor, J., & Theakston, H. (2001). Lower back pain is reduced and range of motion increased after massage therapy. *Int J Neurosci*, 106, 131-45.
959. Hodgson, N., & Anderson, S. (2008). The clinical efficacy of reflexology in nursing home residents with dementia. *J Altern Complem Med*, 14(3), 269-75.
960. Holliday-Welsh, D., Gessert, C., & Renier, C. (2009). Massage in the management of agitation in nursing home residents with cognitive impairment. *Geriatr Nurs*, 30, 108-17.
961. Kolcaba, K., Dowd, T., Steiner, R., & Mitzel, A. (2004). Efficacy of hand massage for enhancing the comfort of hospice patients. *Journal of Hospice and Palliative Nursing*, 6(2), 91-102.
962. Kolcaba, K., Schirm, V., & Steiner, R. (2006). Effects of hand massage on comfort of nursing home residents. *Geriatr Nurs*, 27, 85-91.
963. Kozak, L., Kayes, L., McCarty, R., Walkinshaw, C., Congdon, S., Kleinberger, J., Hartman, V., & Standish, L. (2009). Use of complementary and alternative medicine (CAM) by Washington State hospices. *Am J Hosp Palliat Me*, 5(6), 463-8.
964. Kutnex, J., Smith, M., Corbin, L., Hemphill, L., Benton, K., Mdlis, B., & Fairclough, D. (2008). Massage therapy verses simple touch to improve pain and mood in patients with advanced cancer: A randomized trial. *Ann Intern Med*, 149, 369-79.
965. Lamas, K., Lindholm, L., Stenlund, H., Engstrom, B., & Jacobsson, C. (2009). Effects of abdominal massage in management of constipation: A randomized controlled trial. *Int J Nurs Stud*, 46, 759-67.
966. Lamas, K., Lindholm, L., Engstrom, B., & Jacobsson, C. (2010). Abdominal massage for people with constipation: A cost utility analysis. *J Adv Nurs*, 66(8), 1719-29.
967. MacDonald, G. (2005). *Massage for the hospital patient and medically frail client*. Hagerstown, MD: Lippincott, Williams and Wilkins.
968. Meeks, T., Wetherell, J., Irwin, M., Redwine, L., & Jeste, D. (2007). Complementary and alternative treatments for late-life depression, anxiety, and sleep disturbance: A review of randomized controlled trials. *J Clin Psychiatry*, 68(10), 1461-71.
969. Morien, A., Garrison, D., & Smith, N. (2008). Range of motion improves after massage in children with burns: A pilot study. *J Bodyw Mov Thers*, 12, 67-71.
970. Moyer, C., Rounds, J., & Hannum, J. (2004). A meta-analysis of massage therapy research. *Psychol Bull*, 130(1), 3-18.
971. Munk, N., & Faika, Z. (2011). Relationship between massage therapy usage and health outcomes in older adults. *Journal of Bodywork & Movement Therapies*, 15, 177-85.
972. National Hospice and Palliative Care Organization (NHPCO). (2009). NHPCO facts and figures: Hospice care in America. Retrieved from www.nhpco.org/~files/public/Statistics~Uesearch/NHPCO-facts-and-figures.pdf.
973. Oneschuk, D., Bdnaveas, L., Verhoef, M., Boon, H., Demmer, C., & Chiu, L. (2007). The status of complementary therapy services in Canadian palliative care settings. *Support Care Cancer*, 15, 939-47.
974. Perlman, A., Sabina, A., Williams, A., Njike, V., & Katz, D. (2006). Massage therapy for osteoarthritis of the knee: A randomized controlled trial. *Arch Intern Med*, 166(22), 2533-8.
975. Preyde, M. (2000). Effectiveness of massage therapy for sub-acute low back pain: A randomized controlled trial. *CMAJ*, 162, 1815-20.
976. Reid, M., Papaleontiou, M., Ong, A., Breckman, R., Wethington, E., & Pillemer, K. (2008). Self-management strategies to reduce pain and improve function among older adults in community settings: A review of the evidence. *Pain Med*, 9(4) 409-24.
977. Rose, M. (2010). *Comfort touch: Massage for the elderly and the ill*. Hagerstown, MD: Lippincott, Williams, and Wilkins.
978. Russell, N., Beinhorn, C., & Frenkel, M. (2008). Role of massage therapy in cancer care. *J Altern Complem Med*, 4(2), 209-14.
979. Soden, K., Vincent, K., & Craske, S. (2004). A randomized controlled trial of aromatherapy massage in a hospice setting. *Palliative Med*, 18(2), 87-92.
980. Thompson, D. (2006). *Hands heal: Communication, documentation, and insurance billing for manual Therapists*. 3rd ed. Hagerstown, MD: Lippincott, Williams, and Wilkins.

981. Tsao, J. (2007). Effectiveness of massage therapy for chronic, non-malignant pain: A review. *Evid-Based Complement Altern Med*, 4(2), 165-79.
982. Vaht, M., Birkenfeldt, R., & Ubner, M. (2008). An evaluation of the effect of differing lengths of spa therapy upon patients with osteoarthritis. *Complement Ther Clin Prod*, 14, 60-4.
983. Vaillant, J., Rouland, A., Martigne, P., Braujou, R., Nissen, M., Caillat-Miousse, J., Vuillerme, N., Nougier, V., & Juvin, R. (2009). Massage and mobilization of the feet and ankles in elderly adults: Effect on clinical balance performance. *Manual Ther*, 14(6), 661-4.
984. Walach, H., Guthlin, C., & Konig, M. (2005). Efficacy of massage therapy in chronic pain: A pragmatic randomized trial. *J Altern Complem Med*, 9, 837-46.
985. Williams, A., Selwyn, P., Liberti, L., Molde, S., Njike, V., McCorkle, R., & Katz, D. (2005). A randomized controlled trial of meditation and massage effects on quality of life in people with late stage disease: A pilot study. *J Palliat Med*, 8(5), 939-52.
986. Woods, D., & Dimond, M. (2002). The effect of therapeutic touch on agitated behavior and cortisol on persons with Alzheimer's disease. *Biol Res Nur*, 4(2), 104-14.
987. Woods, D., Craven, R., & Whitney, J. (2005). The effects of therapeutic touch on behavioral symptoms of persons with dementia. *Altern Ther*, 11(1), 66-74.

Emerging Evidence & Conclusion

988. Cerrone, R., Giani, L., Galbiati, B., Messina, G., Castraghi, M., Proserpio, E., & Gardani, G. (2008). Efficacy of ht 7 point acupressure stimulation in the treatment of insomnia in cancer patients and in patients suffering from disorders other than cancer. *Minerva Medical*, 99, 535-7.
989. Eisenberg DM, Kessler RC, Foster C, Norlock FE, Calkins, DR, & Delbanco TL. (1993). Unconventional medicine in the united states: Prevalence, costs, and patterns of use. *N Engl J Med*. 328(4):246-52.
990. Goldblatt, E., Snider, P., Quinn, S., & Weeks, J. (2009). *Clinicians' and educators' desk reference on the licensed complementary and alternative healthcare professions*. ACCAHC.
991. Hsu C, Bluespruce J, Sherman K, & Cherkin D (2010). Unanticipated benefits of CAM therapies for back pain: An exploration of patient experiences. *J Altern Complement Med*, 16(2):157-63.
992. Lamas, K., Lindholm, L., Stenlund, H., Engstrorn, B., & Jacobsson, C. (2009). Effects of abdominal massage in management of constipation: A randomized controlled trial. *Int J Nurs Stud*, 46, 759-67.
993. Lamas, K., Lindholm, L., Engstrom, B., & Jacobsson, C. (2010). Abdominal massage for people with constipation: A cost utility analysis. *J Adv Nun*, 66(8), 1719-29.
994. Rose KE, Taylor HM, Twycross R.(1991). Long-term compliance with treatment in obstructive arm lymphoedema in cancer. *Palliat Med*, 5, 52-5.
995. Woods, D., Craven, R., & Whitney, J. (2005). The effects of therapeutic touch on behavioral symptoms of persons with dementia. *Altern Ther*, 11(1), 66-74.