

OREGON PHYSICIANS FOR SOCIAL RESPONSIBILITY

<http://www.psr.org/chapters/oregon/>

AIRBORNE PARTICULATE MATTER AND PUBLIC HEALTH

BACKGROUND

Outdoor air quality is an increasing concern globally with expanding industrial and transportation emissions. The World Health Organization (WHO) has declared that Particulate Matter (PM) in ambient outdoor air affects more people than any other pollutant. Chronic exposure to particles contributes to the risk of developing or dying from serious disease. (WHO 1) The effects of PM on health occur at levels of exposure currently being experienced by most urban and rural populations in both developed and developing countries. Burning fuel results in two phases of emissions. In addition to the “particulate phase,” there is a “gas phase” containing air pollutants such as benzene, formaldehyde, polycyclic aromatic hydrocarbons (PAHs), and other chemicals, which also contribute to disease. (EPA 1, EPA 2) The combustion of fossil fuel for electricity and transportation, especially coal and diesel, are major contributors to outdoor air pollution.

WHAT IS PARTICULATE MATTER?

Particulate Matter (PM), is composed of very small, solid particles, formed from the incomplete burning of fossil fuels, such as coal, diesel, gasoline, and biomass. PM_{2.5} measure 2.5 microns or less; PM₁₀ are particles 10 microns or smaller. PM consists of a complex mixture of soot, black carbon, absorbed water, aerosolized sulfuric acid droplets, other acids, nitrogen, sulfur, metals, and other toxic substances, especially PAHs (Perera), which are absorbed by the sponge-like particles and carried by them deeply into the smallest compartments of the lung (alveoli) where they gain direct access to the bloodstream and may then contribute to various diseases in organs distant from the lungs.

The World Health Organization (WHO) reports that there is not a threshold below which no damage to human health is observed. Their guidelines for maximizing health within the constraints of a modern world are that PM_{2.5} should not exceed an average of 25 micrograms per cubic meter of air (25mcg/m³) in a 24-hour period, and not exceed an average annual exposure of 10mcg/m³. (WHO 1) To put this into perspective, the PM_{2.5} for Portland, Oregon in January, 2014 ranged from 30-76, while, during same month, the PM_{2.5} in Beijing, China was over 600. In Harbin, China in October of 2013 (the start of the heating season), the PM_{2.5} was over 1000mcg/m³ (Guardian). The major contributors to this astronomically high PM_{2.5} in China are burning coal and vehicle exhaust.

HOW DOES PARTICULATE MATTER AFFECT PEOPLE?

The fine particles less than 2.5 microns (PM_{2.5}) are particularly important in triggering disease because they penetrate deeply into the alveoli of the lungs. Some inhaled particles are taken up by macrophages, resulting in lung inflammation. The final common pathway of the pathologic effects of exposure to particulate matter, as well as gas phase pollutants such as benzene and polycyclic aromatic hydrocarbons, appears to be inflammation. (Avogbe, Bellavia, Dominici, Mustafic, Peters 2011, Pieters). The effects of inflammation on various body organ systems are complex, but increased levels of particulate matter are associated with a number of ill health effects including: increased cancer rates, especially lung and breast (Beeson, Crouse, Demetriou, Dockery, Pope, Wei), congenital lung, heart and immune system anomalies in children (Gauderman, Picciotto Vrijheid,) increased rates of asthma, worsening of preexisting asthma and chronic obstructive pulmonary disease (COPD) (Carlsten, Gowers, Delamater, HEI Panel, Trasande, WHO 2), higher rates of heart attacks and strokes (Chen, Dominici, Lie, Mustafic, Qian, Wellenius, Shaw), and higher rates in children (exposed prenatally) of neurodevelopmental disorders such as autism spectrum disorder (ASD), attention deficit hyperactivity disorder (ADHD), lowered IQ, and adverse behaviors. (Becera, Chiu, Newman, Perera 2013, Perera 2009, Roberts, Volk 2013, Volk 2011) Not surprisingly, the most vulnerable populations are the elderly, pregnant women, children, and people that already have pulmonary diseases like COPD or asthma.

WHAT ARE CURRENT ISSUES IN THE PACIFIC NORTHWEST?

A recent study from the University of Washington, evaluating air quality as a result of increased coal transportation by rail in the Pacific Northwest, found that living within 25 meters of two rail lines in the Blue Ridge Seattle neighborhood was associated with significantly increased exposure to diesel particulate matter (DPM) and airborne coal dust. A proposed 50% increase in diesel train traffic (if coal terminals are built in coastal

harbors) is projected to similarly increase small particulate matter exposure. When added to regional background emissions, this increase would bring total concentrations higher than the new US National Ambient Air Quality Standards (NAAQS). (Jaffe)

In addition, a sharp increase in mile-long trains already transporting oil through the region has considerably increased airborne diesel emissions.

KEY AMERICAN MEDICAL SOCIETIES AND THE WORLD HEALTH ORGANIZATION HAVE ISSUED POSITIONS ON PARTICULATE MATTER AND HEALTH:

The *American Heart Association* (AHA) in 2010 updated and summarized its 2004 Scientific statement position: “The overall evidence is consistent with a causal relationship between PM2.5 exposure and cardiovascular morbidity and mortality. This body of evidence has grown and has been strengthened substantially since publication of the first AHA scientific statement and, ... because the evidence reviewed supports that there is no safe threshold, it appears that public health benefits would accrue from lowering PM2.5 concentrations even below present-day (EPA standards), if feasible, to optimally protect the most susceptible populations.” (Brook)

The American College of Obstetricians and Gynecologists (ACOG) together with the American Society of Reproductive Medicine (ASRM) in October 2013 issued a statement, “The evidence that links exposure to toxic environmental agents and adverse reproductive and developmental health outcomes is sufficiently robust, ... individuals alone can do little about exposure to toxic environmental agents, such as from air and water pollution, ... calling for timely action to identify and reduce exposure.” (ACOG)

The *American Academy of Pediatrics* (AAP) issued a policy statement linking ambient air pollution to adverse health outcomes in children and recommended the National Ambient Air Quality Standards (NAAQS) be promptly reviewed and revised to protect children. (AAP, 2004, reaffirmed 2009)

In October 2013, *WHO's International Agency for Research on Cancer (IARC)*, classified both outdoor air pollution, as a whole, and particulate matter, on its own, as carcinogenic. Therefore, it is vital to implement efficient policies to reduce exposure to pollution worldwide. (World Health Organization (WHO 2) and American Cancer Society).

SPECIFIC DISEASES ASSOCIATED WITH EXPOSURE TO HIGHER LEVELS OF PARTICULATE MATTER IN AIR POLLUTION

Pulmonary - Studies have demonstrated the effects of particulate matter on the lungs:

- decreased lung function (WHO 3)
- inhibited lung development in children (Gauderman)
- causes and makes asthma worse, resulting in increased hospitalization (Delameter, Gowers, HEI Panel, Pandya)

Cardiovascular - Studies have linked increased particulate matter with increased cardiac disease:

- increased cardiovascular disease mortality and morbidity in both short term and long term exposures to PM 2.5 (Brook)
- increased hospital admissions for serious cardiac arrhythmias (Peters 2000)
- increased probability of admission for acute myocardial infarction (Mustafic, Peters 2001)
- increased ischemic heart disease, arrhythmias, congestive heart failure (Dominici) and bio markers (HRV) associated with increased cardiac morbidity and mortality (Pieters)
- increased hospital admissions and death from heart failure (Shaw)
- increased risk of congenital cardiac anomalies in children (Vrijheid)

Cerebrovascular – Studies have shown links between particulate matter and strokes:

- increased hospital admissions for strokes (Dominici, Lue, Wellenius 2005)
- significant increase in stroke mortality associated with increase in PM (Chen, Qian)
- increased risk of stroke associated with increased exposure to small PM, black carbon, and nitrous dioxide (Wellenius 2012)
- increased risk of stroke and death from stroke for post menopausal women (Miller)

Cancer – Studies relating cancer risk and particulate matter

- Exposure to ozone and PM correlated with development of and mortality from lung cancer (Beeson, Dockery, Pope)
- increased biological markers associated with risk of lung cancer (Demetriou)
- increased oxidative DNA damage predictive of cancer risk (Avogbe)
- increased rates of breast cancer (Crouse, Wei)

Neurodevelopmental— Studies associating in-utero exposure to particulate matter and

- Increased incidence of autism spectrum disorder (ASD)—(Becerra, Roberts, Volk 2013, Volk 2011)
- Increased incidence of behaviors associated with attention deficit hyperactivity disorder (ADHD) (Chiu, Newman)
- Lowered IQ (Perera 2009)
- Increased behavioral symptoms of anxiety, depression, social problems, rule breaking, and aggression (Perera 2013)
- Neurobehavioral development in children benefited from the shutdown of a coal-burning plant (Perera 2008)

General

- increased mortality from cardiac, respiratory and kidney disease in all members of communities with coal exposure (15,16,17,18 Hendryx 2007, Hendryx 2010, Hendryx 2008, Hendryx 2009)
- long term exposure linked to decreased life expectancy from cardiopulmonary mortality (Krishnan, WHO 4)
- prenatal exposures linked to altered immune system development (Hertz-Picciotto)

WHAT YOU CAN DO TO PROTECT YOUR FAMILY AND COMMUNITY

The World Health Organization has stated, “Most sources of outdoor air pollution are well beyond the control of individuals and demand action by cities, as well as national and international policymakers in sectors like transport, energy, waste management, buildings and agriculture.” With that in mind, here are some tips to protect your health:

1. Use a **portable air purifier** in your home (Look to consumer organizations for specific product information).
2. Do the best you can to **protect your and your family’s health from the consequences of inflammation** by observing good health habits: Get plenty of sleep and daily exercise, eat six servings of fruits and vegetables daily, consult your health advisors.
3. Push in the **air recycle button on your car dash** when traveling on motorways around other vehicles.
4. **Work for change with environmental and neighborhood groups** (examples: The NW District Association of Portland recently negotiated an agreement with local developers to meet EPA’s highest standards for diesel exhaust emissions, by using exhaust treatment devices during a local construction project. Neighbors for Clean Air negotiated with an industry in the Northwest Portland neighborhood to reduce its toxic air emissions, by installing giant vacuum cleaner bags to capture particulates, and air monitoring equipment in the neighborhood.)
5. **Work for policy change with local and regional governments, and legislatures.**
Ideas for starters (inspired by the Oregon Environmental Counsel): Request “No diesel engine idling” policies, retrofitting of older diesel engines with pollution control devices, advocate for the use of ultra-low sulfur diesel fuel for Oregon, continue negotiating improvement in the emissions standards of heavy-duty, off-road diesel engines, request that all vehicles are using the best available technology for emission control, and that grant funds are available to assure this, especially for small businesses.

RESOURCES THAT INFORMED THIS FACTSHEET

ACOG—The American College of Obstetricians and Gynecologists, Committee Opinion, No. 575, October 2013 - http://www.acog.org/About_ACOG/News_Room/News_Releases/2013/Environmental_Chemicals_Harm_Reproductive_Health

AAP—American Academy of Pediatrics. Committee on Environmental Health. Ambient Air Pollution: Health Hazards to Children, , *Pediatrics*, 2004: 114, 1699-1707, Reaffirmed April 2009 (<http://pediatrics.aappublications.org/content/125/2/e444.extract>)

Avogbe P, Ayi-Fanou L, Autrup H, et al. Ultrafine particulate matter and high-level benzene urban air pollution in relation to oxidative DNA damage. *Carcinogenesis* 2004; 26(3):613-620

Becerra TA, Wilhelm M, Olsen J, et al. Ambient air pollution and autism in Los Angeles County, California. *Environ Health Perspect* 2013; 121(3):380-386

Beeson WL, Abbey DE, Knutsen SF. Long-term concentrations of ambient air pollutants and incident lung cancer in California adults: results from the Adventist Health Study on Smog. *Environ Health Perspect* 1998; 106(12):813-23

Bellavia A, Urch B, Speck M, et al. DNA Hypomethylation, ambient particulate matter, and increased blood pressure: findings from controlled human exposure experiments. *J Am Heart Assoc.* 2013; 2:e000212 doi: 10.1161/JAHA.113.000212

Brook RD, Rajagopalan S, Pope CA, III, et al. Particulate matter air pollution and cardiovascular disease: An update to the scientific statement from the American Heart Association. *Circulation* 2010;121(21):2331-2378

Carlsten C, Dybuncio A, Becker A, et al. Traffic-related air pollution and incident asthma in a high-risk birth cohort. *Occup Environ Med* 2011; 68:291-295

Chen R, Zhang Y, Yang C, et al. Acute effect of ambient air pollution and health effects study. *Stroke* 2013; 44:954-960

Chiu Y-HM, Bellinger D, Coull BA, et al. Associations between traffic-related black carbon exposure and attention in a prospective birth cohort of urban children. *Environ Health Perspect* 2013; 121(7):859-864

Crouse D, Goldberg M, Ross N, et al. Postmenopausal breast cancer is associated with exposure to traffic –related air pollution in Montreal, Canada: a case-controlled study. *Environmental Health Perspectives* 2010; 118(11): 1578-1583

Delamater P, Finley A, Banerjee S. An analysis of asthma hospitalizations, air pollution, and weather conditions in Los Angeles County, California. *Science of the Total Environment* 2012; 425:110-118

Demetriou C, Raaschou-Nielsen O, Loft S, et al. Biomarkers of ambient air pollution and lung cancer: a systematic review. *Occup Environ Med* 2012; 69:619-627

Dockery DW, Pope CA, III, Xu X, et al. An association between air pollution and mortality in six US cities. *N Engl J Med* 1993; 329(24):1753-1759

Dominici F, Peng RD, Bell ML et al. Fine Particulate air pollution and hospital admission for cardiovascular and respiratory diseases. *JAMA* 2006; 295(10):1127-1134

EPA 1—<http://www.epa.gov/region1/eco/airtox/diesel.html>

EPA 2— http://www.epa.gov/teach/chem_summ/BENZ_summary.pdf

Gauderman WJ, Avol E, Gilliland F, et al. The effect of air pollution on lung development from 10 to 18 years of age. *N Engl J Med* 2004;351(11):1057-1067

Gowers A, Cullinan P, Ayres J, et al. Does outdoor air pollution induce new cases of asthma? Biological plausibility and evidence; a review. *Respirology* 2012; 17:887-898

Guardian: <http://www.theguardian.com/environment/chinas-choice/2013/oct/24/china-airpocalypse-harbin-air-pollution-cancer>

HEI Panel on Health effects of traffic related air pollution, Traffic related Air Pollution A Critical review of the literature on emissions, exposure, and health effects. Boston: Health Effects Institute; 2010.

Hendryx M, Ahern MM, Nurkiewicz TR. Hospitalization patterns associated with Appalachian coal mining. *Journal of Toxicology and Environmental Health, Part A.* 2007;70(24):2064-2070

Hendryx M, Fedorko E, Anesetti-Rothermel A. A geographical information system-based analysis of cancer mortality and population exposure to coal mining activities in West Virginia, United States of America. *Geospatial Health* 2010;4(2):243-256

Hendryx M, Ahern MM. Relations between health indicators and residential proximity to coal mining in West Virginia. *American Journal of Public Health* 2008;98(4):669-671

Hendryx M. Mortality from heart, respiratory, and kidney disease in coal mining areas of Appalachia. *International Archives of Occupational and Environmental Health*. 2009;82(2):243-249

Hertz-Picciotto I, Park H-Y, Dostal M, et al. Prenatal exposures to persistent and non-persistent organic compounds and effects on immune system development. *Basic and Clinical Pharmacology and Toxicology* 2007;102:146-154

Jaffe, Daniel A., et al. Diesel particulate matter emission factors and air quality implications from in-service rail in Washington State, USA, *Atmospheric Pollution Research*. Doi: 10.5094/Apr. 2014.040

Krishnan RM, Adar SD, Szpiro AA, et al. Vascular responses to long- and short-term exposure to fine particulate matter, MESA Air (Multi-ethnic study of atherosclerosis and air pollution. *J Am College of Cardiology* 2012; 60(21):2158-2166

Lue S-H, Wellenius GA, Wilker EH, et al. Residential proximity to major roadways and renal function. *J Epidemiol Community Health* 2013; 67:629-634

Miller KA, Siscovick DS, Sheppard L, et al. Long term exposure to air pollution and incidence of cardiovascular events in women. *N Engl J Med* 2007;356(5):447-5

Mustafic H, Jabre P, Caussin C, et al. Main air pollutants and myocardial infarction, a systematic review and meta-analysis. *JAMA* 2012; 307(7):713-721

Newman NC, Ryan P LeMasters G, et al. Traffic-related air pollution exposure in the first year of life and behavioral scores at 7 years of age. *Environ Health Perspect* 2013; 121(6):731-736

Pandya RJ, Solomon G, Kinner A, Balmes JR. Diesel exhaust and asthma: hypotheses and molecular mechanisms of action. *Environmental Health Perspectives*. 2002;110(suppl 1):103-112

Perera FP, Li Z, Whyatt R, et al. Prenatal airborne polycyclic aromatic hydrocarbon exposure and child IQ at age 5 years. *Pediatrics* 2009; 124(2):e195-202

Perera FP, Wang S, Rauh V, et al. Prenatal exposure to air pollution, maternal psychological distress, and child behavior. *Pediatrics* 2013; 132(5):e1284-1294

Perera FP, Li T, Zhou Z, et al. Benefits of reducing prenatal exposure to coal-burning pollutants to children's neurodevelopment in China. *Environmental Health Perspectives* 2008; 116(10):1396-1400

Peters A. Ambient particulate matter and the risk for cardiovascular disease. *Progress in Cardiovascular Diseases*. 2011; 53: 327-333

Peters A, Liu E, Verrier RL et al, Air pollution and incidence of cardiac arrhythmia. *Epidemiology* 2000; 11(1):11-17

Peters A., Dockery DW, Muller JE, Mittleman MA. Increased particulate air pollution and the triggering of myocardial infarction. *Circulation* 2001; 103(23):2810-2815

Pieters N, Plusuin M, Cox B, et al. An epidemiological appraisal of the association between heart rate variability and particulate air pollution: a meta-analysis. *Heart* 2012; 98:1127-1135

Pope CA, III, Burnett RT, Thun MJ et al. Lung cancer, cardiopulmonary mortality, and long term exposure to fine particulate air pollution. *JAMA* 2002; 287(9):1132-1141.

Qian Y, Zhu M, Cai B, et al. Epidemiological evidence on association between ambient air pollution and stroke mortality. *J Epidemiol Community Health* 2013; 67:635-640

Roberts AL, Lyall K, Hart, JE, et al. Perinatal air pollutant exposures and Autism Spectrum Disorder in the children of nurses' health study II participants. *Environ Health Perspect* 2013; 121(8): 978-984

Shaw ASV, Langrish JP, Nair H, et al. Global associations of air pollution and heart failure: a systematic review and meta-analysis. *The Lancet* 2013; 832:1039-1048

Trasande L, Thurston GD. The role of air pollution in asthma and other pediatric morbidities. *J Allergy Clin Immunology* 2005; 115(4):689-699

Vrijheid M, Martinez D, Manzanares, S, et al. Ambient air pollution and risk of congenital anomalies: a systematic review and meta-analysis. *Environmental Health Perspectives* 2011;119(5) 598-606

Volk HE, Lurmann F, Penfold B, et al. *JAMA Psychiatry* 2013; 70(1):71-77

Volk HE, Hertz-Picciotto I, Delwiche L, et al. Residential proximity to freeway and autism in the CHARGE study. *Environ Health Perspect* 2011; 119(6):873-877

Wei Y, Davis J, Bina WF. Ambient air pollution is associated with the increased incidence of breast cancer in US. *Int J Environ Health Res* 2012: 22(1):12-21

Wellenius GA, Schwartz J, Mittleman MA. Air Pollution and hospital admissions for ischemic and hemorrhagic stroke among Medicare beneficiaries. *Stroke* 2005; 36(12):2549-2553

Wellenius GA, Burger, MR, Coull BA, et al. Ambient air pollution and the risk of acute ischemic stroke. *Arch Int Med* 2012; 172(3): 229-234

WHO 1—World Health Organization <http://www.who.int/mediacentre/factsheets/fs313/en/#>

WHO 2— World Health Organization/ American Cancer Society
<http://www.cancer.org/cancer/news/world-health-organization-outdoor-air-pollution-causes-cancer>

WHO 3— World Health Organization. Health aspects of air pollution with particulate matter, ozone, and nitrogen dioxide. Bonn: World Health Organization; 2003

WHO 4—World Health Organization. Outdoor Air Pollution. Global Health Observatory. Available at <http://www.who.int/gho/phe/outdoor-air-pollution/en/index.html>