

DIAGNOSTIC IMAGING PRACTICE GUIDELINES FOR MUSCULOSKELETAL COMPLAINTS IN ADULTS—AN EVIDENCE-BASED APPROACH—PART 3: SPINAL DISORDERS

André E. Bussi eres, DC,^a John A.M. Taylor, DC,^b and Cynthia Peterson, DC, RN, MMedEd^c

ABSTRACT

Purpose: To develop evidence-based diagnostic imaging practice guidelines to assist chiropractors and other primary care providers in decision making for the appropriate use of diagnostic imaging for spinal disorders.

Methods: A comprehensive search of the English and French language literature was conducted using a combination of subject headings and keywords. The quality of the citations was assessed using the Quality of diagnostic accuracy studies (QUADAS), the Appraisal of Guidelines Research and Evaluation (AGREE), and the Stroke Prevention and Educational Awareness Diffusion (SPREAD) evaluation tools. The Referral Guidelines for Imaging (radiation protection 118) coordinated by the European Commission served as the initial template. The first draft was sent for an external review. A Delphi panel composed of international experts on the topic of musculoskeletal disorders in chiropractic radiology, clinical sciences, and research were invited to review and propose recommendations on the indications for diagnostic imaging. The guidelines were pilot tested and peer reviewed by practicing chiropractors, and by chiropractic and medical specialists. Recommendations were graded according to the strength of the evidence.

Results: Recommendations for diagnostic imaging guidelines of adult spine disorders are provided, supported by more than 385 primary and secondary citations. The overall quality of available literature is low, however. On average, 45 Delphi panelists completed 1 of 2 rounds, reaching more than 85% agreement on all 55 recommendations. Peer review by specialists reflected high levels of agreement, perceived ease of use of guidelines, and implementation feasibility. Dissemination and implementation strategies are discussed.

Conclusions: The guidelines are intended to be used in conjunction with sound clinical judgment and experience and should be updated regularly. Future research is needed to validate their content. (*J Manipulative Physiol Ther* 2008;31:33-88)

Key Indexing Terms: *Practice Guideline; Guideline; Diagnostic Imaging; Radiology; Diagnostic X-Ray; Radiography; Adult; Musculoskeletal System; Pain; Cervical Spine; Thoracic Spine; Lumbar Spine; Trauma*

REPORTING OF TOPICS INCLUDED IN THE DEVELOPMENT OF THE DIAGNOSTIC IMAGING PRACTICE GUIDELINES¹

An initial literature review considered 10 clinical questions pertaining to imaging of musculoskeletal conditions to evaluate the pertinence of developing diagnostic

imaging guidelines. This initial review led to a research project divided into 9 phases: (1) literature search; (2) independent literature assessment; (3) guideline development specific recommendations; (4) first external review; (5) consensus panel (modified Delphi); (6) public website; (7) second external review; (8) final draft and grading of the recommendations; and (9) dissemination and implementation. Details of this study are published elsewhere.²

Focus

These diagnostic imaging guidelines concern adult musculoskeletal disorders of the spine where conventional radiography and specialized imaging studies are deemed useful for diagnostic purposes. Special consideration for manual therapy intervention is integrated within these guidelines.

Objectives

Reasons for developing these guidelines include assisting current and future health care providers to make appropriate use of imaging studies, providing indications for the need of

^a Professor, Chiropractic Department, Universit e du Qu ebec   Trois-Rivi eres, Quebec, Canada.

^b Professor of Radiology, D'Youville College, Buffalo, New York.

^c Professor, Canadian Memorial Chiropractic College, Toronto, Ontario, Canada.

Submit requests for reprints to: Andr e E. Bussi eres, DC, Professor, D epartement Chiropratique, Universit e du Qu ebec   Trois-Rivi eres, C.P. 500, Trois-Rivi eres, Qu ebec, Canada G9A 5H7 (e-mail: andre.bussieres@uqtr.ca).

Paper submitted July 15, 2007; in revised form September 28, 2007; accepted October 14, 2007.

0161-4754/\$34.00

Copyright   2008 by National University of Health Sciences.

doi:10.1016/j.jmpt.2007.11.003

imaging studies according to current literature, and expert consensus, and assisting in optimizing the utilization of limited available resources. These proposed guidelines are intended to reduce unnecessary radiation exposure and the use of specialized imaging studies, increase examination precision and decrease health care costs—all without compromising quality of care.

Target Users/Setting

Intended users of the guidelines are chiropractors and other primary health care providers prescribing diagnostic imaging studies. The setting in which these guidelines may be used include private clinics, outpatient clinics, and hospital emergency rooms.

Target Population

The patient population eligible for guideline recommendations are adult patients presenting with musculoskeletal disorders of the spine. Children and pregnant patients are excluded from these guideline recommendations.

Developers

The proposed guidelines are developed from the results of 9 distinct phases overseen by a research team composed of the 3 investigators with postgraduate education from 3 independent teaching institutions. The guidelines were further developed and peer reviewed by more than 60 chiropractic clinicians, academics, and researchers.

Evidence Collection

Electronic searches in English and French language literature occurred and cross references were repeated on 3 different occasions between 2003 and 2006.

METHODS FOR SYNTHESIZING EVIDENCE

- (a) Literature search and independent literature assessment of spinal disorders: Quality of diagnostic accuracy studies (QUADAS),³ Appraisal of Guidelines Research and Evaluation (AGREE),⁴ and Stroke Prevention and Educational Awareness Diffusion (SPREAD).⁵
- (b) Initial draft: template based on European Commission classification (2001).⁶
- (c) Expert consensus: a 2-round modified Delphi process was used to generate consensus among an international panel of more than 60 experts in musculoskeletal disorders.

Recommendation Grading Criteria

The evaluation tool used was designed by the Scottish Intercollegiate Guidelines Network (SIGN) and adapted by the Stroke Prevention and Educational Awareness Diffusion (SPREAD) group.^{5,7}

Patient Preferences

Condition specific imaging guidelines. Integral to evidence-based health care, decisions regarding the use of imaging studies should be based on the best available evidence, the

experience, and judgment of the clinician, while considering the patient preference. A public member reviewed all documents and provided comments and suggestions.

Stakeholders and Editorial Independence

- (a) *Prerelease review*: Before the release of the guidelines, the reliability of proposed recommendations was tested on specialists both in chiropractic and in medicine as well as on practicing chiropractors.
- (b) *Potential conflict of interest*: The research team involved in the development of these guidelines declare no existing or potential conflict of interest. No investigators have received nor will receive any personal financial benefits or derive any salary from this project.
- (c) Funding sources/sponsors:
 1. Canadian Memorial Chiropractic College Post Graduate Education and Research (2005)
 2. National Institute of Health Student Grant (2006)
 3. Canadian Chiropractic Protective Association (2006)

Updating/Revision

The literature review and the guidelines should be updated every 2 to 3 years.

Potential Benefits and Harm

Selection of appropriate radiologic imaging procedures for evaluation of patients with musculoskeletal disorders of the spine; decrease unnecessary ionizing radiation exposure, decrease costs, and improve accessibility.

Dissemination/Implementation Considerations

Publication; applying to National Guideline Clearinghouse; posting of the electronic document on various websites (malpractice insurance carriers, outpatient teaching clinics); educational intervention strategies (e-learning, community pilot studies); referral guidelines; reinforced by request checking and clinical management algorithms; promotion by national, provincial and state organizations, conferences.

Definitions, Patient Presentations, Recommendations, and Rationale

These topics are integral parts of each 1 of the 3 diagnostic imaging guidelines: lower extremity disorders, upper extremity disorders, and spine disorders. Results of the 9 phases of the research project are published elsewhere.²

PRELIMINARY CONSIDERATIONS AND DISCLAIMER

What Is the Role of These Guidelines?

These evidence-based diagnostic imaging practice guidelines are intended to assist primary care providers and students in decision making regarding the appropriate use of diagnostic imaging for specific clinical presentations. The guidelines are intended to be used in conjunction with sound clinical judgment and experience. For example, other special circumstances for radiographic imaging studies may include: patient unable to give a reliable history; crippling cancer phobia focused on back pain; need for immediate decision

about career or athletic future or legal evaluation; history of significant radiographic abnormalities elsewhere reported to patient but no films or reliable report reasonably available; history of finding from other study (eg, NM or gastrointestinal imaging) that requires spine radiographs for correlation.⁸ Application of these guidelines should help avoid unnecessary radiographs, increase examination precision, and decrease health care costs without compromising the quality of care.

The descriptions of clinical presentations and proposed clinical diagnostic criteria, recommendations for imaging studies, and the comments provided throughout this document are a synthesis of the vast body of literature consulted before and during the various phases of this research project. Where the literature was found to be of poor quality or absent, consensus based on expert opinion was used. Although the investigators and collaborators carefully searched for all relevant articles, it is probable that some have been missed. Furthermore, as many new important studies are published in the near future, these will be incorporated in subsequent revisions of the guidelines and recommendations may change accordingly.

What These Guidelines Do and What They Do Not Do

It should be emphasized that these guidelines were developed with the intent of being used for diagnostic purposes and not for therapeutic purposes such as evaluating and monitoring functional or structural rehabilitation of the spine.⁹ In addition, these guidelines are intended to address issues faced by first contact professionals only. These guidelines do not address all possible conditions associated with musculoskeletal disorders, only those that account for the majority of initial visits to a practitioner.

Like other diagnostic tests, imaging studies should only be considered if (a) they yield clinically important information beyond that obtained from the history and physical examination; (b) this information can potentially alter patient management and; (c) this altered management has a reasonable probability to improve patient outcomes.¹⁰⁻¹²

Investigators and collaborators in the development of these imaging guidelines believe that liability insurance companies, third-party payers, and courts of law should not rely solely on descriptions of patient presentations, proposed recommendations, and/or corresponding comments found throughout the documents, as patient presentations are unique and the application of any guideline always requires clinical judgment and thus needs to be considered in the proper context. In addition, laws and regulations may vary between geographical regions and should be considered when applying the proposed indications for any imaging study.

What Is Evidence-Based Health Care?

Evidence based is about tools, not about rules.¹³ Evidence-based health care is an approach in which clinicians and health care professionals use the current best evidence in making decisions about the care of patients. It involves continuously

and systematically searching, appraising, and incorporating contemporaneous research findings into clinical practice. The overall goal is improving patient care through life-long learning.^{13,14}

Potential Disagreements

There are several reasons for disagreement within a guideline development group. These include differences in interpretation of the research literature, differences in personal experience, and different perceptions of the inherent risks and benefits of a procedure.¹⁵ Divergent or competing guidelines on similar topics serve only to further confuse and frustrate practitioners.¹⁶ In addition, the continued lack of unity among chiropractors hinders its growth by limiting integration and cooperation within the greater health care system. Readers of all guidelines are advised to critically evaluate the methods used as well as the content of the recommendations before adopting them for use in practice.¹⁷

Standard Patient Management Activities

Standard patient management activities, including diagnostic assessment and follow-up, are integral components of every patient encounter.¹⁸ Initial triage of patients with spine disorders is a constant recommendation of various clinical guidelines.¹⁹ Imaging studies are used most practically as confirmation studies once a working diagnosis is determined. The objective is to determine the presence of clinical indicators of serious pathologies (red flags) requiring diagnostic imaging, specialist referral or urgent surgical intervention. When a practitioner recommends that a radiograph or other diagnostic imaging study be performed, and the patient refuses, the patient should be advised of the associated risks and implications and this should be recorded in the patient's records.

Duration of Disorders

In clinical practice, spine disorders are generally divided into categories according to the duration of the patient complaint on initial presentation. These diagnostic imaging guidelines therefore consider the following categories of clinical presentations: acute spine disorder (<4 weeks' duration); subacute spine disorder (4-12 weeks' duration); and persistent/chronic spine disorder (>12 weeks' duration).

Defining Radicular Pain

It is generally accepted that patients with neurologic symptoms and signs (pain radiating below the knee or beyond the elbow, as intense as the low back or neck pain, often radiating into the foot or hand with numbness or paresthesia in a dermatomal distribution with positive nerve root tension signs, abnormal motor power, sensation or deep tendon reflexes consistent with a nerve root involvement) have a slower resolution than patients with uncomplicated spine disorders (mechanical pain that varies with time and activity with no neurologic component and a good general health status). Again, clinical presentations are divided accordingly throughout these imaging guidelines.

Table 1. Thoracolumbar, lumbar, and thoracic spine trauma

Patient presentation	Recommendations	Comments
<p>Adult patient with recent (<2 wk) acute thoracolumbar, lumbar or thoracic spine trauma.</p> <p>Absence of pain, normal ROM, and absence of neurologic deficits (normal lower limb motor power, sensation and deep tendon reflexes, and no pathologic reflexes).⁶</p>	<p>Radiographs not routinely indicated [C]</p>	<p>In a recent survey, 53% of patients consulting chiropractors reported having an injury. The most common reported source was non-work related (43%); work-related injuries accounted for only 16%, and 21% were classified as nonspecific.⁴¹ Physical examination is reliable in this region. When the patient is alert and asymptomatic, injury risk is low and radiographs provide no clinical utility.^{6,42}</p> <p>Increased prevalence of traumatic fractures found in patients >75 yoa. Prevalence of lumbar spine traumatic fracture is <1% but overall prevalence of spine fractures for all age groups is 4%.^{43,44}</p>
<p>Adult patient with thoracolumbar, lumbar, or thoracic spine blunt trauma or acute injuries (falls, motor vehicle accidents [MVAs], motorcycle, pedestrian, cyclists, etc)</p> <p>High-risk screening criteria for spinal injuries include any of the following⁴⁵⁻⁴⁹:</p> <ol style="list-style-type: none"> 1. Back pain 2. Midline tenderness on palpation 3. Distracting painful injury and other high-risk mechanism of injury* 4. Neurologic deficits** 5. Altered consciousness (caused by head trauma, intoxication/ethanol, or drugs) <p><i>Sensitivity of 100 % (95% CI, 98%-100%) but low specificity and low positive predictive value for thoracolumbar injuries.</i></p> <p>* High-risk mechanism of injury: Severe mechanism of injury (falls ≥ 10 ft/3 m, ejection from a motor vehicle, and motor vehicle crashes >50 mph/80 km) and the presence of cervical spine fracture are significantly associated with an increase risk of thoracolumbar spine injury.</p> <p>** Neurologic deficits:</p> <ol style="list-style-type: none"> a. <i>Lumbar spine injuries:</i> positive nerve root tension signs decreased motor power, sensation, and/or deep tendon reflexes. b. <i>Thoracolumbar and thoracic spine injuries:</i> lower limb spasticity, presence of pathologic reflexes (Babinski sign/clonus), altered sensations below the injury (sensory level), and altered proprioception (vibration/position sense of the feet), loss of anal sphincter tone. <p>• Patient with spinal injuries may suffer progressive neurologic deterioration, and imaging again has a role in their diagnosis and management.⁷</p> <p>For osteoporotic fracture risk, please see Pain—Osteoporotic Vertebral Collapse section of the Adult Thoracic Disorders Guideline.</p>	<p>Radiographs indicated [B] Lumbar AND thoracic spine: AP (or PA), lateral views N.B. The high-risk screening criteria for spinal injuries was developed to help determine the need for plain film radiography in the emergency room. The lower prevalence of serious post traumatic injuries in ambulatory care likely influence the sensitivity of such instruments. Nonetheless, it appears reasonable to recommend their use in primary care as well as well.</p> <p>Special investigations [C]</p>	<p>Conventional radiography is highly specific but has a sensitivity of ≈ 0.70 and a predictive value of 0.92⁴⁵ Thoracic and lumbar spinal injuries affect 2%-3% of blunt trauma victims with a 40%-50% incidence of neurologic deficits.⁷</p> <p>Major spine traumas are typically seen in hospital settings and mandate immobilization pending radiographic evaluation. Missed spinal injuries resulting in neurologic deficit continue to occur in major trauma centers. Older age (>50), high impact accidents, patients with insufficient imaging, and misread or poor quality radiographs are at highest risk of missed injuries.⁵⁰ The Thoracolumbar Injury Severity Scale (TLISS) may be helpful in the referral decision making process. The scale is based on the mechanism of injury, the integrity of the posterior ligamentous complex, and the neurologic status.⁵¹⁻⁵³ (See Appendix C(1) for details)</p> <p>Advanced imaging and specialist referral recommended:</p> <ol style="list-style-type: none"> a. In presence of a fracture as suggested by history, examination, and/or radiograph. Special attention required in presence of complex lesions and for suspected ligamentous instability or neural injuries; b. If conventional radiographs are difficult to interpret; c. In the absence of clinical improvement after 4-6 wk of therapy; d. Function does not improve or deteriorates; e. Patient has persistent S&S. <ul style="list-style-type: none"> • CT scan (multidetector/multislice CT and spiral CT) highly sensitive (99.3%-100%); specificity of 97%; negative predictive value of 92%^{47,48,54,55} • MRI provides the best evaluation for soft tissue pathology, intrinsic or compressive spinal cord damage and multilevel vertebral fractures. MRI needed when conventional radiographs and CT do not explain patient's symptoms, and when there is a possibility of epidural hematoma or traumatic disc herniation⁵⁶

Table 1 (continued)

Patient presentation	Recommendations	Comments
<p>Adult patient with posttraumatic chest wall pain</p> <p>Minor trauma⁶ Major trauma⁶</p>	<p>Radiographs not routinely indicated [D] Radiographs indicated [GPP] PA, lateral chest radiographs, Specific rib radiographs (AP, oblique)</p> <p>Additional views: PA chest in full expiration, Thoracic and /or lumbar spine views</p> <p>Special investigations [GPP]</p>	<p>Although difficult to visualize, presence of a rib fracture would alter the treatment plan in manual therapy.</p> <p>Chest radiographs frequently underestimate the severity and extent of chest trauma, and in some cases, fail to detect the presence of injury. CT is more sensitive than chest radiography in the detection of pulmonary, pleural, and osseous abnormalities.⁵⁷ Penetrating injuries or clinical findings suggestive of significant air exchange difficulty urgently require more extensive work-up.</p> <p>Advanced imaging and specialist referral recommended:</p> <ul style="list-style-type: none"> • CT for sternum injury, pulmonary, pleural, and osseous abnormalities⁵⁷
<p>Adult patient with pelvis and sacrum trauma (including falls with inability to bear weight)⁶</p> <p>The diagnosis of sacral fractures is frequently missed or delayed, and the treatment is controversial, with significant variations in recommendations regarding nonoperative treatment, neural decompression, and internal fixation.⁵⁸</p> <p>Posterior pelvis physical examination, especially palpation, may accurately detect injuries to the posterior pelvic ring.⁵⁹ Look for ecchymosis. However; physical examination may not be reliable for other pelvic injuries.</p>	<p>Radiographs indicated [D] AP Pelvis and lateral hip “frog leg”</p> <p>Additional views: (1) Lateral lumbar view to better visualize the lateral sacrum (2) Angulated AP or PA sacrum view (15°-45° tube angulation)</p> <p>Special investigations [D]</p>	<p>Be aware that femoral neck fracture may not be visible on initial radiographs even with a good lateral view of the hip.</p> <p>Factors that may impact treatment outcomes include the level and type of sacral fracture, lumbosacral junction and sacroiliac joint involvement, and associated pelvic ring injury.⁵⁸</p> <p>Advanced imaging and specialist referral recommended</p> <ul style="list-style-type: none"> • NM, MRI, or CT may be helpful if radiographs are normal or equivocal.⁶ • Imaging measurement parameters used to evaluate sacral injuries may include AP sacral fracture displacement (axial CT of the pelvis), vertical sacral fracture displacement (coronal CT reconstruction), AP translation and kyphotic angulation (sagittal CT reconstruction), and degree of central canal involvement and foraminal encroachment (axial, coronal, and sagittal CT reconstruction)⁵⁸
<p>Coccyx trauma and coccydynia</p> <p>Consider taking radiographs if unrelieved by initial rest or if manual treatment of this area is contemplated.</p> <ul style="list-style-type: none"> • Obesity, a history of trauma, and transient exacerbation of the pain when standing up from sitting may be associated with posterior dislocation. • Coccyx pain may be referred from a fracture of the distal sacrum. In doubt, consider views of the sacrum if distal sacrum fracture is suspected.⁶⁰⁻⁶² 	<p>Radiographs not routinely indicated [C] (Spot AP, lateral coccyx)^{6,63}</p> <p>Additional views: AP, lateral sacrum Dynamic sitting lateral views of the coccyx⁶⁴</p>	<p>Normal appearance often misleading.⁶³</p> <p>Look for fractures, dislocations, and hypermobility (coccygeal mobility between 5° and 25° is considered normal).⁶¹</p>

Table 2. Cervical spine trauma

Patient presentation	Recommendations	Comments
<p>Adult patient with acute neck injury and negative CCSR (Canadian Cervical Spine Rule for Radiography in Alert and Stable Trauma Patients)⁶⁵⁻⁶⁸</p> <p>Conventional radiographs are unlikely to show clinically significant injuries when <i>all Canadian Cervical Spine Rule</i> criteria are fulfilled⁶⁵⁻⁶⁹</p> <p>(A) Any high-risk factors in alert and stable patient?</p> <ol style="list-style-type: none"> 1. Age >65 (see NEXUS in comment section) 2. Dangerous mechanisms of injury* 3. Paresthesias in extremities <p>(B) Any low-risk factors that allow ROM assessment?</p> <ol style="list-style-type: none"> 1. Simple rear end collision** 2. Patient seated in the waiting room 3. Ambulatory at one time since trauma 4. Delayed cervical pain onset 5. Absence of midline cervical tenderness <p>(C) ROM assessment: Is patient able to actively turn his/her head to 45° in both directions?</p> <p>* Dangerous injury mechanism:</p> <ul style="list-style-type: none"> • Fall >3 ft/5 stairs • Axial cranial force (such as diving) • Road accident >100 km/h, ejected from vehicle or rollover • Motorized recreational vehicle (ATV, snowmobile, etc) <p>** Rear end collision excludes:</p> <ul style="list-style-type: none"> • Being pushed into oncoming traffic • Hit by a bus or large truck • Rollover • Being hit by high speed vehicle (≥100 km/h) <p>Sensitivity 100% (95% CI, 98%-100%) and specificity 42.5% (95% CI, 40%-44%).^{65,66}</p>	<p>Radiographs not routinely indicated [B]</p> <p>N.B. The CCSR and NEXUS instruments are highly sensitive clinical decision rules developed to help determine the need for plain film radiography in the emergency room. The lower prevalence of serious post traumatic injuries in ambulatory care likely influence the sensitivity of such instruments, nonetheless, it appears reasonable to recommend their use in primary care as well.</p>	<p>Whiplash incidence is between 70 and 360 cases per 100 000.⁷⁰ Traumatic unstable injuries are rare (<3%). The CCSR⁶⁵ is superior to physician judgment, more sensitive and specific than the National Emergency X-Radiography Utilization Study criteria/NEXUS^{66,67}</p> <p>In head-turned rear impacts up to 8 g peak acceleration, the alar, transverse, and apical ligaments are not at risk for injury.⁷¹</p> <p>Although patients >65 yoa are more prone to cervical spine fractures, data from NEXUS indicates that being >65 is not, by itself, a risk factor if all criteria are considered.^{7,72} (Sensitivity 100% (95% CI.) Cervical spine radiographs are not indicated <i>if</i> patients fulfill <i>all 5 NEXUS low-risk criteria</i> for head and/or cervical spine trauma⁶⁷:</p> <ol style="list-style-type: none"> (1) Normal level of consciousness (may be temporarily confused, have mild dizziness or headache) (2) No evidence of intoxication (3) Absence of tenderness on palpation of the midline of the neck (4) Absence of focal neurologic deficit (5) Absence of painful traction injury <p>Rare injuries may be missed, when the CCSR or NEXUS is properly applied, findings are generally considered not clinically significant^{65,67}: spinous or transverse process fracture, <25% single anterior compression fracture, isolated avulsion injury not associated with ligamentous injury, type I odontoid fracture, end plate fracture, osteophyte fracture excluding teardrop or corner fracture, trabecular bone injury.</p> <p>NB. "Not clinically significant missed injuries" may be relevant to manual therapy practitioners. However, considering soft tissue and bone healing time; recommendations include early activation and a gradual progression from mobilization to SMT. SMT recommended only when patient can tolerate pre-SMT position test.⁷³⁻⁸²</p> <p>Of interest, the hypothesis that early aggressive conservative care promotes faster recovery was not supported in a recent population-based cohort study⁸³</p>

Table 2 (continued)

Patient presentation	Recommendations	Comments
<p>Adult patient with acute neck injury and positive CCSR (Canadian Cervical Spine Rule for Radiography in Alert and Stable Trauma Patients)⁶⁵⁻⁶⁸</p> <p>Any adult patient not fulfilling all criteria of the CCSR must undergo cervical spine radiographs:</p> <p>(A) Age >65; dangerous mechanisms of injury; paresthasias in extremities</p> <p>(B) Not a simple rear end collision; patient unable to sit in the waiting room; not ambulatory at one time since trauma; immediate cervical pain onset; presence of midline cervical tenderness</p> <p>(C) Patient unable to actively turn his head to 45° in both directions</p> <p>Both physical and psychological factors play a role in recovery or nonrecovery from whiplash injury.^{84,85} Finally, an important association exists between a history of whiplash associated disorders (WAD), pain intensity and disability and comorbidity (headache, LBP, digestive, and cardiovascular disorders).⁸⁶⁻⁸⁸</p> <p><i>Patient history may include:</i></p> <ul style="list-style-type: none"> • Patient age and sex • Pain location/radiation • Delay in S&S appearance • Mechanism of injury • Use of relevant psychometric questionnaires: NDI for neck disability, psychological distress (General Health Questionnaire 28, fear of movement/re-injury (TAMPA Scale of Kinesophobia, TSK), acute post-traumatic stress (impact of events scale and general health and well being [Short Form 36, SF-36]). • a complete system review <p><i>Patient physical examination may include:</i></p> <ul style="list-style-type: none"> • motor function* including cervical ROM and manual spinal palpation • a thorough neurologic examination, including cranial nerves, motor power, deep tendon reflexes, and local and general sensory testing. <p>The above information helps determine the severity of the injury and prognosis. See Appendix C(2) for a modified classification system for acute WADs including physical and psychological impairments.</p> <p>*Motor system function testing may also include muscle recruitment pattern, joint position error, EMG recording, kinematic analysis of the cervical spine, and quantitative sensory testing/algometry.⁸⁹</p>	<p>Radiographs indicated [B] APOM, AP lower cervical, neutral lateral</p> <p>N.B. The CCSR and NEXUS instruments are highly sensitive clinical decision rules developed to help determine the need for plain film radiography in the emergency room. The lower prevalence of serious post traumatic injuries in ambulatory care likely influence the sensitivity of such instruments, Nonetheless, it appears reasonable to recommend their use in primary care as well.</p> <p>Additional views [GPP] CT now considered superior to and should replace oblique, pillar, dynamic flexion/extension (F/E) in suspected fracture.</p> <p>Special investigations [C] CT is becoming the primary investigation for high risk patients on an emergency basis (more accurate, faster, and need less patient mobilization compared with conventional radiography).^{54,90,91} Harboview CT Screening Criteria (prediction rule):</p> <ul style="list-style-type: none"> • High energy trauma • Severe head injury • Focal neurologic deficit • Associated injury • Age (>50 y) • If patient >65 yoa, consider moderate-energy mechanism⁹² 	<p>Conventional radiographs may be difficult to assess; view must clearly include C7-T1 and the odontoid.⁹³</p> <p>Pertinent 3 view radiological findings:</p> <ul style="list-style-type: none"> • ADI >3 mm in adults (>5 mm in children) • Acute intradiscal vacuum cleft • Acute angular segmental kyphosis • Widened spinous process interspaces • Loss of facet parallelism • Presence of prevertebral edema without fracture.⁹⁴ <p>According to the Yale criteria, a sagittal translation >3.5 mm or >11° of sagittal rotation in adults indicates segmental instability & requires surgical fusion.⁹⁴</p> <p>See Appendix C(2) for a new classification of lower cervical spine injuries (Cervical Spine Injury Severity Score)⁹⁵</p> <p>Cervical spine immobilization in awake patient with persistent pain or tenderness and normal initial standard 3-view series (including supplemental CT as necessary) to exclude ligamentous instability may be discontinued after either⁹⁶:</p> <p>(a) normal dynamic flexion/extension (F/E) radiographs as achieved by the patient with no assistance and under proper supervision (consider fluoroscopy)</p> <p>(b) a normal MRI study is obtained within 48 h of injury NB. F/E films may add little value to the acute evaluation of patients with blunt trauma due to muscular spasm. Consider F/E imaging delayed to 14-28 d postinjury when muscular spasm has resolved.^{97,98}</p> <p>Specialist referral is highly recommended in blunt trauma and trauma patients with neurologic signs or fractures/dislocations. In moderate- to high-risk trauma, delays in diagnosis of clinically significant C/S injuries range between 5% and 23%, most of which used conventional radiography as initial screening modality. Neurologic deterioration occurs in 10%-50% of these patients.⁴⁷</p> <ul style="list-style-type: none"> • Consider MDCT (multidetector/multislice CT) or MRI in blunt trauma or if initial radiographs are difficult to interpret or in presence of complex lesions. • MRI for suspected ligamentous instability or neural injuries. In the presence of neurologic S&S, examine patient for signs of acute myelopathy. EMG may be useful.⁶ • Consider CT for occult fractures.⁹⁸

(continued on next page)

Table 2 (continued)

Patient presentation	Recommendations	Comments
<i>The reader is referred to Foreman & Croft textbook Whiplash Injuries⁷³ and the Quebec Task Force on Whiplash-Associated Disorders⁷⁴ for additional reading on WAD</i>	However, MRI is an acceptable option.	The presence of any one of the following parameters places the patient in the high-risk category (>5% risk of cervical spine fracture) and indicates that the patient should undergo helical CT ⁹⁰ : <ol style="list-style-type: none"> 1. High-speed (35 mph [56 km/h] combined impact) MVA 2. Crash with death at scene of MVA 3. Fall from height (10 ft [3 m]) 4. Significant closed head injury (or intracranial hemorrhage seen on CT) 5. Neurologic symptoms or signs referred to the cervical spine 6. Pelvic or multiple extremity fractures See Cervical Artery Dissection section for further details on prior history of major trauma.

Table 3. Nontraumatic lumbar spine disorders

Patient presentation	Recommendations	Comments
<p>General background information</p> <p>Standards of care dictate that careful history & physical examination be done routinely for all patients. See Lurie (2005)⁹⁹ for a comprehensive review of the value of the history & physical examination for LBP.</p> <p>Between 41% and 44% of initial chiropractic visits are for low back–related complaints, of which 45% are acute in origin.^{41,100}</p>		<p>Low back pain on initial presentation is typically classified according to the duration of complaint and type of presentation: uncomplicated/nonspecific LBP, associated with neurologic deficits or to some underlying pathology.¹⁰¹</p> <p>Duration of complaint: Acute LBP: <4 wk of duration Subacute LBP: 4-12 wks' duration Persistent/chronic LBP: >12 wks' duration</p> <p>The cumulative 6-mo incidence of LBP is 18.6% (95% CI, 14.2-23.0) with 17.2% having mild pain (95% CI, 12.9-21.5), 1.0% having intense pain (95% CI, 0.0-2.2), and 0.4% disabling pain (95% CI, 0.0-1.0).¹⁰²</p> <p>Annual course of back pain is as follows¹⁰²:</p> <ul style="list-style-type: none"> • 27.3 of low back complaints resolve • 29% improve • 35.4% persist • 9.3 get worse • 31.3% recur
<p>Adult patient with acute uncomplicated* LBP (<4 wks' duration)</p> <p>* Uncomplicated definition: nontraumatic LBP without neurologic deficits or indicators of potentially serious pathologies—(see red flag list for details).</p>	<p>Radiographs not initially indicated^{42,101,108,109} [B]</p>	<p>Routine use of lumbar spine conventional radiography is not indicated because of very low incidence of unexpected findings on radiographs (only 1 in 2500 radiographs), high radiation dose to gonads, high cost/benefit ratio, and poor association between patient findings and LBP (ie, not specific). Anomalies of no proven clinical relevance include block vertebrae, spina bifida occulta, mild scoliosis, and facet tropism; none of which are considered contraindications to SMT.^{40,43,110-136}</p>
<p>Mechanical LBP accounts for the vast majority of LBP seen in primary care setting. Mechanical designates anatomical or functional abnormality without an underlying pathology.⁴⁴</p> <p>For most young or middle-aged adults, early diagnostic evaluation of low back complaints may focus on 3 basic questions: diagnostic imaging is infrequently required.⁴⁴</p> <p>1. Is there underlying systemic disease?</p>	<p>Special investigations not indicated [B]</p> <p>Outside the setting of suspected systemic disease or neurologic compromise, radiography, scintigraphy, computed-tomography (CT), and MRI add very little to the diagnostic evaluation of LBP.^{42,99}</p>	<p>Several factors influence the reliability and validity of biomechanical and postural analyses on conventional radiographs: Morphologic asymmetries, geometric and positional distortions, measurement errors, poor correlation with symptoms and lack of convincing clinical usefulness.^{43,105,106,121,122,137-145}</p> <p>Acute LBP is generally due to musculoskeletal conditions that respond favorably to SMT.^{146,147} A multidisciplinary panel recommended 2 trial courses of 2 wk each, using</p>

Table 3 (continued)

Patient presentation	Recommendations	Comments
<p>2. Is there neurologic impairment that might require surgical intervention?</p> <p>3. Is social or psychological distress amplifying or prolonging the pain?</p> <p>There is strong evidence that in nonspecific LBP, conventional radiography and MRI findings do not correlate with clinical symptoms or work capacity.^{103,104}</p> <p>In the <i>absence</i> of red flags as revealed by the history and physical exam, radiography in patients with LBP of at least 6 wk duration is not associated with improved physical functioning, severity of pain, disability, or overall health status. Conventional radiography may somewhat improve patient satisfaction. However, minor psychological improvement should be balanced against radiation dose, higher number of doctor visits, and increased disability at 3 mo.¹⁰⁵⁻¹⁰⁸</p>		<p>alternative manipulative procedures.^{148,149} Lumbar spine radiography recommended in the absence of expected treatment response or worsening after 4-6 wk.</p> <p>Serious adverse events are unlikely to occur with chiropractic treatment for LBP.¹¹⁵ Contraindication for lumbar SMT includes progressive neurologic deficits, S&S of cauda equina syndrome, suspicion of abdominal aortic aneurysms or destructive and bone softening diseases such as Paget's disease, and severe osteoporosis. (<i>See LBP with red flag's for details</i>).</p>
<p>Adult patient with uncomplicated subacute (4-12 wks' duration) or persistent LBP (>12 wks' duration) AND no previous treatment trial.^{63,101}</p> <p>Consider a clinical evaluation of risk factors for chronic LBP (<i>yellow flags</i>). Look for functional disability, significant depression, and risk profile for delayed recovery.</p> <p>A trial of 4-6 wk of conservative care is appropriate before radiographs.</p>	<p>Radiographs not initially indicated [B]</p>	<p>As with all cases, recent previous available radiographs should be reviewed by the clinician. Overall, studies have not shown any significant correlation between posture and LBP. Degenerative processes (disc & facet) are common & usually related to age (prevalence >62% in patients >55 yoa). Degenerative processes may account for 10% of all LBP. Abdominal aortic calcification may be associated with lumbar disc degeneration, back pain. However, even if DJD and DDD are present, clinical management remains essentially unchanged.^{42,43,113,121,150-156}</p>
<p>Adult patient with nontraumatic acute LBP (<4 wks' duration) AND sciatica (no red flags)^{6,63,101}</p> <p>The first clinical clue to neurologic impairment usually is a history of sciatica: sharp pain radiating down the posterior or lateral aspect of the leg, often associated with numbness or paresthesia.</p>	<p>Radiographs not initially indicated [B]</p>	<p>The natural course of acute sciatica caused by lumbar disc herniation (LDH) is benign as most patients improve during the first 4 wk,¹⁵⁷ whereas 5%-25% are recurrent or persistent.^{158,159}</p> <p>Patient S&S should not increase while in the premanipulative side posture, in which case SMT should not be attempted. Treatment approach should be adapted to further reduce complication risks.^{160,161} Ongoing clinical assessment is necessary each visit to detect any early signs of CES.^{162,163} — (<i>see red flags for details</i>).</p>
<p>Specific clinical diagnoses:</p>		
<p>Common causes of sciatica</p> <p>Historical and physical findings are the key factors in assessing neurologic compromise.⁹⁹</p> <p>(A) Suspected LDH^{44,163-170}:</p> <ul style="list-style-type: none"> • Symptomatic LDH represents 4%-5% of causes of LBP • Risk factors for LDH include: men (1.6 times more likely), middle-aged (35-54 yoa), repetitive heavy lifting, current smoking, obesity (high BMI), and type of occupation. • Predominantly leg pain, typically involving the foot • >95% of clinically important LDH involve the L5 or S1 nerve root. • Straight leg raise is sensitive (0.8) but nonspecific (0.5); 	<p>Radiographs not initially indicated [B]</p> <p>Unless patient age >50 or has progressive neurologic deficits</p> <p>LDH cannot be seen on conventional radiography.</p> <p>Special investigations not initially indicated [C] (<i>see Suspected causes of sciatica for details</i>)</p>	<p>The natural course of acute sciatica caused by LDH is favorable.¹⁵⁷ A trial of conservative care may first be attempted.^{171,172} SMT appears to be a safe & effective treatment for nonprogressive S&S of lumbar disc protrusion, especially when compared to common treatments for LDH, such as NSAIDs and surgery.^{160,161,173-180}</p> <p>It is estimated only 2% of all patients who have LBP will undergo surgery for disc herniation, representing 5%-10% of all cases of LDH.^{166,176}</p> <p>MRI <i>within the first 4-6 wk</i> of suspected LDH is necessary only if there is progressive neurologic deficit or intolerable pain levels despite conservative care and surgical referral is planned.⁹⁹ Disc abnormalities may be classified on MRI according to the following:</p>

(continued on next page)

Table 3 (continued)

Patient presentation	Recommendations	Comments
<ul style="list-style-type: none"> • Crossed SLR is highly specific (0.8) but not sensitive (0.3) • Look for lower limb sensory, motor, and reflex deficits <p>See Appendix C(3) for a detailed list of clinical criteria for LDH.</p>		<p>Modic classification¹⁸¹:</p> <p>4A—herniated disc protrusion with an intact annulus; 4B—extrusion with rupture of either the annulus or the posterior longitudinal ligament, or both (extrusion appears to be rare in asymptomatic adults); 4C—rupture of the annulus and the posterior longitudinal ligament with sequestration of a disc fragment in the spinal canal.</p>
<p>(B) Suspected degenerative spondylolithesis/lateral stenosis</p> <ul style="list-style-type: none"> • Back pain with or without leg pain • Increase pain with activity • S&S with or without neurologic deficit 	<p>Radiographs indicated if patient age >50 or has progressive neurologic deficits: PA (or AP), lateral lumbar views [GPP]</p> <p>Special investigations not initially indicated [C] (see <i>Suspected causes of sciatica</i> for details)</p>	<p>If clinical signs suggest instability: <i>please see special investigations in: Re-evaluation of LBP for critical exclusionary diagnoses.</i></p> <ul style="list-style-type: none"> • A trial of conservative care may first be attempted
<p>(C) Suspected lumbar degenerative spinal stenosis (3% of all LBP)^{99,182,183}:</p> <ul style="list-style-type: none"> • MC >65 yoa (sensitivity of 0.7; specificity of 0.69) • Neurogenic claudication: severity of back and leg pain commonly increase with ambulation/standing/walking downhill and improve while seated (flexed spine). Absence of pain when seated, rather than improved pain, has low sensitivity (0.46) but is very specific (0.93); • Variable neurologic deficit (numbness, weakness) 	<p>Radiographs indicated if patient age >50 or has progressive neurologic deficits: PA (or AP), lateral lumbar views [C]</p> <p>Special investigations not initially indicated [C] (see <i>Suspected causes of sciatica</i> for details)</p>	<p>Conventional radiography is of limited value in assessing spinal stenosis. As many as 20% of asymptomatic adults >60 yoa have imaging evidence of spinal stenosis and the prevalence of symptomatic stenosis is unknown.^{44,99} Spinal stenosis may be caused by bone (facet hypertrophy, osteophytes, etc), soft tissue (bulging disc or thickened ligamenta flava, etc), or both.</p> <ul style="list-style-type: none"> • A trial of conservative care may be attempted first.
<p>Suspected causes of sciatica:</p> <p>(A) Lumbar disc herniation (B) Degenerative spondylolithesis/lateral stenosis (C) Lumbar degenerative spinal stenosis</p> <p>Results from specialized imaging should always be interpreted in light of the clinical findings. Inappropriate utilization of these highly sensitive examinations will produce false-positive results which may result in labeling of patients and contributing to an unfavorable prognosis.^{99,101,184}</p>	<p>Special investigations not initially indicated^{6,166,185} [C]</p>	<p>Comanagement or specialist referral recommended even if conventional radiographs are unremarkable:</p> <ol style="list-style-type: none"> 1. After failed conservative therapy (4-6 wk) 2. For preoperative planning; 3. If neurologic status is deteriorating (progressive deficit, disabling leg pain)^{186,187}; <ul style="list-style-type: none"> • New disc extrusion and nerve root impingement may be the most important findings on MRI^{135,188} • Substituting rapid MRI for radiographic evaluations in the primary care setting may offer little additional benefit to patients, and it may increase the cost of care because of the increased number of spine surgeries that patients are likely to undergo.^{134,188,189} • MRI sensitivity (0.6-1.0) & specificity (0.72-0.99) for LDH is slightly higher than those for CT (0.62-0.9 & 0.7-0.87) but very similar for the diagnosis of spinal stenosis (0.9 & 0.72-0.99)⁴⁴
<p>Adult patient re-evaluation in the absence of expected treatment response or worsening after 4 to 6 wk</p> <p>Should patient fail to improve as expected or marginally improve within 4-6 wk of initial evaluation, the clinician must review history and physical findings and request appropriate diagnostic imaging studies.^{42,44,99,101,108,109,189-191}</p>	<p>Radiographs indicated [B] PA (or AP), lateral lumbar views</p> <p>Additional views not routinely indicated [C]</p> <p>a. Lateral lumbosacral spot</p>	<p>Probability of encountering disease requiring specialized therapy in patients with LBP using conventional radiography is <0.2 in 100 and the probability of tumors or infection would be <1 in 1000.^{43,196,197}</p> <p>a. If poor visualization of lumbosacral junction on lateral view</p>

Table 3 (continued)

Patient presentation	Recommendations	Comments
<p>Adaptation of the plan of management partly relies on the ability to predict patient disability status or long-term return to work status.¹⁹¹⁻¹⁹⁵ Patient perception of the likelihood of returning to his/her usual activities should be measured after 4 wk of disability or at the time of first consultation if the patient has a history of persistent spinal pain with prolonged disability. Validated questionnaires are available to help clinicians.¹⁰¹</p>	<p>b. Oblique</p> <p>c. Lateral flexion films may be indicated in scoliosis evaluation</p> <p>Special investigations [C]</p>	<p>b. Rarely adds clinical information & doubles radiation exposure to patient.¹⁰⁸ Sometimes used to confirm presence of spondylolysis suspected from the AP & lateral films (also for suggestion of malignancy, fracture, etc).</p> <p>c. Low reliability of lateral flexion film for biomechanical assessment.¹⁹⁸ Lateral bending films useful for scoliosis evaluation (see Scoliosis section).</p> <p>Comanagement or specialist referral recommended even if conventional radiographs are unremarkable AND anyone of the following:</p> <ol style="list-style-type: none"> 1. In presence of a potentially serious pathology as suggested by the patient history, examination and/or radiograph; 2. After failed conservative therapy (4-6 wk), 3. If patient neurologic status is deteriorating (progressive deficit, disabling leg pain); 4. If clinical signs suggest instability. Presumed instability is loosely defined as >10° of angulation or 4 mm of vertebral translation on flexion and extension lateral radiographs. However, diagnostic criteria, natural history, and surgical indications remain controversial.⁴⁴ 5. For preoperative planning; <p>While MRI is excellent for identifying tumor, infection, and nerve compression, it can be overly sensitive with regard to degenerative disease findings and frequently displays pathology that is not responsible for the patient’s symptoms.¹⁹⁹</p> <p>Negative findings (MRI/CT /NM) may be helpful. Structural variables on both MRI and discography have a weak association with back pain episodes and no association with disability or future care.^{188,200}</p> <p>See Carragee and Hannibal (2004)¹⁹⁹ and Saal (2002)²⁰¹ for a review of diagnostic testing in chronic LBP.</p>
<p>Adults with complicated (ie, “red flag”) LBP & indicators of contraindication to SMT (relative/absolute):</p> <p>Presence of the following indicator(s) should alert the clinician to possible underlying pathology.^{38,42,44,63,99,112,114,167,185,197}</p> <p>N.B. Presence of a red flag alone may not necessarily indicate the need for radiology.</p> <ul style="list-style-type: none"> • Patient <age 20 and >age 50, particularly with S&S suggesting systemic disease** <p>Other authors recommend waiting 7 wk in untreated 1st episode of LBP and age over 65 as criteria for radiography²⁰³</p> <ul style="list-style-type: none"> • Absence of expected treatment response or worsening after 4-6 wk • Significant activity restriction >4 wk • Nonmechanical pain (unrelenting pain at rest, constant or progressive S&S) 	<p>Radiographs indicated [B]</p> <p>PA (or AP), lateral lumbar views.</p> <p>Main purpose of lumbar spine radiographs is to exclude LBP caused by: malignancies (sensitivity of 0.6, specificity of 0.95-0.99); infective spondylitis (sensitivity of 0.82, specificity of 0.57); inflammatory SpA (low sensitivity for AS [0.25-0.45], but highly specific); fractures; and instability (sensitivity of 0.51-0.99, specificity of 0.68-0.99). High false-positive and false-negative rates are reported when assessing segmental instability by radiography⁹⁹</p>	<p>**Risks of having a serious pathology may be higher before the age of 20 or over the age of 55. Particular attention to indicators of possible underlying pathology should be given for patients in these age categories.^{6,114,208}</p> <p>The major diagnostic task is to distinguish the vast majority of uncomplicated LBP from the 5% with serious underlying diseases or neurologic impairments.⁴⁴</p> <p>Probability of encountering disease requiring specialized therapy in patients with LBP using radiographs is <0.2 in 100 and the probability of tumors or infection would be <1 in 1000.^{43,121,185}</p> <p>Normal radiographs may falsely reassure clinicians as important causes of LBP are not easily identified with conventional radiographs (ie, not sufficiently sensitive)¹¹⁶</p>

(continued on next page)

Table 3 (continued)

Patient presentation	Recommendations	Comments
<ul style="list-style-type: none"> • Suspected inflammatory—spondyloarthropathies (SpA) (0.3% of all LBP): chronic LBP and asymmetric peripheral arthritis, predominantly of the lower limbs, are the leading symptoms for SpA. Features of inflammatory back pain include significant morning stiffness (>1 h); pain duration ≥3 mo; persisting motion restriction; gradual onset ≤age 40; peripheral joint involvement; UTI, urethral discharge, iritis, skin rash. SI stress tests poorly reproducible. <p>AS is regarded as the most severe subgroup. A new set of criteria for inflammatory back pain has been proposed which gives a sensitivity of 70% and specificity of 81% for the diagnosis of ankylosing spondylitis (AS)²⁰⁴:</p> <p>Individual features of inflammatory back pain (IBP)</p> <ul style="list-style-type: none"> • Morning stiffness for >30 min • Improvement of back pain with exercise but not with rest • Awakening in the second half of the night due to back pain • Alternating buttock pain <p>Presence of 2 features—positive likelihood ratio for AS is 3.7 Presence of 3 features—positive likelihood ratio for AS is 12.4</p> <p>** None of the IBP parameters alone are sufficient for a diagnosis of AS.²⁰⁵</p>	<p>Additional views [C]</p> <p>Spot AP or PA angled lumbosacral, oblique SI views</p> <p>Special investigations in complicated LBP [B]</p> <p>Even if conventional radiographs are negative:</p>	<p>Additional views may be considered in the presence of inflammatory spondyloarthropathy symptoms, although SI joints usually adequately demonstrated on AP lumbar spine.</p> <p>The modified New York criteria for AS diagnosis require the presence of radiographic sacroiliitis to classify the patient as having AS. However, plain radiographic abnormalities of the SI joints are a relatively late feature of disease.²¹³ New approaches for an earlier diagnosis of axial and peripheral SpA have been proposed recently. A combination of clinical, laboratory, and imaging parameters is usually necessary.²¹²</p>
<ul style="list-style-type: none"> • Suspected compression fracture (±4% of all LBP): Recent significant trauma (any age), Hx of repetitive stress of sufficient severity; Hx of high-risk osteoporosis; • Severe onset of pain with minor trauma in patients ≥age 50 (higher risk >age 70 with a likelihood ratio of 5.5), Hx of prolonged corticosteroids intake (>7.5 mg/d >3 mo) has a likelihood ratio of 12; structural deformity^{99,206}—a complaint of LBP in persons at risk may require both thoracic and lumbar radiographs.²⁰⁷ See <i>Osteoporotic Vertebral Collapse in the Thoracic Spine</i> section for details. • Suspected neoplasia (0.7% of all LBP): Considerable LBP starting ≥age 50, Hx of cancer/carcinoma in the last 15 y (likelihood ratio of >15), unexplained weight loss (>4.5 kg over 6 mo), failure of conservative care (4 wk). A combination of the above 4 features has a 100% sensitivity for cancer. Other features include no relief with bed rest (sensitivity of 0.90, but low specificity); ESR >50 mm/h, sciatica, systemically unwell; lymphadenopathy; dermopathy (melanoma).^{6,99,203} 	<p>a. MRI is generally the preferred investigation.</p> <p>b. CT may be needed for bony details. (especially multiplanar reformatted images)</p>	<p>MRI or CT useful in the detection of bone and soft tissue structures. The choice of study depends on the current clinical question, availability of equipment, and costs.²¹⁴</p>
<p>Suspected infection (0.01% of all LBP): age >50, documented fever (>38.3°C/101°F) for >3 wk</p>	<p>Advanced imaging and specialist referral recommended even if conventional radiographs are unremarkable:</p> <ol style="list-style-type: none"> 1. In the presence of a potentially serious pathology as suggested by the patient history, examination, and/or radiograph; 2. In the absence of clinical improvement after 4-6 wk of therapy; 3. If function does not improve or deteriorates; 4. If patient neurologic status is deteriorating (progressive deficit, disabling leg pain); 5. With painful or progressive structural deformity 6. For unstable segment (spondylolisthesis or pathological process); 7. When patient has persisting S&S; 8. In complication from treatment (possible fracture, new/progressive neurologic deficit, considerable pain, or disability, etc) 	<ul style="list-style-type: none"> • Suspected cancer: MRI is especially useful for marrow-based lesions. Sensitivity is highest for MRI (0.83-0.93) and radionuclide scanning (0.75-0.98); specificity was highest for MRI (0.9-0.97) and radiography (0.95-0.99).⁴⁴ • Suspected infection: MRI is more accurate than radiography and bone scan (sensitivity of 0.96 & specificity of 0.92) for suspected osteomyelitis, spondylodiscitis, septic disc, paraspinous abscess, epidural abscess. • Suspected acute inflammation in sacroiliac joints and spine: Contrast-enhanced MRI is both sensitive & specific for active sacroiliitis (0.99 & >0.95).^{99,212,213} Although more sensitive

Table 3 (continued)

Patient presentation	Recommendations	Comments
<p>and no established diagnosis despite appropriate investigation for 1 week (specificity of 98% with likelihood ratio of ± 25, but sensitivity of 50%); elevated ESR >20 mm/h; IV drug abuse; immunosuppression (HIV, transplant patients); UTI, skin infection, diabetes mellitus; alcohol abuse; vertebral tenderness with signs of underlying infection; direct implantation (nail, acupuncture); Hx of spine surgery^{208,209}</p> <ul style="list-style-type: none"> • Suspected failed surgical fusion • Progressive or painful structural deformity (<1% of all LBP): scoliosis, kyphoscoliosis, transitional segments²¹⁰ & LDH • Abnormal laboratory examination and positive S&S: ESR >50 mm/h; Rh factor; HLAB27; reduced hematocrit; elevated white blood count; ALP; PSA; serum calcium for example.^{211,212} 	<p>c. NM</p>	<p>than conventional radiography, CT is not deemed useful to detect active joint inflammation. NM has had mixed results for both sensitivity & specificity.⁹⁹</p> <p>c. NM: commonly used for detection of possible destructive osseous lesions, skeletal metastatic diseases (known primary tumor-skeletal survey), spinal infection, stress fracture of the pars interarticularis and acute inflammatory sacroiliitis. Bone scan, such as gallium-67 may detect spondylodiscitis as early as 48 h after inoculation.^{6,209,215}</p> <p>Abnormal laboratory examination: R/O marrow-based malignancies, including metastasis & primary marrow-based pathology (e.g. myeloma); serum gammopathy and possible infection & arthritis</p>
<ul style="list-style-type: none"> • Suspected cauda equina syndrome (CES) (0.04% of all LBP) <p>The classic syndrome includes LBP, bilateral or unilateral sciatica, saddle anesthesia, motor weakness of the lower extremities that may progress to paraplegia, urinary retention, or bowel and bladder incontinence.</p> <p>The most consistent finding of a CES is urinary retention (sensitivity of 0.90, specificity of 0.95, likelihood ratio of 18). Unilateral or bilateral sciatica, sensory and motor deficits, and abnormal SLR are common, all have a sensitivity of >0.80. The most common sensory deficit occurs over the buttocks, posterior-superior thighs, and perineal regions (saddle anesthesia), with a sensitivity of ≈ 0.75. The saddle anesthesia is characterized by altered sensation in the distribution of sacral dermatomes (S2-S5). Anal sphincter tone is diminished in 60%-80% of cases. Anal wink reflex is absent in the majority of cases.^{44,99,112,167,182,215-219}</p>	<p>Emergency referral without imaging</p> <p>If clinical findings are equivocal, medical referral and specialized imaging recommended [B]</p> <p>Special investigations [C] (see previous page for details)</p>	<p>It is vital to recognize the S&S of CES as this is a surgical emergency.</p> <p>Although several etiologies can cause CES (spinal tumors, epidural abscess, etc), it usually arises from a massive midline posterior disc herniation (represents 1%-2% of all operated LDH). Narrowing of the spinal canal (congenital or degenerative stenosis) is a likely predisposition to CES. To facilitate the early detection of rare but serious complications associated with disc herniations, inquiry should also be made regularly during visits with regard to saddle anesthesia or any changes in bladder or bowel functions which may suggest the development of cauda equina syndrome.</p> <p>The risk of SMT causing a clinically worsened disc herniation or CES in a patient presenting with an LDH has been estimated to be <1 in 3.7 million manipulations, but others have proposed figures as low as 1 in 100 million SMT. The association in some reported cases has been described as temporal rather than causal.^{44,149,161,182,220-223}</p>
<ul style="list-style-type: none"> • Suspected abdominal aortic aneurysm (AAA) <p>Uncommon before age 50, AAA prevalence increases with age (3% in <65 yoa & $>8\%$ in >65). About 25% mimic musculoskeletal LBP pain and 75% may be silent (incidental findings on physical examination and/or radiography) until rupture or size draws attention.</p> <p>Early S&S may include abdominal pain, back-ache, and feeling of fullness or abdominal pulsation. Patient characteristics which may raise clinical suspicion of AAA include older age,</p>	<p>Referral or special investigations [B]</p> <p>In nondissecting AAAs, medical referral and US recommended even if conventional radiographs are negative (calcification, the most reliable radiological sign, is seen in only 50% of AAA)^{229,235}</p>	<p>Recommendation for AAAs (symptomatic or not): Rate of growth of small AAAs is relatively predictable.²³⁶ The sensitivity of ultrasound (US) scanning for AAA is 95%, and the specificity approaches 100%; the examination is safe and reliable.²³³</p> <p>Management^{42,226,227,230-233,236-240,}</p> <ul style="list-style-type: none"> • Ultrasound screening: Recommended in males 65 to 75 yoa (up to 80 in some studies); in females 60 to 85 yoa with cardiovascular risk factors; and in males and females >50 with a first-degree relative with AAA.

(continued on next page)

Table 3 (continued)

Patient presentation	Recommendations	Comments
<p>obese males, smokers or with a significant Hx of smoking, Hx of myocardial infarction, claudication, having a first relative with AAA, and presence of hypertension.²²⁴⁻²²⁷</p> <p>Physical exam including abdominal inspection, auscultation, percussion, & palpation; and peripheral vascular exam. However, clinical exam cannot be relied upon to exclude AAA, especially in smaller AAAs & if abdominal girth >100 cm.^{226,228}</p> <p>Atherosclerosis accounts for 90%-95% of AAA, with the remainder predominantly the inflammatory type. AAAs may be defined as a permanent focal dilation of 1.5 times its normal diameter or an expansion of the aorta measuring >3.8 cm on the lateral view measured between the most distant calcified borders, usually between the renal artery (L2) and the iliac bifurcation (L4). Common iliac arteries are often involved.²²⁹</p> <p>Risk factors^{225,230-234}:</p> <ol style="list-style-type: none"> Abdominal aortic aneurysms (AAA): Male sex, cigarette smoking, a first-degree relative with AAA, hypercholesterolemia, and connective-tissue diseases (eg, Marfan, Ehlers-Danlos). Aneurysm expansion risk factors: Advanced age (>70 y), cardiac/renal transplant, previous stroke, severe cardiac disease, tobacco use. Risk factors for aneurysm rupture: Increases with age (age >60) and aneurysm size (diameter >5.5cm); current smoking and high mean blood pressure. 		<ul style="list-style-type: none"> • Ultrasound monitoring: Yearly screening recommended if AAA between 3.0 and 4.4 cm (possibly every 6 mo if AAA is between 4.0 and 4.4 cm). • Referral to a vascular specialist: Recommended if AAA >4.4 cm (measures based on US or CT). <ul style="list-style-type: none"> ○ Annual rupture rate of small AAA (<4.5 cm) is 1%-3%. • Surgical intervention: Recommended if AAA ≥5.5 cm or growth of >0.6 cm to 0.8 cm/y. <ul style="list-style-type: none"> ○ Annual rupture rate and death from AAA >5.5 cm is 16%, compared with perioperative mortality of 2%-6% (or lower) for open repair.²³¹ • SMT: Relative risk of high velocity, low amplitude (HVLA) lumbar spine manipulation currently undetermined.¹⁹¹ However, considering current recommendations on AAA management, it appears reasonable to propose the following with regard to SMT: <ul style="list-style-type: none"> ○ Small AAA (3.8-4.0 cm): may be regarded as <i>relative</i> contraindication for SMT. Consider expansion risk factors and the need for yearly ultrasound monitoring. ○ AAA between 4.0 and 4.5 cm: consider the need for ultrasound monitoring every 6 mo. May be regarded as <i>absolute</i> contraindication for SMT in the presence of expansion/rupture risk factors. ○ Larger AAAs (≥ 4.5 cm) may be regarded as <i>absolute</i> contraindication for SMT. Refer patient to a vascular specialist. <p>As many as 1 in 3 AAAs eventually rupture if left untreated. Most AAA deaths occur in men 65 yoa and older. Rupture of AAA is associated with a risk of death approaching 80%.^{226,233,236}</p> <p>AAAs confer a risk of spontaneous rupture and death, which is related to the diameter of the aneurysm: a risk of rupture of 0.5%, 1%, 11%, and 26% per year has been associated with a diameter of <4 cm, 4.0-4.9 cm, 5.0-5.9 cm, 6.0-6.9 cm, respectively.²³⁰</p>
<ul style="list-style-type: none"> • Truncal symptoms attributed to presence or worsening of aortic aneurysms including dissection/rupture/occlusion or traumatic aortic injury): Cardiovascular shock and/or syncope, severe, tearing/ripping midabdominal sensation, back, groin, or testicular pain; pressure upon lumbar spine causing excruciating boring pain in the abdomen or back; hypotension; absence of distal lower limb pulses.^{224-226,241,242} <p>At least 25% of AAAs >5 cm will rupture <i>within 5 y</i>, and those <5 cm have a rupture rate of about 5%.²²⁹</p>	<p>Emergency referral without imaging [GPP]</p> <p><i>It is vital to recognize the S&S of dissecting AAA as this is a surgical emergency.</i></p>	<p>Evaluation of acute aortic conditions including dissection/rupture/occlusion or traumatic aortic injury^{224,243}:</p> <ul style="list-style-type: none"> • Because of its speed and proximity to emergency department, CT, helical CT, and CTA are imaging of choice to determine the presence and size of the aneurysms, as well as to R/O a ruptured AAA in a clinically stable patient. • MRI with MRA also plays an important role, particularly when patient is unable to receive intravenous contrast medium. • Conventional angiography is used as a secondary diagnostic tool to clarify equivocal findings. • US is helpful when CT is not readily available or patient is too unstable to undergo MRI

Table 4. *Nontraumatic thoracic spine disorders*

Patient presentation	Recommendations	Comments
<p>Adult patient with uncomplicated* acute (<4 wk duration) thoracic spine pain^{6,38,42,63}</p> <p>AND</p> <p>Adult patient with uncomplicated* subacute (4-12 wk duration) or persistent (>12 wk duration) thoracic spine pain and no previous treatment trial.^{6,38,42,63}</p> <p>* <i>Uncomplicated definition:</i> nontraumatic thoracic pain without neurologic deficits or indicators of potentially serious pathologies—(see red flag list for details).</p> <p>History and physical examination should exclude causes of thoracic referred pain from nonmusculoskeletal origin (see nontraumatic chest wall pain).</p>	<p>Radiographs not routinely indicated [B]</p> <p>Special investigations not indicated [B]</p>	<p>Degenerative changes are generally present in the middle-age adult. Radiographs are rarely useful in the absence of neurologic signs or red flags. However, consider osteoporotic vertebral collapse or other forms of destructive bone processes in the presence of acute sudden onset pain, especially in the elderly.^{6,42,63}</p>
<p>Adult patient: reevaluation in the absence of expected treatment response or worsening after 4 wk.^{38,42}</p> <p>Should patient fail to improve as expected or marginally improve within 4 wk of initial evaluation, the clinician must review history and physical findings and request appropriate diagnostic imaging studies.</p>	<p>Radiographs indicated [B] AP, lateral thoracic spine views</p> <p>Additional views: Swimmer’s view</p> <p>Special investigations [C] • <i>Consider NM if conventional radiographs are unremarkable.</i></p>	<p>Symptoms located in cervicothoracic junction or if this area is not well visualized on lateral view.</p> <p>Comanagement or specialist referral recommended</p> <p>• MRI or CT scan should be considered:</p> <ol style="list-style-type: none"> 1. In suspected pathology as seen on conventional radiography 2. After failed conservative therapy (4 wk), 3. If neurologic status is deteriorating (progressive deficit, disabling leg pain); <p>Negative findings (MRI/CT/NM) may be reassuring to patients.</p>
<p>Adult patient with nontraumatic chest wall pain. History and physical exam first need to rule out life-threatening conditions including pathologies of the heart, lungs, and large vessels.</p>	<p>Emergency referral without imaging in life-threatening conditions [GPP]</p> <p>Special investigations [C]</p>	<p>Nonmusculoskeletal causes of chest wall pain include disorders of the myocardium and pericardium (infarction, angina pectoris, myocarditis, rheumatic carditis, & pericarditis) of the pleura and lung (pleurisy, empyema, pneumothorax, pulmonary infarction, pneumonia, and neoplasm), disorders of the central structures (mediastinitis, sternal thyroiditis, Hodgkin’s disease, oesophagitis, hiatal hernia, aortic aneurysms with or without dissection), intra-abdominal (hepatobiliary), retroperitoneal disorders (pyelonephritis and tumors), and skin disorders such as herpes zoster.²⁴⁴</p> <p>• CT and MRI²⁴⁵</p>
<p>Musculoskeletal chest wall pain Musculoskeletal causes of chest wall pain are a diagnosis of exclusion. Causes are generally identified upon physical examination by reproducing patient complaint (observe for skin lesions, limited and painful movements, positive orthopedic tests such as Schepelmann’s sign, and tenderness on palpation).²³</p>	<p>Radiographs not routinely indicated [D]</p>	<p>Musculoskeletal causes include disorders of the rib, cartilage, and sternum (spondyloarthropathy, costovertebral syndrome, osteochondritis, Tietze’s syndrome, slipping rib syndrome); muscles (intercostals, diaphragm, serratus anterior, and pectoral); and intercostal neuralgia.^{244,246-248}</p>
<p>Adult patient with complicated (i.e., “red flag”) thoracic pain & indicators of contraindication to SMT (relative/absolute):</p> <p>Presence of any of the following indicators should alert the clinician to possible underlying pathology.^{8,38,44,63,112,114,167,185,197,203}</p> <p>N.B. Presence of a red flag alone may not necessarily indicate the need for radiography.</p> <p>• Patient <age 20 and >age 50, particularly with S&S suggesting systemic disease**</p>	<p>Radiographs indicated [B] AP, lateral thoracic spine views.</p> <p>Main purpose of thoracic spine radiograph is to exclude back pain caused by inflammatory spondyloarthropathy, fracture, malignancy, and infection.</p>	<p>Normal radiographs may falsely reassure clinicians as important causes of back pain are not easily identified with conventional radiographs (ie, not sensitive).⁷⁷</p>

(continued on next page)

Table 4 (continued)

Patient presentation	Recommendations	Comments
<ul style="list-style-type: none"> No response to care after 4 wk. Significant activity restriction >4 wk Non mechanical pain (unrelenting pain at rest, constant or progressive S&S) Persistent localized pain (>4 wk) Progressive or painful structural deformity: scoliosis, kyphoscoliosis²¹⁰ Symptoms associated with neurologic signs in the lower extremities 	<p>Additional views: Spot views.</p> <p>In suspected inflammatory spondyloarthropathy, consider spot angulated AP or PA lumbosacral or oblique SI views.</p> <p>Special investigations [B]</p> <p>* Even if conventional radiographs are negative: MRI is generally the preferred investigation.</p> <p>CT may be needed for bony details.</p> <p>NM commonly used for detection of possible destructive osseous lesions, metastatic diseases, infection, or inflammatory processes^{6,215}</p>	<p>Additional views may be considered in the presence of inflammatory spondyloarthropathy symptoms, although SI joints usually adequately demonstrated on AP/PA lumbar spine.</p> <p>Advanced imaging and specialist referral recommended even if conventional radiographs are unremarkable:</p> <ol style="list-style-type: none"> In the presence of a potentially serious pathology as suggested by the history, examination, and/or radiographs; In the absence of clinical improvement after 4 wk of therapy; Function does not improve or deteriorates; If neurologic status is deteriorating (progressive deficit, disabling pain); Painful or progressive structural deformity; unstable segment (scoliosis, kyphoscoliosis, pathological process); Patient has persisting S&S; Complication from treatment (possible fracture, new/progressive neurologic deficit, considerable pain or disability, etc)
<ul style="list-style-type: none"> Suspected inflammatory spondyloarthropathy: significant morning stiffness (>1 h); pain duration ≥ 3 mo; persisting motion restriction; gradual onset \leq age 40; peripheral joint involvement; UTI, urethral discharge, iritis, skin rash. Suspected neoplasia: considerable pain starting \geq age 50; Hx of cancer/carcinoma (in the last 15 y) and/or unexplained weight loss (>10 lb over 4 wk); or failure of conservative care (4 wk); systemically unwell; lymphadenopathy; dermatopathy (melanoma).^{6,203} Suspected infection: documented fever ($>38.3^{\circ}\text{C}/101^{\circ}\text{F}$) for >3 wk and no established diagnosis despite appropriate investigation for 1 week; elevated ESR >20 mm/h; IV drug abuse; immunosuppression (HIV, transplant patients); UTI, skin infection, diabetes mellitus; alcohol abuse; vertebral tenderness with underlying infection; direct implantation (nail, acupuncture); Hx of spine surgery.^{208,209} 	<ol style="list-style-type: none"> MRI or CT NM MRI especially for marrow-based lesion and neurologic disorders. MRI (Gadolinium) may increase specificity and detect spondylodiscitis as early as 48 h after inoculation²⁰⁹ 	<ol style="list-style-type: none"> MRI or CT: Suspected primary bone tumor NM: Known primary tumor, skeletal metastases, inflammatory arthritis, or spinal infection MRI: More accurate than radiography and bone scan (sensitivity of 0.96 & specificity of 0.92) for suspected osteomyelitis, spondylodiscitis, septic disc, paraspinal abscess, epidural abscess; in suspected myelopathy/long tract signs, spinal cord tumors, inflammation, infection, infarction, etc
<ul style="list-style-type: none"> Suspected failed surgical fusion 		
<ul style="list-style-type: none"> Abnormal laboratory examination and positive S&S: ESR >50 mm/h; RA factor; HLA-B27; BCB; ALP; PSA; serum calcium, for example.^{211,212} 		<p>Abnormal laboratory examination: R/O marrow-based malignancies, including metastasis & primary marrow-based pathology (e.g. myeloma); serum gammopathy and possible infection & arthritis</p>
<ul style="list-style-type: none"> Suspected acute thoracic aortic aneurysms dissection/rupture/occlusion or traumatic aortic injury: Severe, tearing/ripping chest sensation, back pain; hypotension; absent distal pulse. High index of suspicion in connective tissue disorders and diseases with genetic predisposition for ascending aortic aneurysms such as Marfan or Ehlers-Danlos syndrome.^{249,250} 	<p>Emergency referral without imaging [GPP]</p>	
<ul style="list-style-type: none"> In recent significant trauma (any age) 	<p>See Blunt Trauma/Acute Injuries section.</p>	
<ul style="list-style-type: none"> Suspected compression fracture ($\pm 4\%$ of all back pain)⁶³: specific pathologies and severe trauma account for only 3% and 14%, respectively, of all clinically evident vertebral fractures. The majority in North America are related to osteoporosis. S&S: Severe onset of pain (with or without appearance of spinal deformity) after trauma in older patients. Patients with thoracic or lumbar spine osteoporotic fractures report pain mainly in the lumbosacro-gluteal area. Look for Hx of repetitive stress of sufficient severity or Hx of high-risk osteoporosis.^{107,251} Prevalence of asymptomatic vertebral fracture significantly increases with age and glucocorticoid use in postmenopausal women.²⁵² 	<p>Radiographs indicated [B]</p> <p>AP, lateral thoracic</p> <p>Additional views [D]</p> <p>Supine cross-table lateral view in suspected osteoporotic vertebral pseudoarthrosis (VPA)²⁷⁵</p>	<p>Although compression fractures can be visualized on conventional radiography, a clear definition for identification of these vertebral "deformities" is still lacking (absence of a true gold standard).²⁵³ Furthermore, radiography may be insensitive for revealing osteoporotic fractures^{254,276,277} and there is no consistent relationship between back pain and changes in vertebral shape on radiographs.²⁵³</p> <p>Vertebral compression fractures represent a serious public health problem by virtue of the associated disability and cost. Physical function, such as cooking meals, shopping, bending, lifting, and descending stairs, self-esteem, body image, and mood are negatively affected by vertebral fractures.²⁵³</p>

Table 4 (continued)

Patient presentation	Recommendations	Comments
<ul style="list-style-type: none"> Only about one third of all affected women in the community have the severe vertebral deformities most likely to produce chronic symptoms including height loss, kyphosis and postural changes, and persistent pain interfering with daily activities.²⁵³ 		<p>Age-adjusted incidence rates for clinically evident vertebral fractures in women are twice those in men, partly due to their greater liability to falls.²¹⁷ Lifetime risk of clinically evident osteoporotic fracture among postmenopausal North American white women >age 50 has been estimated at 40% (vertebrae 15.5%; wrist 16%; hip 17.5%)^{254,278} and at 22% in men.²⁷⁹ The incidence of osteoporotic fractures is increasing and the spectrum of fractures is changing. Fractures of the proximal femur, proximal humerus, pelvic, and many fractures around the knee should also be considered osteoporotic fractures.²⁸⁰</p>
<p>Risk factors for additional vertebral fractures²⁵³:</p> <ol style="list-style-type: none"> History of a previous fracture (4-fold among postmenopausal white women); Greater age; Lower femoral neck bone mass density (BMD); Shorter height. 		<p>Vertebral pseudarthrosis, a rare posttraumatic vertebral delayed collapse resulting in bone ischemia, may cause vertebral instability: look for an intravertebral cleft containing gas. These patients tend to have ongoing back pain despite conservative care, severe pain during motion.²⁷⁵</p>
	<p>Special investigations [D]</p>	<p>In suspected fracture:</p> <ul style="list-style-type: none"> Consider MRI or CT if initial radiographs are positive for fracture, equivocal, in the presence of complex lesions, for suspected ligamentous instability or neural injuries. Small fractures can also go undetected on CT scans.²⁸¹ NM or MRI may be used to determine if fracture is recent or not, as it may impact on treatment. (NM has a 72-hour delay in elderly patients.)⁶³
<p>Suspected osteoporosis</p> <p>Osteoporosis predisposing risk factors^{42,206,252,254-265}:</p> <p>Include nonmodifiable (age, sex, and period of amenorrhea as in anorexia nervosa and female athlete) and modifiable factors (dietary calcium intake, low BMI, smoking, inactivity, parental Hx of fracture, and high alcohol intake). See Appendix C(4) for a list of other important risk factors.</p>	<p>Radiographs are <i>unreliable</i> for assessment of bone mass changes before at least a 30%-50% loss.²²⁹ [B]</p>	<p>There is a concern that spinal manipulation (HVLA)^{282,283} and spinal mobilization^{281,284} could cause fractures in individuals with osteoporosis. Spinal manipulation is generally regarded as a relative contraindication in osteoporotic patients.^{239,285-287} However, in vitro studies and some case reports suggest forces generated in some manual techniques may be lower than the load required for vertebral failure (fracture) in this population.^{273,281} A number of control strategies are available that allow treatment to be applied safely and effectively.²⁸⁸</p>
<p>Osteoporosis clinical decision rules:</p> <p>Several highly sensitive clinical decision rules based on weighted indices and major osteoporosis risk factors have been developed to identify women with low bone mineral density in need of DXA.^{260,266-271}</p>	<p>Special investigations [B]</p> <p>If clinical decision rules are positive</p>	<p>When DXA results are unavailable, use clinical decision rules to estimate the presence of osteoporosis.</p> <p>See Appendix C(4) for details.</p>
<ol style="list-style-type: none"> In healthy peri- and early postmenopausal women (45-64 yoa), consider using the OST score (Osteoporosis Self Assessment Tool). OST score considers only 2 variables: (weight in kg - age)/5. <i>The cut-off for a positive test is <2, indicating this woman should be referred for DXA.</i> In higher-risk patients, use either the ORAI (Osteoporosis Risk Assessment Instrument), the more complex calculation of SCORE (Simple Calculation of Osteoporosis by Estimation) or the AMMED as all 3 were found to be highly sensitive clinical tools. See Appendix C(4) for details. <ol style="list-style-type: none"> ORAI considers age, weight and estrogen use; SCORE considers 6 variables (race, presence of RA, history of fracture, age, estrogen, weight [in pounds]); AMMED considers 5 variables (age, years after menopause, age at menarche, BMI). 		<p>In suspected osteoporosis</p> <ul style="list-style-type: none"> Bone densitometry or dual-energy x-ray absorptiometry (DXA) recommended in the presence of osteoporotic fractures or with positive osteoporosis clinical decision rules (elevated risk of osteoporosis). DXA is considered the gold standard for bone mass evaluation in the diagnosis and treatment of osteoporosis (DXA >2.5 SD). In postmenopausal females, each SD decrease in lumbar spine BMD increases the risk of any fractures by 1.5-fold and of new vertebral fracture by 2.3-fold. Lateral DXA also accurate for thoracic spine evaluation.^{253,284} <p>Of interest:</p> <ul style="list-style-type: none"> Advanced atherosclerosis of the abdominal aorta is associated with lower bone mass density, accelerated bone loss, and increase risk of incident fractures at the proximal femur. Findings in the lumbar spine have been inconsistent.²⁸⁹

(continued on next page)

Table 4 (continued)

Patient presentation	Recommendations	Comments
<p>Physical examination and osteoporosis:</p> <p>When likelihood of significant osteoporosis is low according to clinical decision rules, examination may include asking patient to rise from a chair and to bend forward at the trunk as vertebral fracture risk associated with loading activities increases risk by about 8- and 10-fold respectively compared with upright standing.²⁷² It should be noted, however, that vertebral tolerance to PA thoracic compressive forces is much less than axial loading.²⁷³ Simple physical tests that could assist in determining ability to tolerate thoracic spine SMT would be useful.</p> <p>Quality of life among adults with vertebral osteoporosis may be measured using a self-administered questionnaire such as the one from the European Foundation for Osteoporosis (QUALEFFO).²⁷⁴</p>		<p>Physical examination: Vertebral Fracture Index (rising from a chair & bending forward) has a lower predictive value with a lower sensitivity and specificity than axial bone mineral density measured by DXA in assessing vertebral fracture risk.²⁷²</p>
<p>Adult patient with nonpainful and nonprogressive scoliosis</p> <p>The prevalence rate in healthy adults is >30% and may be as high as 68% in the elderly.²⁹⁰ Untreated adults with late-onset idiopathic scoliosis (LIS) are productive and functional at a high level at 50-year follow-up. Untreated LIS causes little physical impairment other than back pain (most only have minimal or moderate back pain) and cosmetic concerns.²⁹¹ Patients with nonpainful and nonprogressive scoliosis are unlikely to benefit from repeat evaluation.</p>	<p>Radiographs not routinely indicated [C]</p>	<p>In a skeletally mature patient, scoliosis is defined as a spinal deformity with a Cobb angle of >10° in the coronal plane.²⁹²</p>
<p>Adult patient with painful or progressive scoliosis</p> <p>Some patients with known scoliosis may present with significant disability. Back pain is the most common clinical problem presenting as a multiform mosaic of symptoms. Constant & nonspecific back pain has a poor prognosis. Other S&S: Radicular pain & claudication when standing or walking (from nerve traction or compression), neurologic deficit may include sphincter dysfunction. Curve progression needs monitoring.²⁹²</p> <p><i>Clinical evaluation:</i> Look for spinal deformity, uneven shoulder heights, limb length inequality, pelvic obliquity, hip pathology, and claudication. Neurologic exam includes nerve root tension signs, motor power, sensations, deep tendon reflexes, and pathological reflexes.²⁹³⁻²⁹⁵</p> <ol style="list-style-type: none"> Adam's forward bend test is sensitive to detect trunk asymmetry Maximal lateral flexion into convexity to evaluate curve functional aspect. Scoliometer (a trunk angle of 7° indicates a structural curve >20°.) <p><i>Spinal deformity has an important impact on the general health status of adults as measured by the Medical Outcomes Study Short-form 36 (SF-36) and the modified Scoliosis research Society Outcome instrument (SRS-22). The SRS-22 is a disease-specific instrument assessing pain, self-image, function, mental health, and satisfaction. It is a valid, reliable, and reproducible outcomes instrument for adult deformity.</i>^{296,297}</p>	<p>Radiographs indicated [B]</p> <p>Erect sectional radiographs (better detail) or standing full-length PA (14 × 36 in) and lateral sectionals</p> <p>PA films* significantly reduce breast and thyroid dose. Effective doses to the digestive and respiratory systems are comparable, but are higher in the bone marrow compared to AP views.^{299,300}</p> <p>Full spine radiographs not recommended for patients with an AP measurement >28 cm or for older patients due to poor film quality. Consider using sectional radiographic views instead.²²⁹</p> <p>Additional views:</p> <ol style="list-style-type: none"> Right and left lateral bending 	<p>Full spine radiograph for scoliosis to determine other areas of implication</p> <ol style="list-style-type: none"> Cobb angle Nash-Moe method (pedicle rotation) <p>NB. Consider ± 5° measurement error</p> <p>Clinically relevant radiographic parameters in adult scoliosis including lateral vertebral listhesis, L3 and L4 endplate obliquity angles, increased lumbar lordosis and thoracolumbar kyphosis significantly correlated with pain,³⁰¹ sagittal imbalance & significant coronal imbalance of >4 cm,³⁰² lateral spondylolisthesis (lateral translation & vertebral rotation) may cause radicular pain syndrome.³⁰³</p> <p>Types of lumbar scoliosis may be classified according to the lordosis and L3 obliquity. Self-reported pain & disability may increase with increasing scoliosis type (from I to III).³⁰⁴</p> <ol style="list-style-type: none"> Most commonly used to determine fusion levels. May help differentiate between structural and nonstructural curves and help assess primary from secondary scoliotic curves. <p>Adult degenerative scoliosis: pathomechanism includes a viscous cycle of asymmetric degeneration (disc & facet joints) leading to asymmetric loading, leading to asymmetric deformity which in turn triggers asymmetric degeneration and loading and enhancing curve progression.²⁹²</p>

Table 4 (continued)

Patient presentation	Recommendations	Comments
<p>There are many causes of spinal deformity in the adult. They may be grouped as follows:</p> <p>(1) Primary degenerative scoliosis (mostly lumbar or thoracolumbar—triggered by disc degeneration).²⁹⁸</p> <p>(2) Progressive idiopathic adolescent scoliosis of the thoracic, thoracolumbar, and/or lumbar spine.</p> <p>(3) Secondary degenerative scoliosis</p> <p>(3a) Scoliosis after idiopathic, neuromuscular, or congenital scoliosis, or occurring in the context of pelvis obliquity, leg length discrepancy, hip pathology, or a lumbosacral transitional anomaly.</p> <p>(3b) Deformity secondary to metabolic bone disease (mostly osteoporosis) combined with asymmetric arthritis and/or vertebral fractures.²⁹²</p>	<p>Follow-up evaluation [C]</p> <p>Repeat radiographs [B]</p> <p>Special investigations [C]</p>	<p>Follow-up evaluation dictated by clinical progression. Long-term follow-up of <i>adolescent idiopathic scoliosis</i> has revealed little curve progression even 20 y after bracing (7.9°) or surgical treatment (3.5°). Degenerative disc changes are more common in both patient groups compared to a control group.³⁰⁵</p> <p>Repeat radiographs, advanced imaging, and specialist referral recommended²⁹²</p> <ol style="list-style-type: none"> 1. In the absence of clinical improvement; after 4 to 6 wk of therapy; 2. If function does not improve or deteriorates; 3. In the presence of persisting S&S or considerable pain; 4. If patient neurologic status is deteriorating (progressive deficit, disabling leg pain); 5. With painful or progressive structural deformity (scoliosis, kyphoscoliosis); 6. With suspected segmental instability (this is common in adult scoliosis and should be considered with all manual therapy intervention). 7. With suspected pathological process; 8. With new or progressive neurologic deficit including claudication, significant radiculopathy, or suspected syrinx; 9. To plan surgical intervention. <ul style="list-style-type: none"> • Spiral CT useful in rapid reconstruction of the spine. • MRI to evaluate the spinal cord and nerves. • Sequential discograms, facet blocks, epidural blocks

Table 5. *Nontraumatic cervical spine disorders*

Patient presentation	Comments
<p>General background information</p> <p>Most neck pain is not the result of serious disease. Factors associated with neck pain include³⁰⁶:</p> <ul style="list-style-type: none"> • Headache • Low back pain • Poor psychological health • History of whiplash injury • Poorer general or self-rated health • Possibly cigarette smoking <p>• In patients presenting with head and/or neck pain, the physical examination should flow as a natural extension of the information obtained from the chief complaint, history, and system review. (see Honet and Ellenberg³⁰⁷ for a review of history & physical examination for neck pain).</p> <p>• In complaints of neck and arm pain, peripheral (weakness, dermatomal pattern, and loss of deep tendon reflexes) and central nervous system exam (clonus, Babinski, and pathological reflexes) should be an integral part of initial examination. Findings of long tract signs mandate further evaluation.</p>	<p>Standards of care dictate that careful history and physical examination be done routinely for all patients on the initial visit.</p> <ul style="list-style-type: none"> • Self-reported patient assessment to evaluate perceived pain, function, disability, and psychosocial status are recommended. • The validity and utility of the clinical assessment (history and examination) in patients with neck pain with out radiculopathy or red flags needs to be confirmed. • Diagnostic tests such as X-rays, CT or MRI scans are only required in a minority of cases when a thorough physical examination and patient history indicate further investigation is needed. Routine imaging is unlikely to increase understanding of causation. <p>Duration of complaint:</p> <ul style="list-style-type: none"> • Acute neck pain: <4 wks' duration • Subacute neck pain: 4-12 wks' duration • Persistent/chronic neck pain: >12 wks' duration <p>Neck pain affects nearly two third (66.7%) of the adult population over their lifetime representing a significant and costly health problem in terms of care, suffering, and work absenteeism.³¹⁵⁻³¹⁷ Contrary to prior belief, most individuals with neck pain do not experience complete resolution of their symptoms and disability.</p> <p>The age-standardized 6-month point prevalence is around 22% with nearly 40% having low-intensity and low-disability neck pain, 10% having high-intensity and low-disability neck pain and 4.6% (95% CI, 3.3-5.8) having disabling pain.³¹⁶</p>

(continued on next page)

Table 5 (continued)

Patient presentation		Comments
<ul style="list-style-type: none"> In headache complaints, vital signs (to R/O severe hypertension or fever) and testing of the cranial nerves (to R/O vascular events, space occupying lesions, etc) should be an integral part of initial examination. Significant positive findings mandate further evaluation. In suspected connective tissue disorders, examination should also include inspection of the integument (skin elasticity, discoloration), the back (scoliosis), the thorax (deformity, abnormal respiratory pattern), and extremities (disproportion), auscultation of the heart (valvular defect) and abdominal aorta (bruit), abdominal palpation (organomegaly in fibromuscular dysplasia-kidney, aorta), and musculoskeletal examination including peripheral joint laxity using the 1998 Beighton criteria and examination of chronic joint and limb complaints.³⁰⁸ <p>Auscultation of the neck and the use of common premanipulation vascular screening tests (eg, Houles, George, etc) are of no diagnostic value. Functional vascular tests may themselves put patient at risk of a serious complication.³⁰⁸⁻³¹⁴</p>		<p>The age and sex annual incidence of neck pain is 14.6% (95% CI, 11.3-17.9) and each year, 0.6% develop disabling neck pain. Annual course of neck pain is as follows: 36.3% resolve and 32.7% improve. Among neck pain sufferers, 33.3% report persistent problems, 9.9% get worse, and 22.8% recur.³¹⁷</p> <p>Between 22% and 25% of initial chiropractic visits are for neck-related complaints. Although most patients seeking chiropractic care experienced their symptoms for <3 wk (45%), over 20% had symptoms for >6 mo.⁴¹ In addition, 5%-6% of chiropractic patients consult for headache. Neck pain is slightly more common in females and increases with age, the mean age being in the early 40s.^{41,100} Not surprisingly, these figures parallel the most common age range seeking consultation for chiropractic spinal manipulative therapy (61% of females; mean age of 42 years old).</p>
<p>Adult patient with acute uncomplicated* neck pain (<4 wks' duration)</p> <p>* Definition of "uncomplicated": nontraumatic neck pain without underlying neurologic deficits or red flag (see red flags list for details).^{42,63,318}</p>	<p>Radiographs not routinely indicated [C]</p> <p>Special investigations not indicated [C]</p>	<p>The clinical value of conventional radiographs for nontraumatic acute neck pain <i>without</i> red flags or arm pain is not well established. Acute neck pain is generally due to conditions that cannot be seen on conventional radiographs.^{38,42,128} The clinical significance of the cervical spine curvature remains controversial.^{138,155,319-323}</p> <p>Consider cervical radiography, advanced imaging studies, and a second opinion in the absence of expected treatment response or worsening after 4 wk.^{38,42}</p>
<p>Adult patient with nontraumatic neck pain and radicular symptoms</p> <p>May be caused by HNP, spondylosis, and rarely by other pathological processes (tumor, abscess, etc).³¹⁸</p> <p>(A) Suspected acute cervical disc herniation (CDH)^{42,307,324-333}:</p> <ul style="list-style-type: none"> 35-55 yoa <p>(B) Suspected acute cervical spondylotic radicular syndrome/lateral canal stenosis (LCS)^{307,330,331}:</p> <ul style="list-style-type: none"> 45-54 yoa <p>Common S&S of acute cervicobrachial syndrome (A and B)^{38,307,330-333}:</p> <ul style="list-style-type: none"> Predominant arm & scapular pain with or without neurologic deficit: Specific weakness and/or hypoesthesia and/or hyporeflexia. Provocation tests (Spurling's, axial compression) & possible nerve root tension signs (upper limb tension test, median nerve & medial cord stretch test, shoulder depression, brachial-plexus compression, or arm pain relieved by cervical traction and/or Bakody). Possible increased intra-theal pressure (Valsalva/Dejerine triad) <p>N.B. Consider past imaging studies for evidence of disc pathology and/or moderate to advanced cervical spondylosis.</p>	<p>Radiographs indicated [D/consensus]</p> <p>APOM, AP lower cervical, neutral lateral</p> <p>Additional views:</p> <p>a. Oblique views</p>	<p>Disc lesions cannot be visualized on radiographs. IVF narrowing by facet & uncinat process hypertrophy are very often associated with radicular symptoms.³³⁴⁻³³⁶ Spinal stenosis (sagittal diameter <13 mm) may also be present, in which case special care should be taken if manual therapy is considered. Always examine the patient's lower extremities for upper motor neuron signs indicating cervical radiculomyelopathy (Hoffman sign, lower limb hyper-reflexia, spasticity, Babinski, and/or Clonus).^{307,335,336}</p> <p>Clinical diagnosis of CDH possible in only 50%-75% of cases of cervical radiculopathy as a wide variation of presentations exists even with only one nerve root involved (C6 more common than C7 which is more common than C8 level). Patients should be monitored as S&S may progress in the first 2-3 wk.^{38,337}</p> <p>When applied by properly trained and experienced practitioners, SMT is potentially a safe treatment option for patients with <i>nonprogressive</i> cervical radiculopathy secondary to CDH and LCS.³³⁸⁻³⁴⁰ Premanipulative position test may help determine whether patient can tolerate cervical manipulative procedure. Technique adaptation/mobilization recommended in patients with acute cervicobrachial pain.²³⁹</p> <p>a. MRI provides more useful information than conventional radiography oblique views. Oblique views may be considered in cervical radiculopathy, spondylotic myelopathy, and extensive degenerative, apophyseal joints, pedicles.^{229,341}</p>

Table 5 (continued)

Patient presentation		Comments
<p>When consistent with the history and other physical findings, a positive Spurling's, traction/neck distraction, and Valsalva's might be indicative of a cervical radiculopathy, whereas a negative upper limb tension test (contralateral neck rotation and extension of the arms and fingers) might be used to rule out radiculopathy.³³³</p>	<p>b. Swimmer's view</p>	<p>b. Consider swimmer's view if symptoms located in cervicothoracic (C/T) junction or if this area is not well visualized on lateral view.</p>
<p>See Appendix C(5) for a detailed list of clinical criteria for cervicobrachial syndrome.</p>	<p>Special investigations [C] Combined with radicular symptoms, specific findings on examination (manual provocation tests), and possibly needle EMG findings, MRI may be helpful in confirming the site and level of root compression.</p>	<p>Comanagement or specialist referral or MRI recommended even if conventional radiographs are unremarkable^{38,42,63}</p> <ol style="list-style-type: none"> 1. After failed conservative therapy (4 wk); 2. Major neurological deficits at onset, disabling radicular pain or progression of deficits; 3. For preoperative planning
<p>1. Adult patient with uncomplicated* subacute neck pain (4-12 wks' duration) with or without arm pain³¹⁸</p> <p>* Definition of "uncomplicated": nontraumatic neck pain without underlying neurologic deficits or red flags (see red flag list below for details).</p> <p style="text-align: center;">AND</p> <p>2. Adult patient with persistent neck pain (>12 wk) with or without arm pain^{63,318}</p> <p>Psychological distress/depressive mood, somatization, perceived disability, and passive coping behavior are important predictors of chronicity and poor outcome in persistent neck pain.^{202,342} An important association exists between a Hx of a WAD, pain intensity, & disability and co-morbidity (headache, LBP, digestive, & cardiovascular disorders).⁸⁶⁻⁸⁸ Complete history should therefore include a system review.</p>	<p>Radiographs not initially indicated [Consensus] APOM, AP lower cervical, neutral lateral NB. This recommendation was modified according to the recent findings of The Bone and Joint Decade 2000-2010 Task Force on Neck Pain and its Associated Disorders (see articles published in Spine 2008; 33(4S). A majority of Delphi panelists agreed with this change (92% of 50 respondents). Additional views: a. Oblique views b. Swimmer's view c. Flexion/extension</p>	<p>Vertebral degeneration and anterior slippage (anterolisthesis) increases with age. However, there appears to be no correlation between neck pain (with or without prior trauma) and the severity or the number of levels of disc, facet or uncovertebral joint degeneration, or degree of anterior slippage, but for the exception of a weak association between chronic neck pain (>6 month) & the severity of the degenerative disc disease (when present in the lower cervical spine).^{42,320,343-347}</p> <ol style="list-style-type: none"> a. Oblique views may be considered in cervical radiculopathy, spondylotic myelopathy, and extensive degenerative disease to evaluate IVFs, uncovertebral joints, apophyseal joints, pedicles.²¹⁸ However, MRI provides more useful information than conventional radiography oblique views. b. Consider swimmer's view if symptoms located in cervicothoracic (C/T) junction or if this area is not well visualized on lateral view. c. Flexion/extension views are indicated in suspected segmental instability: high-risk ligament laxity populations (Down's, RA and other inflammatory spondylo-arthropathies), failed surgical fusion, trauma and advanced degenerative disc disease.^{6,348} Patients should be supervised during procedure. An ADI interval >3 mm is abnormal.³⁴⁹
<p>Adult patient reevaluation in the absence of expected treatment response or worsening after 4 wk^{60,318}</p> <p>See Appendix C(6) for a list of psychological risk factors (yellow flags) and clinical indicators of significant anxiety or depression in whiplash patients.</p>	<p>Radiography indicated [C] APOM, AP lower cervical, neutral lateral</p> <p>Special investigations [B] (1. subacute neck pain, 2. persistent neck pain, and 3. reevaluation in absence of treatment response or worsening after 4 weeks)</p>	<p>Comanagement or specialist referral or MRI recommended even if conventional radiographs are unremarkable.^{38,42,63}</p> <ol style="list-style-type: none"> 1. If conventional radiography reveals suspected pathology; 2. After failed conservative therapy (4 wk); 3. If patient neurologic status is deteriorating (progressive deficit, disabling arm pain); 4. If clinical signs suggest subaxial cervical spine instability; 5. For preoperative planning; <p>NB. The frequency of all degenerative MRI findings increases linearly with age with disc degeneration being the most frequent ranging from 12% in asymptomatic males & females in their 20s to 89% in those over age 60. Posterior disc protrusion and even slight cord compression are not rare in asymptomatic subjects over age 40 with reduction of the cross-sectional area of the spinal cord not exceeding 16%.³⁵⁰⁻³⁵²</p>

(continued on next page)

Table 5 (continued)

Patient presentation		Comments
<p>Adult patient with complicated (ie, “red flag”) neck pain & indicators of contraindication to SMT</p> <p>Presence of the following indicator (s) should alert the clinician to possible underlying pathology^{38,42,63,308,318,353-363}:</p> <p>NB. Presence of a red flag alone may not necessarily indicate the need for radiography.</p> <ul style="list-style-type: none"> • Patient <age 20 and >age 50, particularly with S&S suggesting systemic disease** • No response to care after 4 wk. • Significant activity restriction >4 wk • Nonmechanical pain (unrelenting pain at rest, constant or progressive S&S) • Neck rigidity in the sagittal plain in the absence of trauma (discitis, infection, tumor, meningitis, etc); • Dysphasia; • Impaired consciousness; • Central nervous system S&S (cranial nerves, pathological reflexes, long tract signs); • High risk ligament laxity populations/suspected atlantoaxial instability (see details below); • Arm or leg pain with neck movements; • Suspected cervical myelopathy and radiculomyelopathy (see details below); • Sudden onset of acute and unusual neck pain and/or headache (typically occipital) with or without neurologic symptoms, suspected cervical artery dissection (VAD, CAD), TIA (VBI, carotid artery ischemia), stroke (see details below); • Hx of severe trauma (see Trauma section). 	<p>Radiographs indicated [B]</p> <p>APOM, AP lower cervical, neutral lateral</p> <p>Additional views:</p> <p>a. Flexion/extension</p> <p>b. Oblique views</p> <p>c. Pillar</p> <p>Special investigations^{6,38,42,63} [B]</p>	<p>Normal radiographs may falsely reassure clinicians as important causes of neck pain are not easily identified with conventional radiographs (i.e., not sensitive)</p> <p>APOM useful for trauma, inflammatory arthropathy, high-risk atlantoaxial instability populations, malignancy, known congenital anomaly.</p> <p>** Risk of having a serious pathology may be higher before age 20 & over age 55. Particular attention to indicators of possible underlying pathology should be given for patients in these age categories.</p> <p>a. Flexion/extension views indicated in suspected segmental instability: high-risk ligament laxity populations (Down’s, RA, and other inflammatory spondylo-arthropathies), failed surgical fusion, trauma, and advanced degenerative disc disease.^{6,348} Patients should be supervised during procedure. An ADI interval >3 mm is abnormal.³⁴⁹</p> <p>b. Oblique views may be considered in suspected fracture, malignancy, and other equivocal findings on standard AP and lateral radiographs (radiculopathy, spondylotic myelopathy, and extensive degenerative disease).</p> <p>c. Pillar views may be considered in severe trauma (CT usually done instead—see Trauma section)</p> <p>Advanced imaging and specialist referral recommended:</p> <ul style="list-style-type: none"> • MRI for suspected myelopathy (tumors, inflammation, infection, infarction, etc). <p>In cases of atlantoaxial instability, MRI (F/E) shows effects on cord compression when radiograph is positive or neurologic signs are present.</p>
<p>In addition, also consider general red flags (relative utility and cost-effective of screening patients for non traumatic neck pain for serious structural disease (“Red Flags”) need to validate) which may apply to the cervical spine</p> <ul style="list-style-type: none"> • Suspected neoplasia: considerable pain starting ≥age 50; Hx of cancer/carcinoma (in the last 15 y) or failure of conservative care (4 wk), lymphadenopathy; dermatopathy (melanoma). S&S of cervical spine tumors tend to be variable (slight pain, neck stiffness, scoliosis, presence of a mass and/or severe neurologic deficit). Night pain remains a classic indicator of nonmechanical neck pain and neoplasm. Other systemic symptoms may include unexplained weight loss (>10 lb over 4 wks) and systemically unwell.^{6,38,355,356} • Suspected infection (discitis, osteomyelitis, tuberculosis): documented fever (>38.3°C/101°F) for >3 wk and no established diagnosis despite appropriate investigation for 1 week; elevated ESR >20 mm/h; IV drug abuse; immunosuppression (HIV, transplant patients); UTI, skin infection, diabetes mellitus; alcohol abuse; vertebral tenderness with underlying infection; direct implantation (nail, acupuncture); Hx of spine surgery.^{208,209} • Suspected failed surgical fusion; • Progressive or painful structural deformity; 		

Table 5 (continued)

Patient presentation	Comments
<p>• Abnormal laboratory examination and positive S&S: ESR >50 mm/h; RA factor; HLA-B27; BCB; ALP; PSA; serum calcium, for example.^{211,212}</p> <p>Note: It may be reasonable to consider many of the lumbar spine red flags in complaints of neck pain. However, their level of evidence as such is currently unknown.¹⁵⁵</p>	<p>Abnormal laboratory examination: R/O marrow-based malignancies, including metastasis & primary marrow-based pathology (eg, myeloma); serum gammopathy and possible infection & arthritides</p>
<p>1. Suspected atlantoaxial instability (AAI): High-risk ligament laxity populations/possible atlantoaxial instability include^{6,63,229,357-363}:</p> <p>(a) Active inflammatory arthritis: known rheumatoid arthritis (RA), psoriatic arthropathy, chronic juvenile RA, and less commonly in AS (2% of cases). S&S of presumed inflammatory spondyloarthropathy include significant morning stiffness (>1 h); pain duration ≥3 mo; persisting motion restriction; gradual onset ≤age 40; peripheral joint involvement; UTI, urethral discharge, iritis, skin rash.³⁶⁴</p> <p>(b) Congenital disorders & hereditary connective tissues disorders such as dysplasia, os odontoideum, Klippel-Feil, Morquios, Down’s syndrome (20% are born without a transverse ligament), Ehlers-Danlos syndrome type III, Marfan syndromes.</p> <p>Screening for high-risk ligament laxity populations is important for manual therapists. Little is known about the value of commonly used manual tests. Although agreement and reliability between 2 examiners is difficult to achieve for general cervical spine mobility & intersegmental passive mobility,³⁶⁵ acceptable levels of predictive value, sensitivity, and specificity have been reported for the Sharp-Purser test.³²⁰ Cattrysse et al³⁶⁶ were not able to reproduce such results in a small group of children with Down’s syndrome, however. Preliminary data support the use of the upper cervical flexion test for those patients.</p>	<p>Radiographs indicated [B] APOM, AP lower cervical, neutral lateral</p> <p>Additional views [D] a. Flexion/extension laterals*⁶</p> <p>* Flexion/extension views indicated in suspected AAS. However, a single lateral cervical radiograph with the patient in supervised comfortable <i>flexion</i> should reveal any subluxation in patients with rheumatoid arthritis, Down’s syndrome, etc.⁶</p> <p>Special investigations [C] Monitoring, advanced imaging and specialist referral recommended^{63,318,348,369,371,372}:</p> <ol style="list-style-type: none"> 1. ADI >3 mm, vertical dislocation, lateral, posterior, or subaxial subluxations 2. Upward odontoid translocation (pseudobasilar invagination) 3. In the presence of neurologic S&S <p>◆ CT and MRI provide detailed images of the bone and spinal cord lesions.</p>
<p>2. Suspected cervical compressive myelopathy (CCM) and radiculo-myelopathy Causes of cord compression include trauma, tumors, infection, vascular disease, degenerative conditions, demyelinating disorders, spinal stenosis, & central cervical disc herniation.</p> <p><i>Cervical spondylotic myelopathy (CSM)</i>^{63,372-375}</p> <ol style="list-style-type: none"> 1. MC >50 y; 2. Variable neurologic deficit of upper (hyporeflexia) and lower extremities (hyperreflexia); axial skeleton sensory and motor dysfunction that skip the head & face; 3. Arm or leg pain with neck movements; 4. Long tract signs and pathological reflexes (Clonus, Babinski, Hoffman); Lhermitte’s sign (cervical flexion and extension producing electrical shocks down the arm and leg); 5. Subtle gait abnormality (loss of balance, spasticity, unsteadiness, loss of leg power, broad base, shuffle and disruption in smooth, rhythmic function); 	<p>Radiographs indicated [C] APOM, AP lower cervical, neutral lateral, and bilateral oblique views.</p> <p>Additional views: Swimmer’s view</p> <p>For cervical spondylotic myelopathy, patients should be treated conservatively if they have a spinal transverse area >70 mm², are of older age, and have normal central motor conduction time. Surgery is more suitable for patients with clinically worse status (function expressed as modified Japanese Orthopaedic Association score and slower walk) and a smaller transverse area of spinal cord.³⁷⁶</p> <p>Central spinal stenosis, acute cervical myelopathy, and/or radiculo-myelopathy is generally considered a contraindication for manual therapy although some patients with nonacute & nonprogressive S&S have safely received such intervention. Caution is advised in those patients as even minor trauma in patients with preexisting severe spondylosis has potentially serious consequences.^{318,336,339,340,377-385}</p> <p>Consider Swimmer’s view if symptoms are located in cervicothoracic junction or if this area is not well visualized on lateral view.</p> <p>a. A sagittal diameter <13 mm indicates spinal stenosis and is critical for the development of a cervical</p>

(continued on next page)

Table 5 (continued)

Patient presentation	Comments
<p>6. Bowel & bladder dysfunction (rare).</p> <p>Neurologic examination in patients with CSM is moderately accurate and reliable for determining the neurologic level of disease. Among the neurologic tests (deep tendon reflex, pinprick response, muscle weakness, and numbness in the hand only), patient-perceived location of numbness in the hands may be the most useful for establishing the affected level.³⁷⁴</p> <p style="text-align: center;">Special investigations [C]</p>	<p>radiculo-myelopathy, even in the presence of mild degenerative spondylosis (normal is 17-18 mm between C3-C7). Caution is recommended if AP diameter <11 mm and manual therapy is contemplated.³¹⁸</p> <p>b. A Torg' or Pavlov ratio (spinal canal/vertebral body) <0.8 indicates spinal stenosis.^{334,383,386}</p> <p>NB. It is important to recognize that conventional radiography may not be very reliable to determine central canal stenosis as there is some variation in the size of the spinal cord and stenosis may be due to nonosseous factors (ligamentum flava, cartilage hypertrophy etc).</p> <p>Refer patient for investigation and possible surgical intervention:</p> <ol style="list-style-type: none"> 1. After failed conservative therapy (4 wk); 2. If neurologic status is deteriorating (progressive deficit, disabling arm pain); 3. For preoperative planning: <ul style="list-style-type: none"> • MRI (CT-myelography if MRI not available) <p>Electrophysiologic testing such as somatosensory-evoked potentials (SSEP) may be useful.</p>
<p>3. Suspected cervical artery dissection (VAD, CAD), TIA (VBI, carotid artery ischemia), stroke.</p> <p>In most cases, there is little that could prospectively alert even the astute practitioner to an impending vascular event. Most patients are young, apparently healthy individuals, suffering from musculoskeletal complaints without any predisposing stroke risk factors.</p> <p>The most important points in the history and chief complaint, which would warn of a possible cervical artery disease, are^{308,318,354,387,388}:</p> <ol style="list-style-type: none"> a. S&S of VBI—the “5D’s and 3 N’s”: dizziness, dysphasia, dysarthria (hoarseness), drop attacks, diplopia (or other visual problems), ataxia of gait (hemiparesis), nausea (possibly with vomiting), numbness (hemianesthesia), nystagmus. b. S&S of carotid artery ischemia/stenosis: confusion, dysphasia, headache, anterior neck and/or facial pain, hemianesthesia, hemiparesis or monoparesis, visual field disturbances. c. Neck or occipital pain with sharp quality and severe intensity <i>or</i> severe & persistent headache that is sudden and unlike any previous experienced pain or headache (even when it is suspected the pain is of a musculoskeletal or neuralgic origin) <p style="text-align: center;">Special investigations [C]</p> <p>Any reports consistent with TIAs, prior stroke history (1% risk of recurrence/year), severe hypertension (noted in half of cases of spontaneous/idiopathic vertebral artery dissection), Hx of a major cervical spine trauma (attributed to 10% of vertebrobasilar syndrome), cardiac abnormalities that predispose to thrombus formation or myolytic arteriopathy should prompt close observation for neurologic status.^{308,318,389,390}</p> <p>Appropriate consultation and/or diagnostic procedures to evaluate the status of the cerebral circulation required in patients presenting with significant risk factors for cervical artery dissection. In such cases, approach the treatment with caution until a specific determination is made.^{308,318}</p>	<p>If patient reports active/existing VAD, CAD, TIA/stroke, DO NOT proceed with cervical manipulation</p> <p>Emergency care:</p> <p><i>It is vital to recognize the S&S of cerebrovascular ischemia and stroke in which case patient must be referred immediately to emergency facility for early and appropriate treatment.</i></p> <p>Additional information:</p> <ol style="list-style-type: none"> a. Sudden onset of acute and unusual headache and/or neck pain is present in 92% of cases of cervical artery dissection and may occur either spontaneously, after a specific activity, a significant neck trauma, or after cervical spine manipulation. b. Immediate or delayed neurologic S&S developing spontaneously, after a specific activity, a significant neck trauma, or after cervical manipulation; including the 5 D’s and 3 N’s are absolute contraindications to SMT. c. S&S of internal carotid artery dissection: most common presentation: headache and neck pain (90%), ischemic symptoms (50%-95%), Horner’s syndrome (<52%), visual scintillation (33%), monocular blindness (6%-30%), subjective bruit (25%-48%), impairment of taste (10%-19%).⁴⁰² <p>Should cervical artery problems be suspected, a thorough workup is indicated which may include²⁴⁵:</p> <ul style="list-style-type: none"> • Gold standard diagnostic tests (MRI/MRA) should be performed before implementing any manipulative procedure.³⁰⁸ • US imaging of the cervical vessels is typically used as a fast, noninvasive screen when pathology is suspected. Color-coded duplex ultrasonography can help diagnose vertebral artery (VA) occlusion and VBI by measuring blood flow velocity & flow volume and diameter of the VAs (including hypoplasia & asymmetry).^{354,403-406} However, US is operator dependant, lacks resolution, and has some limitations associated with vertebral artery evaluation. For these reasons, ultrasound may not be a viable screening tool.^{407,408}

Table 5 (continued)

Patient presentation	Comments
<p>Although several risk factors have been proposed for cervical artery dissection,^{308,318,387-399} current literature <i>does not assist</i> in identification of precipitating causes. VBIs may be considered the result of cumulative events over time/multifactorial.^{400,401} The patient history should include a system review and family history.</p>	<ul style="list-style-type: none"> • Catheter digital angiography is an invasive procedure with associated risk of stroke (≈1%). MRI/MRA considered method of choice as initial diagnosis and follow-up of craniocervical artery dissection. <p>See Mullins⁴⁰⁹ and Foerster et al²⁴⁵ or a review on stroke imaging</p>
<p style="text-align: center;">Further consideration</p>	<p style="text-align: center;">Further considerations</p> <ol style="list-style-type: none"> 1. External structures such as facet & uncovertebral joint osteophytes and presence of a retroarticular canal may potentially compress the vertebral artery.⁴¹⁰⁻⁴¹² However, no cases were found to be associated with either retroarticular canal or posterior ponticle in young adults.³⁰⁸ Although 1 case of VBI and one stroke have been associated with vertebral artery stenosis from osteophytes, cervical spondylosis does not appear to increase the risk of vertebrobasilar accidents after vertebral manipulation. If such were the case, one would expect older patients (>45 yoa) to be at heightened risk, but they are not.³⁸⁸⁻³⁹⁰ There is currently no evidence that in the absence of red flags in the adult patient, routine cervical spine radiographs before initiating manipulative therapy have any value as a mean to reduce rare, random, unpredictable vascular complications associated with neck movement, trivial trauma, or cervical manipulation.^{308,318,390} 2. Limiting neck rotation during SMT and applying the smallest force required to achieve therapeutic objective is advisable.^{311,413} 3. Migraine headache is a risk factor for stroke in young patients. When possible, treat all stroke modifiable risk factors aggressively, especially in women over age 35: smoking, oral contraceptives, presence of aura, >12 attacks of migraines/year, >12 y of migraine history.⁴¹⁴ 4. Tests and imaging studies that would help screen patients at risk of adverse vascular events before receiving cervical SMT would be of great help. However, such studies are not available presently.³⁰⁸ There is currently no evidence that in the absence of red flags, routine cervical spine radiographs before initiating manipulative therapy have any value as a means to reduce rare, random, unpredictable vascular complications associated with neck movement, trivial trauma, or cervical manipulation. 5. In patients with past a history of <i>major trauma to the cervical spine*</i>, MRA or angiography recommended before treating the neck as follow-up MRA indicates that these damaged vertebral arteries usually do not recanalize.⁴¹⁵⁻⁴²¹ <p>In a review of 80 cases of traumatic vertebral ischemia, MVA were thought to be responsible for 77.5% of cases, with delays between trauma and the onset of neurologic S&S of up to 5 y. The severity of trauma caused loss of consciousness (LOC) in 70% either at the time of injury or with subsequent episode of vertebrobasilar ischemia and short-term memory deficit in 64%. However, severity of trauma was mild for many, with no LOC, and no blow to the head.⁴²²</p>

ARE THERE POTENTIAL RISKS ASSOCIATED WITH CONVENTIONAL RADIOGRAPHS?

Although somewhat controversial,²⁰⁻²³ it is important to remember that health hazards of all forms of radiation are cumulative.²³⁻³⁰ The Biological Effects of Ionizing Radiation (BEIR) 2005 report released by the National Academy of Sciences adds further support to the “linear-no-threshold” model of cancer risk from ionizing radiation exposure.³¹ In summary, this report concludes that ionizing radiation is dangerous even at low doses and that there are no safe limits. Given the potential risks associated with conventional radiographs, only appropriate clinical indications can justify its use. In this regard, the need to confirm pathology, to follow the evolution of a pathology possibly affecting therapy, or to identify a clinically suspected contraindication to manipulative therapy is the best-documented reason. The benefits of all diagnostic studies must outweigh the risks and the inherent costs to the patient.^{10,32-40}

Tables 1-5 list the patient presentations, recommendations, and comments regarding diagnostic imaging for spine disorders. A list of abbreviations and glossary of terms used in the recommendations is in Appendix A. Appendix B provides a summary of the recommendations. Pertinent information for spinal disorders is presented in Appendix C. General indications for advanced imaging are in Appendix D. Typical effective ionized radiation dose for common imaging procedures is listed in Appendix E. Further recommended reading pertaining to magnetic resonance imaging of the spine is offered in Appendix F.

Practical Applications

- History taking and physical examination to exclude underlying sinister cause (red flags) and serious injuries (fracture and/or dislocation)
- Physical examination for signs of nerve compression
- Diagnostic triage (non specific back, neck, and extremity pain, with or without restriction of activity of daily living, radicular syndrome, presence of red flags)
- Radiographs not initially indicated for non specific acute, subacute, or persistent back and neck pain (no red flags)
- Consider conventional radiography after blunt trauma (according to decision rules), and if there is no improvement after 4-6 weeks of conservative care or increasing disability
- Consider conventional radiography and specialized imaging for radicular syndrome and presence of red flags
- Urgent specialized imaging for back and neck pain with adverse features: sphincter or gait disturbance, saddle anesthesia, severe or progressive neurologic deficit, systemic illness (cancer, infection), vascular causes (suspected abdominal/thoracic aorta aneurysm), or cervical artery dissection.

ACKNOWLEDGMENT

We appreciate the efforts of the many people who assisted in this process, listed in Appendix G.

SUPPLEMENTARY DATA

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.jmpt.2007.10.003 and in tables D1-D3 in Appendix D of Reference 2.

REFERENCES

1. Shiffman RN, Shekelle P, Overhage JM, Slutsky J, Grimshaw J, Deshpande AM. Standardized reporting of clinical practice guidelines: a proposal from the conference on guideline standardization. *Ann Intern Med* 2003;139:493-8.
2. Bussi eres AE, Peterson C, Taylor JAM. Diagnostic imaging practice guidelines for musculoskeletal complaints in adults— an evidence-based approach: introduction. *J Manipulative Physiol Ther* 2007;30:617-83.
3. Whiting P, Rutjes AWS, Dinnes J, Reitma JB, Bossuyt PMM, Kleijnen JK. Development of validation of methods for assessing the quality of diagnostic accuracy studies. Chap 9. *Health Technol Assess (Rockv)* 2004;8:59-65.
4. The Agree Collaboration. Development and validation of an international appraisal instrument for assessing the quality of clinical practice guidelines: the AGREE project. *Qual Saf Health Care* 2003;12:18-23.
5. Ricci S, Celani MG, Righetti E. Development of clinical guidelines: methodological and practical issues. *Neurol Sci* 2006;27(Suppl 3):S228-30.
6. European Commission. Radiation protection 118. Referral guidelines for imaging in conjunction with the UK Royal College of Radiologists; Luxembourg; 2001.
7. Harbour R, Miller J. for the SIGN grading review group. A new system for grading recommendations in evidence based guidelines. *BMJ* 2001;323:334-6.
8. Simmons ED, Guyer RD, Graham-Smith A, Herzog R. Radiographic assessment for patients with low back pain. *Spine* 1995;20:1839-41.
9. Cooperstein R, Perle SM, Gleberzon BJ, Peterson DH. Flawed trials, flawed analysis: why CPB should avoid rating itself. *J Can Chiropr Assoc* 2006;50:97-102.
10. Sackett DL, Haynes RB, Guyatt GH, Tugwell P. *Clinical epidemiology: a basic science for clinical medicine*. 2nd ed. Boston: Little Brown and Company; 1991. p. 441.
11. Sackett DL, Haynes RB, Tugell PX, Trout KS, Stoddard GL. How to read clinical journals: II. To learn about diagnostic test. *Can Med Assoc J* 1981;124:703-9.
12. Thornbury JR. Clinical efficacy of diagnostic imaging: love it or leave it. *AJR Am J Roentgenol* 1994;162:1-8.
13. Belsey J, Snell T, editors. What is evidence-based medicine?; 2000. [Serial on the internet]; [cited 2007 Jul 14]; 1(2): [about 6 p.]. Available from <http://www.evidence-based-medicine.co.uk>.
14. Evidence-Based Health care Project. [Home page on the Internet], Minnesota. Regents of the University of Minnesota; [updated: 2007, Nov 9; cited 2007 Jul 13]. Available from: <http://evidence.ahc.umn.edu/>.
15. Raine R, Sanderson C, Black N. Developing clinical guidelines: a challenge to current methods. *BMJ* 2005;331:631-3.
16. Mootz RD. Chiropractic’s current state: impact for the future. *J Manipulative Physiol Ther* 2007;30:1-3.

17. Broughton R, Rathbone B, editors. What makes a good clinical guideline? 2001. [Serial on the internet]; [cited 2007 Jul 9]; 1(11): [about 8 p.]. Available from: <http://www.evidence-based-medicine.co.uk>.
18. Mootz RD, Cherkin DC, Odegard CE, Eisenberg DM, Barassi JP, Deyo RA. Characteristics of chiropractic practitioners, patients, and encounters in Massachusetts and Arizona. *J Manipulative Physiol Ther* 2005;28:645-53.
19. Koes BW, van Tulder MW, Ostelo R, Burton KA, Waddell G. Clinical guidelines for the management of low back pain in primary care: an international comparison. *Spine* 2001;26:2504-13.
20. Oakley PA, Harrison DD, Harrison DE, Hass JW. On "phantom risks" associated with diagnostic ionizing radiation: evidence in support of revising radiography standards and regulations in chiropractic. *J Can Chiropr Assoc* 2005;49:264-9.
21. Bussi eres AE, Ammendolia C, Peterson C, Taylor JAM. Ionizing radiation exposure—more good than harm? The preponderance of evidence does not support abandoning current standards and regulations. *J Can Chiropr Assoc* 2006;50:1003-6.
22. Luckey TD. Nurture with ionizing radiation: a provocative hypothesis. *Nutr Cancer* 1999;34:1-11.
23. Masse R. Ionizing radiation. *CR Acad Sci III* 2000;323:633-40.
24. Trosko JE. Role of low-level ionizing radiation in multi-step carcinogenic process. *Health Phys* 1996;70:812-22.
25. Evans BF, Wennberg JE, McNeil BJ. The influence of diagnostic radiography on the incidence of breast cancer and leukemia. *N Engl J Med* 1996;315:810-5.
26. Ron E. Ionizing radiation and cancer risk: evidence from epidemiology. *Radiat Res* 1998;150(5 Suppl):S30-41.
27. Kleinerman RA. Cancer risks following diagnostic and therapeutic radiation exposure in children. *Pediatr Radiol* 2006;36(14 suppl):S121-5.
28. Levy AR, Golberg MS, Mayo NE, Hanley JA, Poitras B. Reducing the life time risk of cancer from spinal radiographs among people with adolescent idiopathic scoliosis. *Spine* 1996;21:1540-8.
29. Assessment and Management of Cancer Risks from Radiological and Chemical Hazards 1998 Health Canada, Atomic Energy Control Board. [Monograph on the Internet], Ottawa: [cited 2007 Jul 14]. Available from: http://www.hc-sc.gc.ca/ewh-semt/alt_formats/hecs-sesc/pdf/pubs/radiation/98ehd-dhm216/98ehd-dhm216_e.pdf.
30. UNSCEAR. Sources and effects of ionizing radiation. Report of the United Nations Scientific Committee on the effects of atomic radiation to the General Assembly. [Monograph on the Internet], Geneva: UNSCEAR; 2000 Vol I and II, [cited 2007 Jul 14]. Available from: <http://www.unscear.org/unscear/en/publications.html>.
31. Committee to Assess Health Risks from Exposure to Low Levels of Ionizing Radiation, National Researcher Council, Health risks from exposure to low levels of ionizing radiation: BEIR VII—Phase 2. [Monograph on the Internet], Washington: National Academy Press; 2006 [cited 2007 Jul 14]. Available from: <http://books.nap.edu/catalog/11340.html>.
32. Berrington de Gonzalez A, Darby S. Risk of cancer from diagnostic X-rays: estimates for the UK and 14 other countries. *Lancet* 2004;363:345-51.
33. Peterson C, Hsu W. Indications for and use of x-rays. In: Haldeman S, editor. Principles and Practice of Chiropractic. 3rd ed. New York: McGraw-Hill; 2005. p. 661-81.
34. Owens JP, Ruth G, Keir MJ, Spencer H, Richardson D, Richardson A, et al. A survey of general practitioners opinions on the role of radiology in patients with low back pain. *Br J Gen Pract* 1990;40:98-101.
35. Halpin SF, Yeoman L, Dundas DD. Radiographic examination of the lumbar spine in a community hospital: an audit of current practice. *BMJ* 1991;303:813-5.
36. Liang M, Katz JN, Frymoyer JW. Conventional radiographs in evaluating the spine. In: Frymoyer JW, editor. *The Adult Spine*. New York: Raven Press; 1991. p. 699-718.
37. Mootz RD, Hoffman LE, Hansen DT. Optimising clinical use of radiography and minimizing radiation exposure in chiropractic practice. *Top Clin Chiro* 1997;4:34-44.
38. Nachemson A, Vingard E. Assessment of patients with neck and back pain: A best-evidence synthesis, chap 9. In: Nachemson A, Jonsson E, editors. *Neck and back pain—The scientific evidence of causes, diagnosis and treatment*. Philadelphia: Lippincott Williams & Wilkins; 2000. p. 189-235.
39. Ammendolia C, Bombardier C, Hogg-Johnson S, Glazier R. Views on radiography use for patients with acute low back pain among chiropractors in an Ontario community. *J Manipulative Physiol Ther* 2002;25:511-20.
40. Vader JP, Terraz O, Perret L, Aroua A, Valley JF, Burnand B. Use of and irradiation from plain lumbar radiography in Switzerland. *Swiss Med Wkly* 2004;134:419-22.
41. Coulter I, Shekelle P. Chiropractic in North America: A descriptive analysis. *J Manipulative Physiol Ther* 2005;28:83-9.
42. Brooks P, March L, Bogduk N, Bellamy N, Spearing N, Fraser M, et al. Acute Australian Musculoskeletal Pain Guidelines Group. Evidence-based Management of acute musculoskeletal pain. Brisbane: National Health and Medical Research Council Australian Academic Press Pty Ltd; 2003 [Monograph on the Internet]; [cited 2006 Jan 30]. Available from: http://www.nhmrc.gov.au/publications/synopses/_files/cp94.pdf.
43. Van den Bosch MAAJ, Hollingworth W, Kinmonth AL, Dixon AK. Evidence against the use of lumbar spine radiography for low back pain. *Clin Radiol* 2004;59:69-76.
44. Jarvik JG, Deyo RA. Diagnostic evaluation of low back pain with emphasis on imaging. *Ann Intern Med* 2002;137:586-97.
45. Holmes JF, Panacek EA, Miller PQ, Lapdis AD, Mower WR. Prospective evaluation of criteria for obtaining thoracolumbar radiographs in trauma patients. *J Emerg Med* 2003;24:1-7.
46. Hsu JM, Joseph T, Ellis AM. Thoracolumbar fractures in blunt trauma patients: guidelines for diagnosis and imaging. *Injury* 2003;34:426-33.
47. Bagley LJ. Imaging of spinal trauma. *Radiol Clin North Am* 2006;44:1-12.
48. Berry GE, Adams S, Harris MB, Boles CA, McKernan MG, Collinson F, Hoth JJ, Meredith JW, Chang MC, Miller PR. Are plain radiographs of the spine necessary during evaluation after blunt trauma? Accuracy of screening torso computed tomography in thoracic/lumbar spine fracture diagnosis. *J Trauma* 2005;59:1410-3.
49. Braddock EJ, Greenlee J, Hammer RE, Johnson SF, Martello MJ, O'Connell MR, et al. Manual medicine guidelines for musculoskeletal injuries [online publication]. California: Academy for Chiropractic Education; 2004. p. 25. [39 references]. Available from: <http://www.guideline.gov/>.
50. Reichtine GR. Nonoperative management and treatment of spinal injuries. *Spine* 2006;31(Suppl):S22-7.
51. Harrop J, Vaccaro AR, Hurlbert RJ, Wilsey JT, Baron EM, Shaffrey CI, Fisher C, Dvorak M, Oner FC, Wood KB, et al. Intrarater and interrater reliability and validity in the

- assessment of the mechanism of injury and integrity of the posterior ligamentous complex: a novel injury severity scoring system for thoracolumbar injuries. Invited submission from the joint section meeting on disorders of the spine and peripheral nerves, March 2005. *J Neurosurg Spine* 2006;4:118-22.
52. Vaccaro AR, Lehman RA, Hurlbert RJ, Anderson PA, Harris M, Hedlund R, Harrop J, Dvorak M, Wood K, Fehlings MG, Fisher C, et al. A new classification of thoracolumbar injuries: the importance of injury morphology, the integrity of the posterior ligament complex, and neurologic status. *Spine* 2005;30:2325-33.
 53. Vaccaro AR, Baron EM, Sanfilippo J, Jacoby S, Steuve J, Grossman E, et al. Reliability of a novel classification system for thoracolumbar injuries: The Thoracolumbar Injury Severity Score. *Spine* 2006;31(Suppl):S62-9.
 54. Van Goethem JWM, Maes M,  zsarlak  , van den Hauwe L, Parizel PM. Imaging in spinal trauma. *Eur Radiol* 2005;15:582-90.
 55. Brown CV, Antevil JL, Sise MJ, Sack DI. Spiral computer tomography for the diagnosis of cervical, thoracic, and lumbar spine fractures: its time has come. *J Trauma* 2005;58:890-5.
 56. Gray L, Vandemark R, Hays M. Thoracic and lumbar trauma. *Semin Ultrasound CT MR* 2001;22:125-34.
 57. Miller LA. Chest wall, lung, and pleural space trauma. *Radiol Clin North Am* 2006;44:213-24.
 58. Kuklo TR, Potter BK, Ludwig SC, Anderson PA, Lindsey RW, Vaccaro AR. Spine Trauma Study Group. Radiographic measurement techniques for sacral fractures consensus statement of the Spine Trauma Study Group. *Spine* 2006;31:1047-55.
 59. McCormick JP, Morgan SJ, Smith WR. Clinical effectiveness of the physical examination in diagnosis of posterior pelvic ring injuries. *J Orthop Trauma* 2003;17:257-61.
 60. Hodges SD, Eck JC, Humphreys SC. A treatment and outcomes analysis of patients with coccydynia. *Spine J* 2004;4:138-40.
 61. De Andres J, Chaves S. Coccygodynia: a proposal for an algorithm for treatment. *J Pain* 2003;4:257-66.
 62. Maigne JY, Doursounian L, Chatellier G. Causes and mechanisms of common coccydynia: role of body mass index and coccygeal trauma. *Spine* 2000;25:3072-9.
 63. RCR Working Party. Making the Best Use of a Department of Clinical Radiology: Guidelines for Doctors. 5th ed. London: Royal College of Radiologists; 2003. Available from: <http://www.rcr.ac.uk>.
 64. Maigne JY, Lagauche D, Doursounian L. Instability of the coccyx in coccydynia. *J Bone Joint Surg Br* 2000;82:1038-41.
 65. Stiell IG, Wells G, Vandemheen KL, Clement CM, Lesiuk H, De Maio VL, Laupacis A, Schull M, et al. The Canadian cervical spine rule for radiography in alert and stable trauma patients. *JAMA* 2001;286:1841-8.
 66. Stiell IG, Clement CM, McKnight RD, Brison R, Schull MJ, Rowe BH, et al. The Canadian C-Spine Rule versus the NEXUS low-risk criteria patients with trauma. *N Engl J Med* 2003;349:2510-8.
 67. Hoffman JR, Mower WR, Wolfson AB, Tood KH, Zucker M, National Emergency X-Radiography Utilisation Study Group. Validity of a set of clinical criteria to rule out injury to the cervical spine in patients with blunt trauma. *N Engl J Med* 2000;343:94-9.
 68. Bandiera G, Stiell IG, Wells GA, Clement C, De Maio V, Vandemheen KL. The Canadian C-Spine Rule performs better than unstructured physician judgment. *Ann Emerg Med* 2003;42:395-402.
 69. No authors listed. Radiographic assessment of the cervical spine in asymptomatic trauma patients. *Neurosurgery* 2002;50(3 Suppl):S30-5.
 70. Provinciali L, Baroni M. Clinical approaches to whiplash injuries: a review. *Crit Rev Phys Rehab Med* 1999;11:339-68.
 71. Maak TG, Tominaga Y, Panjabi MM, Ivancic PC. Alar, transverse, and apical ligament strain due to head-turned rear impact. *Spine* 2006;31:632-8.
 72. Touger M, Gennis P, Nathanson N, Lowery DW, Pollark CV, Hoffman JR, Mower WR. Validity of a decision rule to reduce cervical spine radiography in elderly patients with blunt trauma. *Ann Emerg Med* 2002;40:287-93.
 73. Foreman SM, Croft AC. Whiplash injuries—the acceleration/deceleration syndrome. 2nd ed. Baltimore: William & Wilkins; 1995.
 74. Spitzer WO, Skovron ML, Salmi LR, Cassidy JD, Duranceau J, Suissa S, Zeiss E. Scientific monograph of the Quebec Task Force on Whiplash-Associated Disorders: redefining “whiplash” and its management. *Spine* 1995;20(8 Suppl):1S-73S.
 75. Duranceau JA. Les maladies du syst eme locomoteur. La gu erison des tissus mous. *Le Clinicien* 2001:121-32.
 76. Barnsley L, Lord S, Bogduk N. Whiplash injury. *Pain* 1994;58:283-307.
 77. Rogers LF. Traumatic lesions of bones and joints. In: Juhl JH, Crummy AB, editors. *Essentials of radiologic imaging*. 5th ed. Philadelphia: J.B. Lippincott Company; 1987. p. 35-84.
 78. Rengachary SS. Bone morphogenetic proteins: basic concepts. Article 2. *Neurosurg Focus* 2002;13:1-6.
 79. Rosen H. Fracture healing and pseudoarthrosis. In: Taveras JM, Ferrucci JT, Dalinka MK, editors. *Radiology Diagnosis—Imaging Intervention. Skeletal Radiology*. Philadelphia: Lippincott-Raven; 1998. p. (5)110:3.
 80. Pilitsis J, Lucas D, Regachary SR. Bone healing and spinal fusion. Article 1. *Neurosurg Focus* 2002;13:1-6.
 81. Walker DH, Wright NM. Bone morphogenetic proteins and spinal fusion. Article 3. *Neurosurg Focus* 2002;13:1-13.
 82. Verhagen AP, Scholten-Peeters GG, de Bie RA, Bierma-Zeinstra SM. Conservative treatments for whiplash. *Cochrane Database Syst Rev* 2004:CD003338.
 83. C  te P, Hogg-Johnson S, Cassidy JD, Carroll L, Frank JW, Bombardier C. Initial pattern of clinical care and recovery from whiplash injuries: a population-based cohort study. *Arch Intern Med* 2005;165:2257-63.
 84. Sterling M, Jull G, Vicenzino B, Kenardy J, Darnell R. Physical and psychological factors predict outcome following whiplash injury. *Pain* 2005;114:141-8.
 85. Sterling M, Jull G, Kenardy J. Physical and psychological factors maintain long-term predictive capacity post-whiplash injury. *Pain* 2006;122:102-8.
 86. C  te P, Cassidy JD, Carroll L. Is a lifetime history of neck injury in a traffic collision associated with prevalent neck pain, headache and depressive symptomatology? *Accid Anal Prev* 2000;32:151-9.
 87. Berglund A, Alfredsson L, Cassidy JD, Jensen I, Nygren A. The association between exposure to a rear-end collision and future neck or shoulder pain: a cohort study. *J Clin Epidemiol* 2000;53:1089-94.
 88. Berglund A, Alfredsson L, Jensen I, Cassidy JD, Nygren A. The association between exposure to a rear-end collision and future health complaints. *J Clin Epidemiol* 2001;54:851-6.
 89. Sterling M, Jull G, Vicenzino B, Kenardy J. Characterization of acute whiplash-associated disorders. *Spine* 2004;29:182-8.
 90. Hanson JA, Blackmore CC, Mann FA, Wilson AJ. Cervical spine injury: a clinical decision rule to identify high-risk patients for helical CT screening. *AJR Am J Roentgenol* 2000;174:713-7.

91. Antevil JL, Sise MJ, Sack DI, Kidder B, Hopper A, Brown CV. Spiral computed tomography for the initial evaluation of spine trauma: a new standard of care? *Trauma* 2006;61:382-7.
92. Bub LD, Blackmore CC, Mann FA, Lomoschitz FM. Cervical spine fractures in patients 65 years and older: a clinical prediction rule for blunt trauma. *Radiology* 2005;234:143-9.
93. Miller JS, Craw MM. Diagnostic imaging of the cervical spine following whiplash-induced injury. *Top Clin Chiro* 1997;4:26-33.
94. Jonsson H, Cesarini K, Sahlstedt B, Rauschnig W. Findings and outcome in whiplash-type neck distortions. *Spine* 1994;19:2733-42.
95. Moore TA, Vaccaro AR, Anderson PA. Classification of lower cervical spine injuries. *Spine* 2006;31(11 Suppl):S37-43.
96. Radiographic assessment of the cervical spine in symptomatic trauma patients. *Neurosurgery* 2002;50(3 Suppl):S36-43.
97. Insko EK, Gracias VH, Gupta R, Goettler CE, Gaieski DF, Dalinka MK. Utility of flexion and extension radiographs of the cervical spine in the acute evaluation of blunt trauma. *J Trauma* 2002;53:426-9.
98. Pollack CV, Hendey GW, Martin DR, Hoffman JR, Mower WR. Use of flexion-extension radiographs of the cervical spine in blunt trauma. *Ann Emerg Med* 2001;38:8-11.
99. Lurie JD. What diagnostic tests are useful for low back pain? *Best Pract Res Clin Rheumatol* 2005;19:557-75.
100. Mootz RD, Cherkov DC, Odegard CE, Eisenberg DM, Barassi JP, Deyo RA. Characteristics of chiropractic practitioners, patients, and encounters in Massachusetts and Arizona. *J Manipulative Physiol Ther* 2005;28:645-53.
101. Rossignol M, Arsenault B. Guide de pratique—Clinique des lombalgies interdisciplinaires en premi ere ligne, Direction de la Sant e Publique de Montr eal; 2006. p. 40.
102. Cassidy JD, Cote P, Carroll LJ, Kristman V. Incidence and course of low back pain episodes in the general population. *Spine* 2005;30:2817-23.
103. Waddell G, Burton AK. Occupational health guidelines for the management of low back pain at work: evidence review. *Occup Med* 2001;51:124-35.
104. Borenstein DG, O'Mara JW, Boden SD, Lauerman WC, Jacobson A, Platenberg C, et al. The value of magnetic resonance imaging of the lumbar spine to predict low-back pain in asymptomatic subjects: a seven-year follow-up study. *J Bone Joint Surg Am* 2001;83-A:1306-11.
105. Kendrick D, Fielding K, Bentley E, Kerslake R, Miller P, Pringle M. Radiography of the lumbar spine in primary care patients with low back pain: randomised controlled trial. *BMJ* 2001;322:400-5.
106. Kerry S, Hilton S, Dundas D, Rink E, Oakeshott P. Radiography for low back pain: a randomised controlled trial and observational study in primary care. *Br J Gen Pract* 2002;52:469-74.
107. Miller P, Kendrick D, Bentley E, Fielding K. Cost-effectiveness of lumbar radiography in primary care patient with low back pain. *Spine* 2002;27:2291-7.
108. Institute for Work & Health, editor. Guidelines: Special studies—Plain X-Rays—Summary; 1999. [Monograph on the Internet]; [cited 2003 Nov 30]. Available from: <http://www.backguide.com/guidelin/speplsu.htm>.
109. Kendrick D, Fielding K, Bentley E, Miller P, Kerslake R, Pringle M. The role of radiography in primary care patients with low back pain of at least 6 weeks duration: a randomised (unblinded) controlled trial. *Health Technol Assess* 2001;5:1-69.
110. Nachemson AL. The lumbar spine: an orthopaedic challenge. *Spine* 1976:1.
111. Bigos SJ, Bowyer O, Braen GR, et al. Acute low-back pain problems in adults: clinical guideline no.14. AHCPR publication no. 95-0642. Rockville, MD: Agency for Health Care Policy and Research, Public Health Service, United States Department of Health Human Services; 1994.
112. Bradley WG, Seidenwurm DJ, Brunberg JA, Davis PC, La Paz DE, Dormont D, et al. In: Expert Panel on Neurologic Imaging, editor. *Low Back Pain*. Reston (VA): American College of Radiology (ACR); 2005. p. 7. [online publication] [23 references]. Available from: <http://www.guideline.gov>.
113. Van M, Tulder W, Assendelf WJ, Koes BW, Bouter LM. Spinal radiographic findings and low back pain. A systematic review of observational studies. *Spine* 1997;22:447-534.
114. Koes BW, van Tulder MW, Ostelo R, Burton AK, Waddell G. Clinical guidelines for the management of low back pain in primary care—an international comparison. *Spine* 2001;26:2504-14.
115. Brown A, Angus D, Chen S, Tang Z, Milne S, Pfaff J, Li H, Mensinkai S, editors. *Costs and Outcomes of Chiropractic Treatment for Low Back Pain*; 2005. [Monograph on the Internet] Technology Report: Chiropractic treatment. (56). July. Available from: <http://www.ccohta.ca>.
116. Isaac DM, Marinac J, Sun C. Radiograph use in low back pain: a United States emergency department database analysis. *J Emerg Med* 2004;26:37-45.
117. Fickel TE. Organ-specific dosimetry in spinal radiography: an analysis of genetic and somatic effects. *J Manipulative Physiol Ther* 1988;11:3-9.
118. Gonzalez L, Va no E, Oliete S, Manrique J, Hern ez JM, Lahuerta J, Ruiz J. Report of an image quality and dose audit according to directive 97/43/Euratom at Spanish private radiodiagnostics facilities. *Br J Radiol* 1999;72:186-92.
119. Johnston DA, Brennan PC. Reference dose levels for patients undergoing common diagnostic X-ray examinations in Irish hospitals. *Br J Radiol* 2000;73:393-402.
120. Luomas K, Vehma T, Raininko R, Luukkonen R, Riihimaki H. Lumbosacral transitional vertebra: relation to disc degeneration and low back pain. *Spine* 2004;29:200-5.
121. Ammendolia C, Bombardier C, Hogg-Johnson S, Glazier R. Views on radiography use for patients with acute low back pain among chiropractors in an Ontario community. *J Manipulative Physiol Ther* 2002;25:511-20.
122. Ammendolia C. Radiographic anomalies that may alter chiropractic intervention strategies found in a New Zealand population. *J Manipulative Physiol Ther* 2005;28:375.
123. Assessment and Management of Cancer Risks from Radiological and Chemical Hazards0-662-82719-8; 1998 [Monograph on the Internet], Health Canada, Atomic Energy Control Board. Available from: http://www.hc-sc.gc.ca/ewh-semt/alt_formats/hecs-sesc/pdf/pubs/radiation/98ehd-dhm216/98ehd-dhm216_e.pdf.
124. Berrington de Gonz alez A, Darby S. Risk of cancer from diagnostic X-rays: estimates for the UK and 14 other countries. *Lancet* 2004;363:345. <http://www.thelancet.com>.
125. Beutler WF, Fredrickson BE, Murtland A, Sweeney CA, Grant WD, Baker D. The natural history of spondylolysis and spondylolisthesis: 45-year follow-up evaluation. *Spine* 2003;28:1027-35.
126. Mireau D, Cassidy JD, McGregor M, Kirkaldy-Willis WH. A comparison of the effectiveness of spinal manipulative therapy for low back pain with or without spondylolysthesis. *J Manipulative Physiol Ther* 1987;10:49-55.
127. Ventura JM, Justice B. Need for multiple diagnoses in the presence of spondylolisthesis. *J Manipulative Physiol Ther* 1988;11:41-2.
128. Halpin SF, Yeoman L, Dundas DD. Radiographic examination of the lumbar spine in a community hospital: an audit of current practice. *BMJ* 1991;303:813-5.

129. Health Risks from Exposure to Low Levels of Ionizing Radiation : BEIR VII—Phase 2. [Monograph on the Internet], National Academy of Sciences 2005. Available from: <http://books.nap.edu/catalog/11340.html>.
130. Mootz RD, Hoffman LE, Hansen DT. Optimising clinical use of radiography and minimizing radiation exposure in chiropractic practice. *Top Clin Chiro* 1997;4:34-44.
131. Liang M, Kattz JN, Frymoyer JW. Plain film radiographs in evaluating the spine. In: Frymoyer JW, editor. *The Adult Spine*. New York: Raven Press; 1991. p. 699-718.
132. Leonardi M, Simonnetti L, Agati R. Neuroradiology of spine degenerative disease. *Best Pract Res* 2002;16:59-87.
133. Owens JP, Ruth G, Keir MJ, Spencer H, Richardson D, Richardson A, et al. A survey of general practitioners opinions on the role of radiology in patients with low back pain. *Br J Gen Pract* 1990;40:98-101.
134. Hollingworth W, Todd CJ, King H, Dixon AK, Karia KR, Kinmonth AL. Primary care referrals for lumbar spine radiography: diagnostic yield and clinical guidelines. *Br J Gen Pract* 2002;52:475-80.
135. Jarvik JG, Hollingworth W, Haegerty PJ, Haynor DR, Boyko EJ, Deyo RA. Three-years incident of low back pain in an initially asymptomatic cohort: clinical and imaging risk factors. *Spine* 2005;30:1541-8.
136. Peterson CK, Bolton J, Hsu W, Wood A. A cross-sectional study comparing pain and disability levels in patients with low back pain with and without transitional lumbosacral vertebrae. *J Manipulative Physiol Ther* 2005;28:570-4.
137. Taylor JAM. The role of radiography in evaluating the vertebral subluxation. In: Gatterman M, editor. *Foundations of chiropractic: the subluxation*. St-Louis (Mo): Mosby-Year Book; 1995. p. 68-86.
138. Hass M, Taylor JAM, Gillette RG. The routine use of radiographic spinal displacement analysis: A dissent. *J Manipulative Physiol Ther* 1999;22:254-9.
139. Schultz G, Bassano J. Is radiography appropriate for detecting subluxations? *Top Clin Chiro* 1997;4:1-8.
140. Harrison DD, Harrison DE, Troyanovich SJ, Hansen DJ. The anterior–posterior full spine view: the worst radiographic view for determination of mechanics of the spine. *Chiro Techn* 1996;8:163-70.
141. Coleman RR, Bernard BB, Harrison DE. The effects of x-axis vertebral translation on projected y-axis vertebral rotation. *J Manipulative Physiol Ther* 1998;21:333-40.
142. Coleman RR, Bernard BB, Harrison DE. Correlation and quantification of projected 2-dimensional radiographic images with actual 3-dimensional y-axis vertebral rotations. *J Manipulative Physiol Ther* 1999;22:21-5.
143. Coleman RR, Harrison DE, Fischer T, Harrison SO. Correlation and quantification of relative 2-dimensional projected vertebral endplate z-axis rotations with 3-dimensional y-axis vertebral rotations and focal spot elevations. *J Manipulative Physiol Ther* 2000;23:414-9.
144. Coleman RR, Harrison DE, Bernard BB. The effects of combined x-axis translations and y-axis rotations on projected lamina junction offset. *J Manipulative Physiol Ther* 2001;24:509-13.
145. Coleman RR, Thomas IW. Movement of the projected pedicles relative to the projected vertebral body in a fourth lumbar vertebra during axial rotation. *J Manipulative Physiol Ther* 2004;27:403-7.
146. Nordin M, Balagus F, Cedraschi C. Nonspecific lower-back pain: surgical versus nonsurgical treatment. *Clin Orthop Relat Res* 2006;443:156-67.
147. Bronfort G, Haas M, Evans RL, Bouter LM. Efficacy of spinal manipulation and mobilization for low back pain and neck pain: a systematic review and best evidence synthesis. *Spine J* 2004;4:335-56.
148. Shekelle PG, Adams AH, et al. The appropriateness of spinal manipulation for low back pain: Indication and ratings by a multidisciplinary expert panel. Santa Monica (Calif): RAND; 1991. Monograph No. R-4025/3-CCR/FCCR.
149. Shekelle PG, Adams AH, Chassin MR, Hurwitz EL, Brook RH. Spinal manipulation for low back pain. *Ann Intern Med* 1992;117:590-8.
150. Peterson CK, Bolton JE, Wood AR. A cross-sectional study correlating lumbar spine degeneration with disability and pain. *Spine* 2000;25:218-23.
151. Kauppila LI, McAlindon T, Evans S, Wison PW, Kiel D, Feldon DT. Disc degeneration/back pain and calcification of the abdominal aorta. A 25-year follow-up study in Framingham. *Spine* 1997;22:1642-7.
152. Karasik D, Kiel DP, Kiely DK, Cupples LA, Wilson PW, O'Donnell CJ, Felson DT. Abdominal aortic calcification and exostoses at the hand and lumbar spine: the Framingham Study. *Calcif Tissue Int* 2006;78:1-8.
153. Pye SR, Reid DM, Smith R, Adams JE, Nelson K, Silman AJ, O'Neill TW. Radiographic features of lumbar disc degeneration and self-reported back pain. *J Rheumatol* 2004;31:753-8.
154. Phillips RB, Frymoyer JW, Mac B, Pherson V, Newburg AH. Low back pain: A radiographic enigma. *J Manipulative Physiol Ther* 1986;9:183-7.
155. Peterson C, Hsu W. Indications for and use of x-rays. In: Haldeman S, editor. *Principles and Practice of Chiropractic*. 3rd ed. New York: McGraw-Hill; 2005. p. 661-81.
156. Adams MA, Dolan P. Spine biomechanics. *J Biomech* 2005;38:1972-83.
157. Weber H, Holme I, Amlie E. The natural course of acute sciatica with nerve root symptoms in a double-blind placebo-controlled trial evaluating the effect of piroxicam. *Spine* 1993;18:1433-8.
158. Vucetic N, Astrand P, Guntner P, Svensson O. Diagnosis and prognosis in lumbar disc herniation. *Clin Orthop Relat Res* 1999;116-22.
159. Vroomen PC, de Krom MC, Knottnerus JA. Predicting the outcome of sciatica at short-term follow-up. *Br J Gen Pract* 2002;52:119-23.
160. Cassidy JD, Thiel HW, Kirkaldy-Willis WH. Side posture manipulation for lumbar intervertebral disc herniation. *J Manipulative Physiol Ther* 1993;16:96-103.
161. Oliphant D. Safety of spinal manipulation in the treatment of lumbar disk herniation: A systematic review and risk assessment. *J Manipulative Physiol Ther* 2004;27:197-210.
162. Crawford CM, Hannan RF. Management of acute disk herniation presenting as mechanical low back pain. *J Manipulative Physiol Ther* 1999;22:235-44.
163. Busse JW, Hsu WS. Rapid progression of acute sciatica to cauda equina syndrome. *J Manipulative Physiol Ther* 2001;24:350-5.
164. Vroomen PC, de Krom MC, Knottnerus JA. Diagnostic value of history and physical examination in patients suspected of sciatica due to disc herniation: a systematic review. *J Neurol* 1999;246:899-906.
165. Heli ovaara M, Impivaara O, Sievers K, Melkas T, Knekt P, Korpi J, Aromaa A. Lumbar disc syndrome in Finland. *J Epidemiol Community Health* 1987;41:251-8.
166. Deyo RA, Loeser JD, Bigos SJ. Herniated lumbar intervertebral disc. *Ann Intern Med* 1990;112:598-603.
167. Deyo RA, Rainville J, Kent DL. The rational clinical examination. What can the history and physical examination tell us about low back pain? *JAMA* 1992;268:760-5.
168. Andersson GB, Deyo RA. History and physical examination in patients with herniated lumbar discs. *Spine* 1996;21:10S-8S.

169. Ioto K, Aebi M, Alini M. Aging and degeneration: Cause and effect of intervertebral disc damage. In: Gunzburg R, Szpalski M, editors. *Lumbar disc herniation*. Philadelphia: Lippincott Williams & Wilkins; 2002. p. 72-83.
170. Leino-Arjas P, Kaila-Kangas L, Kauppinen T, Notkola V, Keskim aki I, Multane P. Occupational exposure and inpatient hospital care for lumbar intervertebral disc disorders among Finns. *Am J Ind Med* 2004;46:513-20.
171. Awad JN, Moskovich R. Lumbar disc herniations: surgical versus nonsurgical treatment. *Clin Orthop Relat Res* 2006;443:183-97.
172. Weinstein JN, Lurie JD, Tosteson TD, Skinner JS, Hanscom B, Tosteson AN, Herkowitz H, Fischgrund J, Cammisa FP, Albert T, Deyo RA. Surgical vs nonoperative treatment for lumbar disk herniation: the Spine Patient Outcomes Research Trial (SPORT) observational cohort. *J Am Med Ass* 2006;296:2451-9.
173. Quon JA, Cassidy DJ, O'Connor SM, Kirkaldy-Willis WH. Lumbar intervertebral disc herniation: treatment by rotational manipulation. *J Manipulative Physiol Ther* 1989;12:220-7.
174. Hubka MJ, Taylor JAM, Schultz GD, Traina AD. Lumbar intervertebral disc herniation: Chiropractic management using flexion, extension, and rotational manipulative therapy. *Chiropractic Technique* 1991;3:5-12.
175. Peterson DH. Principles of adjustive technique. In: Bergmann TF, Peterson DH, Lawrence DJ, editors. *Chiropractic Technique*. Philadelphia: Churchill Livingstone; 1993. p. 123-95.
176. Stern PJ, C ot e P, Cassidy JD. A series of consecutive cases of low back pain with radiating leg pain treated by chiropractors. *J Manipulative Physiol Ther* 1995;18:335-42.
177. Bergmann TF, Jongeward BV. Manipulative therapy in lower back pain with leg pain and neurological deficit. *J Manipulative Physiol Ther* 1998;21:288-94.
178. Jensen A, Nykoliation J. Successful resolution of signs and symptoms from L4 radiculopathy with spinal manipulation: a case report. *J Can Chiropr Assoc* 2003;47:116-20.
179. Lisi AJ, Holmes EJ, Ammendolia C. High-velocity low-amplitude spinal manipulation for symptomatic lumbar disk disease: a systematic review of the literature. *J Manipulative Physiol Ther* 2005;28:429-42.
180. Santilli V, Beghi E, Finucci S. Chiropractic manipulation in the treatment of acute back pain and sciatica with disc protrusion: a randomized double-blind clinical trial of active and simulated spinal manipulations. *Spine J* 2006;6:131-7.
181. Modic MT, Steiberg PM, Ross JS, Masaryk TJ, Carter JR. Degenerative disk disease: assessment of changes in vertebral body marrow with MR imaging. *Radiology* 1988;166:193-9.
182. Orend acov a J, Cizkova D, Kafka J, Lukacova N, Marsala M, Sulla I, et al. Cauda equina syndrome. *Prog Neurobiol* 2001;64:613-37.
183. Benoist M. The natural history of lumbar degenerative spinal stenosis. *Joint Bone Spine* 2002;69:450-7.
184. Savage RA, Whitehouse GH, Roberts N. The relationship between the magnetic resonance imaging appearance of the lumbar spine and back pain, age and occupation in males. *Eur Spine J* 1997;6:106-14.
185. Waddel G, McIntosh A, Hutchinson A, Feder G, Lewis M. Low back pain evidence review. [Online publication]. London: Royal College of General Practitioners; 1999. Available from: <http://www.rcgp.org.uk>.
186. Fournier E, editor. *Atlas d' lectromyographie*. Paris:  dition M dicales Internationales; 2000.
187. Leis AA, Trapanive VC, editors. *Atlas of Electromyography*. New York: Oxford University Press; 2000. p. 199.
188. Jarvik JJ, Hollingworth W, Heagerty P, Haynor DR, Deyo RA. The Longitudinal Assessment of Imaging and Disability of the Back (LAIDBack) Study: baseline data. *Spine* 2001;26:1158-66.
189. Hollingworth W, Gray DT, Martin BI, Sullivan SD, Deyo RA, Jarvik JG. Rapid magnetic resonance imaging for diagnosing cancer-related low back pain. *J Gen Intern Med* 2003;18:303-12.
190. Frymoyer J. Point of view. *Spine* 2002;27:1926-33.
191. Nachemson A, Spitzer WO, LeBlanc FE, et al. Scientific approach to the assessment and management of activity-related spinal disorders. A monograph for clinicians: report of the Quebec Task Force on Spinal Disorders. *Spine* 1987;12 (Suppl 7):S1-S59.
192. Carroll LJ, Cassidy JD, Cote P. Depression as a risk factor for onset of an episode of troublesome neck and low back pain. *Pain* 2004;107:134-9.
193. Dionne CE, Boubonnais R, Fr emont P, Rossignol M, Stock SR, Laroque I. A clinical return-to-work rule for patients with back pain. *Can Med Assoc J* 2005;172:1559-67.
194. Maher C. Clinical prediction rule for return to work after back pain. *Can Med Assoc J* 2005;172:1575-6.
195. Waddell G, Burton AK, Main CJ. Screening to identify people at risk of long-term incapacity for work. London: Royal Society of Medicine Press Ltd; 2003.
196. Liang M, Komaroff AL. Roentgenograms in primary care patients with acute low back pain: a cost effectiveness analysis. *Arch Intern Med* 1982;142:1108-12.
197. Deyo RA, Diehl AK. Cancer as a cause of back pain: frequency, clinical presentation, and diagnostic strategies. *J Gen Intern Med* 1988;3:320-8.
198. Hass M, Nyiendo J, Peterson C, Thiel H, Seller T, Cassidy D, Yong-Hing K. Interrater reliability of roentgenological evaluation of the lumbar spine in lateral bending. *J Manipulative Physiol Ther* 1990;13:179-89.
199. Carragee EJ, Hannibal M. Diagnostic evaluation of low back pain. *Orthop Clin North Am* 2004;35:7-16.
200. Carragee EJ, Alamin TF, Miller JL, Carragee JM. Discographic, MRI and psychosocial determinants of low back pain disability and remission: a prospective study in subjects with benign persistent back pain. *Spine J* 2005;5:24-35.
201. Saal JS. General principles of diagnostic testing as related to painful lumbar spine disorders: a critical appraisal of current diagnostic techniques. *Spine* 2002;27:2538-45.
202. Tunks E. Neck pain and whiplash. In: Gray J, editor. *Therapeutic choices*. 4th ed. Ottawa, Ontario, Canada: Canadian Pharmacists Association; 2003. p. 583.
203. Simmons ED, Guyer RD, Graham-Smith A, Herzog R. Radiographic assessment for patients with low back pain. *Spine* 1995;20:1839-41.
204. Rudwaleit M, Metter A, Listing J, et al. Inflammatory back pain in ankylosing spondylitis: a reassessment of the clinical history for application as classification and diagnostic criteria. *Arthritis Rheum* 2006;54:569-78.
205. Zochling J, Braun J, van der Heijde D. Assessments in ankylosing spondylitis. *Best Pract Res Clin Rheumatol* 2006;20:521-37.
206. Naganathan V, Jones G, Nash P, et al. Vertebral fracture risk with long-term corticosteroid therapy. Prevalence and relation to age, bone density, and corticosteroid use. *Arch Intern Med* 2000;160:2917-22.
207. Friedrich M, Gittler G, Pieler-Bruha E. Misleading history of pain location in 51 patients with osteoporotic vertebral fractures. *Eur Spine J* 2006;15:1797-800.
208. Jarvik JG, Hollingworth W, Martin B, Emerson SS, Gray DT, Overman S, et al. Rapid magnetic resonance imaging vs radiographs for patients with low back pain: a randomized controlled trial. *JAMA* 2003;289:2810-8.
209. Kim PSY, Hsu W. Discitis in an adult following acupuncture treatment: a case report. *J Can Chiropr Assoc* 2004;48:132-6.

210. Otani K, Konno S, Kikuchi S. Lumbosacral transitional vertebrae and nerve-root symptoms. *J Bone Joint Surg Br* 2001;83:1137-40.
211. Joines JD, McNutt RA, Carey TS, Deyo RA, Rouhani R. Finding cancer in primary care outpatients with low back pain: a comparison of diagnostic strategies. *J Gen Intern Med* 2001; 16:14-23.
212. Sieper J, Rudwaleit M, Khan MA, Braun J. Concepts and epidemiology of spondyloarthritis. *Best Pract Res Clin Rheumatol* 2006;20:401-17.
213. Maksymowych WP, Landew e R. Imaging in ankylosing spondylitis. *Best Pract Res Clin Rheumatol* 2006;20: 507-19.
214. Scutellari PN, Rizzati R, Antinolfi G, Malfaccini F, Leprotti S, Campanati P. The value of computed tomography in the diagnosis of low back pain. A review of 2012 cases. *Minerva Med* 2005;96:41-59.
215. Turpin S, Lambert R. Role of scintigraphy in musculoskeletal and spinal infections. *Radiol Clin North Am* 2001;39:169-89.
216. Shapiro S. Medical realities of cauda equina syndrome secondary to lumbar disc herniation. *Spine* 2000;25:348-51.
217. Ahn UM, Ahn NU, Buchowski JM, Garrett ES, Sieber AN, Kostuik JP. Cauda equina syndrome secondary to lumbar disc herniation: a meta-analysis of surgical outcomes. *Spine* 2000; 25:515-22.
218. Kostuik JP, Harrington I, Alexander D, Rand W, Evans D. Cauda equina syndrome and lumbar disc herniation. *J Bone Joint Surg Am* 1986;68:386-91.
219. Marshall KG. La colonne lombo-sacr ee 4e partie: le syndrome de la queue de cheval. *MedActuel FMC* 2004;11-2.
220. Assendelft WJ, Bouter LM, Knipschild PG. Complications of spinal manipulation: a comprehensive review of the literature. *J Fam Pract* 1996;42:475-80.
221. Dupeyron A, Vautravers Ph, Lecoq J, Isner-Horobeti ME. Complications following vertebral manipulation—a survey of a French region physicians.  valuation de la fr quence des accidents li s aux manipulations vert brales   partir d'une enq ete r trospective r alis ee dans quatre d partements fran ais. *Ann Readapt Med Phys* 2003;46:33-40.
222. Haldeman S, Rubinstein SM. Cauda equina syndrome in patients undergoing manipulation of the lumbar spine. *Spine* 1992;17:1469-73.
223. Patijn J. Complications in manual medicine: a review of the literature. *J Manual Med* 1991;6:89-92.
224. Bhalla S, Menias CO, Heiken JP. CT of acute abdominal aortic disorders. *Radiol Clin North Am* 2003;41:1153-69.
225. Ayling J. Gut feeling. *Emerg Med Serv* 2003;32:37.
226. Crawford CM, Hurtgen-Grace K, Talarico E, Marley J. Abdominal aortic aneurysm: an illustrated narrative review. *J Manipulative Physiol Ther* 2003;26:184-95.
227. Upchurch GR, Schaub TA. Abdominal aortic aneurysm. *Am Fam Physician* 2006;73:1198-204.
228. Lynch RM. Accuracy of abdominal examination in the diagnosis of non-ruptured abdominal aortic aneurysm. *Accid Emerg Nurs* 2004;12:99-107.
229. Rowe LJ, Yochum TR. Masquaraders of musculoskeletal disease. In: Yochum TR, Rowe LJ, editors. *Essentials of skeletal radiology*. 3rd ed. Philadelphia: Lippincott Williams & Wilkins; 2005. p. 1818-24.
230. Mastracci TM, Cina CS, Canadian Society for Vascular Surgery. Screening for abdominal aortic aneurysm in Canada: review and position statement of the Canadian Society for Vascular Surgery. *J Vasc Surg* 2007;45:1268-76.
231. Powell JT, Brown LC. The natural history of abdominal aortic aneurysms and their risk of rupture. *Acta Chir Belg* 2001;101: 11-6.
232. Wanhainen A, Bergqvist D, Boman K, Nilsson TK, Rutegard J, Bjorck M. Risk factors associated with abdominal aortic aneurysm: a population-based study with historical and current data. *J Vasc Surg* 2005;41:390-6.
233. Fleming C, Whitlock EP, Beil TL, Lederle FA. Screening for abdominal aortic aneurysm: a best-evidence systematic review for the U.S. Preventive Services Task Force. *Ann Intern Med* 2005;142:203-11.
234. Powell JT. Smoking, lung function and the prognosis of abdominal aortic aneurysm. The UK small aneurysm trial participant. *Eur J Vasc Endovasc Surg* 2000;19:636-42.
235. Braunwald E, Fauci AS, Kasper DL, et al. *Harrison's principles of internal medicine*. 15th ed. New York: McGraw-Hill; 2001. p. 1431.
236. Kent KC, Zwolak RM, Jaff MR, Hollenbeck ST, Thompson RW, Schermerhorn ML, American Association of Vascular Surgery, et al, Society for Vascular Surgery, Society for Vascular Medicine and Biology. Screening for abdominal aortic aneurysm: a consensus statement. *J Vasc Surg* 2004;39:267-9.
237. Grimshaw G, Thompson JM. Changes in diameter of the abdominal aorta with age: an epidemiological study. *J Clin Ultrasound* 1997;25:7-13.
238. Ballard DJ, Fowkes FG, Powell JT. Surgery for small abdominal aortic aneurysms. *Cochrane Database Syst Rev* 2000:CD001835.
239. Henderson D, Chapman-Smith D, Miors S, Vernon H. Toronto, Ontario, Canada: Clinical guidelines for chiropractic practice in Canada. Proceedings of a consensus conference commissioned by the Canadian Chiropractic Association. *J Can Chiropr Assoc* 1994;38(Suppl):145.
240. Beck RW, Holt KR, Fox MA, Hurtegen-Grace K. Radiographic anomalies that may alter chiropractic intervention strategies found in a New Zealand population. *J Manipulative Physiol Ther* 2004;27:554-9.
241. Leo E, Biancari F, Nesi F, Pogany G, Bartolucci R, De Pasquale F, et al. Risk-scoring methods in predicting the immediate outcome after emergency open repair of ruptured abdominal aortic aneurysm. *Am J Surg* 2006;192: 19-23.
242. Cates JR. Abdominal aortic aneurysms: clinical diagnosis and management. *J Manipulative Physiol Ther* 1997;20: 557-61.
243. Sparks AR, Johnson PL, Meyer MC. Imaging of abdominal aortic aneurysms. *Am Fam Phys* 2002;65:1565-70.
244. Collins RD. *Differential diagnosis in primary care*. 3rd ed. Philadelphia: Lippincott Williams & Wilkins; 2003. p. 595.
245. Foerster V, Murtagh J, Lentle BC, Reed MH, Husereau D, Mensinkai S, editors. *CT and MRI for selected clinical disorders: A systematic review of clinical systematic reviews; 2005. [Monograph on the Internet] Technology report (59), CCOHTA*. Available from: <http://www.ccohta.ca>.
246. Scaringe JG, Ketner C. Manual method for the treatment of rib dysfunction and associated functional lesions. *Top Clin Chiro* 1999;6:20-38.
247. Erwin WM, Jackson PC, Homonko DA. Innervation of the human costovertebral joint: implications for clinical back pain syndromes. *J Manipulative Physiol Ther* 2000;23: 395-403.
248. Gregory PL, Biswas AC, Batt ME. Musculoskeletal problems of the chest wall in athletes. *Sports Med* 2002;32:235-50.
249. Jondeau G, Boileau C, Chevallier B, Delorme G, Digne F, Guiti C, et al. Marfan syndrome. *Arch Mal Coeur Vaiss* 2003; 96:1081-8.
250. Jondeau G, Muti C, Boileau C. Aortic aneurysms excluding Marfan's syndrome. *Arch Mal Coeur Vaiss* 2003;96: 1074-80.

251. Kaptoge S, Armbrecht G, Felsenberg D, Lunt M, O'Neill TW, Silman AJ, Reeve J, EPOS Study Group. When should the doctor order a spine X-ray? Identifying vertebral fractures for osteoporosis care: results from the European Prospective Osteoporosis Study (EPOS). *J Bone Miner Res* 2004;19:1982-93.
252. Angeli A, Guseppe G, Dovio A, Capelli G, de Feo D, Giannini S, et al. High prevalence of asymptomatic vertebral fractures in post-menopausal women receiving chronic glucocorticoid therapy: a cross-sectional outpatient study. *Bone* 2006;39:253-9.
253. Melton LJ, Kallmes DF. Epidemiology of vertebral fractures: implications for vertebral augmentation. *Acad Radiol* 2006;13:538-45.
254. Plan d'action contre l'ost eoporose: Une strat egie de lutte contre l'ost eoporose en Ontario. Rapport du Comit e du Plan d'action contre l'ost eoporose au minist ere de la Sant e et des Soins de longue dur ee. [Monograph on the Internet.] Pr epar e en partenariat avec la Soci ete de l'ost eoporose du Canada. 2003. [cited 2005 Jul 9]. Available from: <http://www.health.gov.on.ca>.
255. Nguyen HV, Ludwig S, Gelb D. Osteoporotic vertebral burst fractures with neurologic compromise. *J Spinal Disord Tech* 2003;16:10-9.
256. Hanley DA. Osteoporosis. In: Gray J, editor. *Therapeutic choices*. 4th ed. Ottawa: Canadian Pharmacists Association; 2003. p. 636.
257. Brown JP, Josse RG. Lignes directrices de pratique clinique 2002 pour le diagnostic et le traitement de l'ost eoporose au Canada. *CMAJ* 2003;168(6 Suppl):SF1-SF38.
258. Cappuccio FP, Meilahn E, Zmuda JM, et al. High blood pressure and bone mineral loss in elderly white women: a prospective study. *Lancet* 1999;354:971-5.
259. Yaffe K, Browner W, Cauley J, et al. Association between bone mineral density and cognitive decline in older women. *J Am Soc Geriatr Dent* 1999;47:1176-82.
260. D'Amelio P, Tamone C, Pluviano F, Di Stefano M, Isaia G. Effects of lifestyle and risk factors on bone mineral density in a cohort of Italian women: suggestion for a new decision rule. *Calcif Tissue Int* 2005;77:72-8.
261. Pafumi C, Ciotta L, Farina M, Bosco P, Chiarenza M, Pernicone G, Calcagno A, Evola F, Cavallaro A, Iemmola A, Cianci A. Evaluation of bone mass in young amenorrheic women with anorexia nervosa. *Minerva Ginecol* 2002;54:487-91.
262. Misra M, Aggarwal A, Miller KK, Almazan C, Worley M, Soyka LA, Herzog DB, Klibanski A. Effects of anorexia nervosa on clinical, hematologic, biochemical, and bone density parameters in community-dwelling adolescent girls. *Pediatrics* 2004;114:1574-83.
263. Cooper C, Westlake S, Harvey N, Javaid K, Dennison E, Hanson M. Review: developmental origins of osteoporotic fracture. *Osteoporos Int* 2006;17:337-47.
264. Cockayne S, Adamson J, Lanham-New S, Shearer MJ, Gilbody S, Torgerson DJ. Vitamin K and the prevention of fractures: systematic review and meta-analysis of randomized controlled trials. *Arch Intern Med* 2006;166:1256-61.
265. Goodman LR, Warren MP. The female athlete and menstrual function. *Curr Opin Obstet Gynecol* 2005;17:466-70.
266. Rud B, Jensen JEB, Mosekilde L, Nielsen SP, Hilden J, Abrahamsen B. Performance of four clinical screening tools to select peri- and early postmenopausal women for dual X-ray absorptiometry. *Osteoporos Int* 2005;16:764-72.
267. Cook RB, Collins D, Tucker J, Zioupos P. Comparison of questionnaire and quantitative ultrasound techniques as screening tools for DXA. *Osteoporos Int* 2005;16:1565-75.
268. Mauck KF, Cuddihy MT, Atkinson EJ, Melton LJ. Use of clinical prediction rules in detecting osteoporosis in a population-based sample of postmenopausal women. *Arch Intern Med* 2005;165:530-6.
269. Cadarette SM, McIsaac WJ, Hawker GA, Jakkimainen L, Culbert A, Zarifa G, Ola E, Jaglal SB. The validity of decision rules for selecting women with primary osteoporosis for bone mineral density testing. *Osteoporos Int* 2004;15:361-6.
270. Goulay ML, Miller WC, Richy F, Garrett JM, Hanson LC, Reginster JY. Performance of osteoporosis assessment tools in postmenopausal women aged 45-64 years. *Osteoporos Int* 2005;16:921-7.
271. Brenneman SK, Lacroix AZ, Bruist DS, Chen YT, Abbot TA. Evaluation of decision rules to identify postmenopausal women for intervention related osteoporosis. *Dis Manag* 2003;6:159-68.
272. Duan Y, Duboeuf F, Munoz F, Delmas PD, Seeman E. The fracture risk index and bone mineral density as predictors of vertebral structural failure. *Osteoporosis Int* 2006;17:54-60.
273. Sran MM, Khan KM. Is spinal mobilization safe in severe secondary osteoporosis? A case report. *Man Ther* 2006;11:344-51.
274. Lips P, Cooper C, Agnusdei D, Caulin F, Egger P, Johnell O, et al. Quality of life in patients with vertebral fractures: validation of the quality of Life Questionnaire of the European Foundation for Osteoporosis (QUALEFFO). Working Party for Quality of Life of the European Foundation for Osteoporosis. *Osteoporos Int* 1999;10:150-60.
275. Hashidate H, Kamimura M, Nakagawa H, Takahara K, Uchiyama S. Pseudoarthrosis of vertebral fracture: radiographic and characteristic clinical features and natural history. *J Orthop Sci* 2006;11:28-33.
276. Sran MM, Khan KM. Physiotherapy and osteoporosis: practice behaviour and clinicians' perceptions—a survey. *Man Ther* 2004;10:21-7.
277. Takahara K, Kamimura M, Nakagawa H, Hashidate H, Uchiyama S. Radiographic evaluation of vertebral fractures in osteoporotic patients. *J Clin Neurosci* 2007;14:122-6.
278. Osteoporosis overview; 2006 [Monograph on the Internet], National Institutes of Health Osteoporosis and Related Bone Disease-Resource Center; [cited 2006 May 15]. Available from: http://www.niams.nih.gov/Health_Info/Bone/Osteoporosis/default.asp.
279. Johnell O, Kanis J. Epidemiology of osteoporotic fractures. *Osteoporos Int* 2005;16:3-7.
280. Court-Brown CM, Caesar B. Epidemiology of adult fractures: a review. *Injury* 2006;37:691-7.
281. Sran MM, Kahn KM, Zhu Q, McKay HA, Oxland TR. Failure characteristics of the thoracic spine with a posteroanterior load: investigating the safety of spinal mobilization. *Spine* 2004;29:2382-8.
282. Haldeman S, Rubinstein SM. Compression fractures in patients undergoing spinal manual therapy. *J Manipulative Physiol Ther* 1992;15:450-4.
283. Shvartzman P, Abelson A. Complication of chiropractic treatment for back pain. *Postgrad Med* 1988; 83:57-8, 61.
284. Sran M, Khan KM, Keiver K, Chew JB, McKay HA, Oxland TR. Accuracy of DXA scanning of the thoracic spine: cadaveric studies comparing BMC, areal BMD and geometric estimates of volumetric BMD against ash weight and CT measures of bone volume. *Eur Spine J* 2005;14:971-6.
285. Maitland GD, Banks K, English K, Hengeveld E, editors. *Maitland's Vertebral Manipulation*. 6th ed. Boston: Butterworth Heinemann; 2001. p. 472.
286. Ernst E. Chiropractic spinal manipulation for (osteoporotic) back pain. *Br J Sports Med* 2003;37:195-6.
287. Haldeman S, Chapman-Smith D, Peterson DM. Guidelines for chiropractic quality assurance and practice parameters. Gaithersburg (MD): Aspen Publisher; 1992. p. 174.

288. Bougie JD. Geriatric practice-specific issues. Chap 54. In: Haldeman S, Dagenais S, et al, editors. *Principles and practice of chiropractic*. 3rd ed. New York: McGraw-Hill; 2005. p. 1223.
289. Bagger YZ, Tank o LB, Alexandersen P, Qui G, Christiansen C. Radiographic measure of aorta calcification is a site-specific predictor of bone loss and fracture at the hip (prospective epidemiological risk factors study group). *J Intern Med* 2006;259:598-605.
290. Schwab F, Dubey A, Gamez L, El-Fegoun AB, Hwang K, Pagala M, Farcy JP. Adult scoliosis: prevalence, SF-36, and nutritional parameters in an elderly volunteer population. *Spine* 2005;30:1082-5.
291. Weinstein SL, Dolan LA, Spratt KF, Peterson KK, Spoonamore MJ, Ponseti IV. Health and function of patients with untreated idiopathic scoliosis: a 50-year natural history study. *JAMA* 2003;289:559-67.
292. Aebi M. The adult scoliosis. *Eur Spine J* 2005;14:925-48.
293. Scoliosis Research Society [monograph on the Internet]. *Depth review of scoliosis*. 2000. Available from: <http://www.srs.org>.
294. Boulanger YL, Faucet R. Les d eviations du rachis. In: Dupuis M, Leclaire R, editors. *Pathologie m edicale de l'appareil locomoteur*. Qu ebec: Edisem; 1991. p. 309-40.
295. Karachalios T, Sofianos J, Roidis N, Sapkas G, Korres D, Nikolopoulos K. Ten-year follow-up evaluation of a school screening program for scoliosis. Is the forward-bending test an accurate diagnostic criterion for the screening of scoliosis? *Spine* 1999;24:2318-24.
296. Schwab F, Dubey A, Pagala M, Gamez L, Farcy JP. Adult scoliosis: a health assessment analysis by SF-36. *Spine* 2003;15:602-6.
297. Berven S, Deviren V, Demir-Deviren S, Hu SS, Bradford DS. Studies in the modified Scoliosis Research Society Outcome Instrument in adults: validation, reliability, and discriminatory capacity. *Spine* 2003;28:2164-9.
298. Murata Y, Takahashi K, Hanaoka E, Utsumi T, Yamagata M, Moriya H. Changes in scoliotic curvature angle during early phase of degenerative lumbar scoliosis. *Spine* 2002;27:2268-73.
299. Levy AR, Golberg MS, Mayo NE, Hanley JA, Poitras B. Reducing the life time risk of cancer from spinal radiographs among people with adolescent idiopathic scoliosis. *Spine* 1996;21:1540-8.
300. Morin DM, Lonstein JE, Stovall M, Hacker DG, Luckyanov N, Land CE. Breast cancer mortality after diagnostic radiography: findings from the U.S. Scoliosis Cohort Study. *Spine* 2000;25:2052-63.
301. Schwab F, Smith VA, Biserni M, Gamez L, Farcy JP, Pagala M. Adult scoliosis: a quantitative radiographic and clinical analysis. *Spine* 2002;27:387-92.
302. Glassman SD, Berven S, Bridwell K, Horton W, Dimar JR. Correlation of radiographic parameters and clinical symptoms in adult scoliosis. *Spine* 2005;30:682-8.
303. Toyone T, Tanaka T, Kato D, Kaneyama R, Otsuka M. Anatomic changes in lateral spondylolisthesis associated with adult lumbar scoliosis. *Spine* 2005;30:E671-5.
304. Schwab F, el-Fegoun AB, Gamez L, Goodman H, Farcy JP. A lumbar classification of scoliosis in the adult patient: preliminary approach. *Spine* 2005;15:1670-3.
305. Danielsson AJ, Nachemson AL. Radiographic findings and curves progression 22 years after treatment for adolescent idiopathic scoliosis: comparison of brace and surgical treatment with matching control group of straight individuals. *Spine* 2001;26:516-25.
306. Cote P, Cassidy JD, Carroll L. The factors associated with neck pain and its related disability in the Saskatchewan population. *Spine* 2000;25:1109-17.
307. Honet JC, Ellenberg MR. What you always wanted to know about the history and physical examination of neck pain but were always afraid to ask. *Phys Med Rehabil Clin N Am* 2003;14:473-91.
308. Triano JJ, Kawchuk G, Gudavalli MR, Haneline MT, Hill MD, Wynd S, editors. *Current concepts spinal manipulation and cervical arterial incidents*. Clive, Iowa: NCMIC Insurance Company; 2006.
309. C  t e P, Kreitz B, Cassidy D, Thiel H. The validity of the extension-rotation test as a screening procedure before neck manipulation: a secondary analysis. *J Manipulative Physiol Ther* 1996;19:159-64.
310. Licht P, Christensen HW,  ilund-Carlsen HPF. Is there a role for premanipulative testing before cervical manipulation? *J Manipulative Physiol Ther* 2000;23:175-9.
311. Mann T, Refshauge M. Causes of complications from cervical manipulation. *Aust J Physiother* 2001;47:255-66.
312. Zeina C, Grant R, Johnson C, Dansie B, Taylor J, Spyropolous P. The effect of cervical rotation on blood flow in the contralateral vertebral artery. *Man Ther* 2003;8:103-9.
313. Westaway MD, Stratford P, Symons B. False-negative extension/rotation pre-manipulative screening test on a patient with an atretic and hypoplastic vertebral artery. *Man Ther* 2003;8:120-7.
314. Thiel H, Rix G. Is it time to stop functional pre-manipulation testing of the cervical spine? *Man Ther* 2005;10:154-8.
315. M akel a M, Heliovaara M, Sievers K, Impivaara O, Knekt P, Aromaa A. Prevalence, determinants, and consequences of chronic neck pain in Finland. *Am J Epidemiol* 1991;134:1356-67.
316. C  t e P, Cassidy JD, Carroll L. The Saskatchewan Health and Back pain survey—The prevalence of neck pain and related disability in Saskatchewan adults. *Spine* 1998;23:1689-98.
317. Cote P, Cassidy JD, Carroll LJ, Kristman V. The annual incidence and course of neck pain in the general population: a population-based cohort study. *Pain* 2004;112:267-73.
318. The Canadian Chiropractic Association and the Canadian Federation of Chiropractic Regulatory Boards, Clinical Practice Guidelines Development Initiative, Guidelines Development Committee (GDC) comprising, Anderson-Peacock E, Blouin R, Bryans N, Danis A, et al. *Chiropractic Practice Guidelines for Nontraumatic Acute Neck Pain in Adults*. *J Can Chiropr Assoc* 2005;49:158-209.
319. Gay RE. The curvature of the cervical spine: Variations and significance—Reviews of the literature. *J Manipulative Physiol Ther* 1993;16:591-4.
320. Cote P, Cassidy JD, Yong-Hing K, Sibley J, Loewy J. Apophysial joint degeneration, disc, degeneration, and sagittal curve of the cervical spine. *Spine* 1997;22:859-64.
321. McAviney J, Schulz D, Bock R, Harrison DE, Holland B. Determining the relationship between cervical lordosis and neck complaints. *J Manipulative Physiol Ther* 2005;28:187-93.
322. Grob D, Frauenfelder H, Mannion AF. The association between cervical spine curvature and neck pain. *Eur Spine J* 2007;16:669-78.
323. Harrison DD, Harrison DE, Janik TJ, Cailliet R, Ferrantelli JR, Haas JW, et al. Modeling of the sagittal cervical spine as a method to discriminate hypolordosis: results of elliptical and circular modeling in 72 asymptomatic subjects, 52 acute neck pain subjects, and 70 chronic neck pain subjects. *Spine* 2004;29:2485-92.
324. Dreyer SJ, Boden SD. Nonoperative treatment of neck and arm pain. *Spine* 1998;23:2746-54.
325. Kleinrensink GJ, Stoeckart R, Mulder PGH, Hoek GVD, Broek T, Vleeming A, Snijders CJ. Upper limb tension tests as

- tools in the diagnosis of nerve and plexus lesions anatomical and biomechanical aspects. *Clin Biomech* 2000;15:9-14.
326. Uitylught G, Idenbaun S. Clinical assessment of atlantoaxial instability using the Sharp-Purser test. *Arthritis Rheum* 1988; 84:370-4.
327. Uchihara T, Furukawa T, Tsukagoshi H. Compression of brachial plexus as a diagnostic test of cervical cord lesion. *Spine* 1994;19:2170-3.
328. Viikari-Juntura E, Prsa M, Laasonen EM. Validity of clinical tests in the diagnosis of root compression in cervical disc diseases. *Spine* 1989;14:253-7.
329. Viikari-Juntura E, Takala E, Riihim aki H, Martikainen R, J ppinen P. Predictive validity of signs in the neck and shoulders. *J Clin Epidemiol* 2000;53:800-8.
330. Radhakrishnan K, Litchy WJ, O'Fallon M, Kurlan LT. Epidemiology of cervical radiculopathy. A population study from Rochester, Minnesota, 1976 through 1990. *Brain* 1994; 117:325-35.
331. Malanga GA, Landes P, Nadler SF. Provocative tests in cervical spine examination: historical basis and scientific analyses. *Pain Physician* 2003;6:199-205.
332. Wainner RS, Fritz JM, Irrgang JJ, Boringer ML, Delitto A, Allison S. Reliability and diagnostic accuracy of the clinical examination and patient self-measures for cervical radiculopathy. *Spine* 2003;28:52-62.
333. Rubinstein SM, Pool JJ, van Tulder MW, Riphagen II, de Vet HC. A systematic review of the diagnostic accuracy of provocative tests of the neck for diagnosing cervical radiculopathy. *Eur Spine J* 2007;16:307-19.
334. Fehling MG, Skaf G. A review of the pathophysiology of cervical spondylotic myelopathy with insights for potential novel mechanisms drawn from traumatic spinal injury. *Spine* 1998;23:2730-7.
335. Duus P. Innervation musculaire segmentaire et p eriph erique. *Diagnosis neurologique—Les bases anatomiques*. 6th ed. Bruxelles: De Boeck Universit ; 1998. p. 85.
336. Debois V, Herz R, Berghmans D, Hermans B, Herregots P. Soft cervical disc herniation. Influence of cervical canal measurements on development of neurologic symptoms. *Spine* 1999;24:1996-2002.
337. Ellenberg M, Honet JC, Treanor WJ. Cervical radiculopathy—Review article. *Arch Phys Med Rehabil* 1994;75:342-52.
338. Dougherty P, Bajwa S, Burke J, Dishman JD. Spinal manipulation postepidural injection for lumbar and cervical radiculopathy: a retrospective case series. *J Manipulative Physiol Ther* 2004;27:449-56.
339. Oppenheim JS, Spitzer DE, Segal DH. Nonvascular complications following spinal manipulation. *Spine J* 2005;5: 660-6.
340. Murphy DR, Hurwitz EL, Gregory A, Clary R. A nonsurgical approach to the management of patients with cervical radiculopathy: a prospective observational cohort study. *J Manipulative Physiol Ther* 2006;29:279-87.
341. Souza TA. Differential diagnosis and management for the chiropractor—Protocols and algorithms. 2nd ed. Gaithersburg, Maryland: Aspen Publication; 2001. p. 43.
342. Mercado AC, Carroll LJ, Cassidy JD, Cote P. Passive coping is a risk factor for disabling neck or low back pain. *Pain* 2005; 11:51-7.
343. van der Donk J, Schouten JS, Passchier J, Romunde LK, Valkenburg HS. The association of neck pain with radiological abnormalities of the cervical spine and personality traits in a general population. *J Rheumatol* 1991;18:1884-9.
344. Marchiori DM, Henderson CN. A cross-sectional study correlating cervical radiographic degenerative findings to pain and disability. *Spine* 1996;21:2747-52.
345. Gore DR, Sepic SB, Gardner GM, Murray P. Neck pain: A long-term follow-up of 205 patients. *Spine* 1987;12:1-5.
346. Gore DR. Roentgenographic findings in the cervical spine in asymptomatic persons. A ten year follow-up. *Spine* 2001;26: 2463-6.
347. Peterson C, Bolton J, Wood AR, Humphreys BK. A cross-sectional study correlating degeneration of the cervical spine with disability and pain in United Kingdom patients. *Spine* 2003;28:129-33.
348. Kauppi M, Neva MH. Sensitivity of lateral view cervical spine radiographs taken in the neutral position in atlantoaxial subluxation in rheumatic diseases. *Clin Rheumatol* 1998;17: 511-4.
349. Swinkels RAHM, Oostendorp RAB. Upper cervical instability: facts or fiction. *J Manipulative Physiol Ther* 1996;19:185-94.
350. Boden SD, McCowin PR, Davis DO, Dina TS, Mark AS, Wiesel S. Abnormal magnetic resonance scans of the cervical spine in asymptomatic subjects. *J Bone Joint Surg (Am)* 1990; 72:1178-84.
351. Lehto IJ, Tertti MO, Komu ME, Paajanen HE, Tuominen J, Kormanen MJ. Age-related MRI changes at 0.1 T in cervical discs in asymptomatic subjects. *Neuroradiology* 1994;36: 49-52.
352. Matsumoto M, Fujimura Y, Suzuki N, Nishi Y, Nakamura M, Yabe Y, et al. MRI of cervical discs in asymptomatic subjects. *J Bone Joint Surg* 1998;80-B:19-24.
353. Dvor ak J. Epidemiology, physical examination and neurodiagnostics. *Spine* 1998;23:2663-73.
354. Haldeman S, Kholbeck FJ, McGregor M. Stroke, cerebral artery dissection, and cervical spine manipulative therapy. *J Neurol* 2002;249:1098-104.
355. Abdu WA, Provencher ML. Primary bone and metastatic tumors of the cervical spine. *Spine* 1998;23:2767-77.
356. Shah AN, Pietrobon R, Richardson WJ, Myers BS. Patterns of tumor spread and risk of fracture and epidural impingement in metastatic vertebrae. *J Spinal Disord Tech* 2003;16: 83-9.
357. Melsom RD, Benjamin JC, Barnes CG. Spontaneous atlantoaxial subluxation: an unusual presenting manifestation of Reiter's syndrome. *Ann Rheum Dis* 1989;48:170-2.
358. Ramos-Remus C, Gomez-Vargas A, Gizman-Guzman JL, Jimenez-Gil F, Gamez-Nava JJ, et al. Frequency of atlantoaxial subluxation and neurologic involvement in patients with ankylosing spondylitis. *J Rheumatol* 1995;22:2120-5.
359. Rogers LF. The congenital malformation syndromes: osteochondrodysplasias, dysostoses, and chromosomal disorders. In: Jhul JH, Crummy AB, editors. *Essential of radiologic imaging*. 5th ed. Philadelphia: J.B. Lippincott Company; 1987. p. 314.
360. Toussiroit E, Benansour A, Bonneville JF. Atlantoaxial subluxation in an ankylosing spondylitis patient with cervical spine ossification. *Br J Rheumatol* 1997;36:293-5.
361. Taybi H. *Handbook of syndromes and metabolic disorders. Radiologic and clinical manifestations*. Mosby, St. Louis (Mo). 1998. p. 46, 58, 174.
362. Harger BL, Hoffman LE. Diagnostic imaging of the cervical spine. In: Murphy DR, editor. *Conservative management of cervical spine syndromes*. New York: McGraw-Hill Company; 2000. p. 376.
363. Guilles JT, Sherk HH. Congenital osseous anomaly of the upper and lower cervical spine in children—Current concepts review. *J Bone Joint Surg Am* 2002;84-A:277-88.
364. Reiter MF, Boden S. Inflammatory disorders of the cervical spine. *Spine* 1998;23:2755-66.
365. Pool JJ, Hoving JL, de Vet HC, van Mameren H, Bouter LM. The interexaminer reproducibility of physical examination of

- the cervical spine. *J Manipulative Physiol Ther* 2004;27:84-90.
366. Cattrysse E, Swinkels RAHM, Oostendorp RAB, Duquet W. Upper cervical instability: are clinical tests reliable? *Man Ther* 1997;2:91-7.
 367. Riise T, Jacobsen BK, Gran JT. High mortality in patients with rheumatoid arthritis and atlantoaxial subluxation. *J Rheumatol* 2001;28:2425-9.
 368. Baddoura R, Haddad S, Awada H, Al-Masri AF, Merheb G, Attoui S, Okais J, Messayke J, Ghandour F. Severity of rheumatoid arthritis: the SEVERA study. *Clin Rheumatol* 2005;15:1-5.
 369. Wolfs JF, Peul WC, Boers M, van Tulder MW, Brand R, van Houwelingen HJ, Thomeer RT. Rationale and design of The Delphi Trial-I (RCT) 2: international randomized clinical trial of rheumatoid craniocervical treatment, an intervention-prognostic trial comparing 'early' surgery with conservative treatment [ISRCTN65076841]. *BMC Musculoskelet Disord* 2006;7:14.
 370. Zatarain E, Strand V. Monitoring disease activity of rheumatoid arthritis in clinical practice: contributions from clinical trials. *Nat Clin Pract Rheumatol* 2006;2:611-8.
 371. Naranjo A, Carmona L, Gavrilu D, Balsa A, Belmonte MA, Tena X, Rodriguez-Lozano C, Sanmarti R, Gonzalez-Alvaro I. EMECAR Study Group. Prevalence and associated factors of anterior atlantoaxial luxation in a nation-wide sample of rheumatoid arthritis patients. *Clin Exp Rheumatol* 2004;22:427-32.
 372. Bouchaud-Chabot A, Liote F. Cervical spine involvement in rheumatoid arthritis. A review. *Joint Bone Spine* 2002;69:141-54.
 373. Matsumoto M, Fujimura Y, Toyama Y. Usefulness and reliability of neurological signs for level diagnosis in cervical spine myelopathy caused by soft cervical disc herniation. *J Spinal Disord* 1996;9:317-21.
 374. Matsumoto M, Ishikawa M, Ishii K, Nishizawa T, Maruiwa H, Nakamura M, et al. Usefulness of neurological examination for diagnosis of the affected level in patients with cervical compressive myelopathy: prospective comparative study with radiological evaluation. *J Neurosurg Spine* 2005;2:535-9.
 375. Harrop JS, Hanna A, Silva MT, Sharan A. Neurological manifestations of cervical spondylosis: an overview of signs, symptoms, and pathophysiology. *Neurosurgery* 2007;60 (1 Supp11):S14-20.
 376. Kadanka Z, Mares M, Bednarik J, Smrcka V, Krbec M, Chaloupka R, Dusek L. Predictive factors for spondylotic cervical myelopathy treated conservatively or surgically. *Eur J Neurol* 2005;12:55-63.
 377. Browder DA, Erhard RE, Piva SR. Intermittent cervical traction and thoracic manipulation for management of mild cervical compressive myelopathy attributed to cervical herniated disc: a case series. *J Orthop Sports Phys Ther* 2004;34:701-12.
 378. Kukurin GW. The amelioration of symptoms in cervical spinal stenosis with spinal cord deformation through specific chiropractic manipulation: a case report with long-term follow-up. *J Manipulative Physiol Ther* 2004;27:e7.
 379. Andrews FJ. Transient neuropraxia associated with cervical spine stenosis. *Emerg Med J* 2002;19:172-3.
 380. Gilles LGF. Mechanisms of neurovascular compression within the spinal and intervertebral canals. *J Manipulative Physiol Ther* 2000;23:1007-11.
 381. Merli J, Staas E. Acute transverse myelopathy: association with body position. *Arch Phys Med Rehabil* 1985;66:325-8.
 382. Milkawa Y, Watanabe R, Fuse K. Quadraplegia caused by push-up exercises. *Arch Orthop Trauma Surg* 1994;113:174-5.
 383. Patten J. *Neurological differential diagnosis*. 2nd ed. London: Springer-Verlag; 1996. p. 258.
 384. Vertichel P, Berthelot JL, Randriananja H, Crozier S, Masson C. Traumatic spinal complications of cervical arthrosis. *Presse Med* 1996;17:230-4.
 385. Pake WW, Whalen JL. Phrenic paresis—A possible additional spinal cord dysfunction induced by neck manipulation in cervical spondylotic myelopathy (CSM): A report of two cases with anatomical and clinical considerations. *Clin Anat* 2001;14:173-8.
 386. An HS. Cervical spine trauma. *Spine* 1998;23:2713-29.
 387. Terret AGJ, editor. *Current concepts in Vertebrobasilar complications following spinal manipulation*. 2nd ed. Iowa: NCMIC Group Inc; 2001. p. 35-53.
 388. Terret AJ. Vertebrobasilar stroke following spinal manipulation therapy. In: Murphy D, editor. *Conservative management of cervical spine syndromes*. New York: McGraw-Hill; 2000. p. 562.
 389. Haldeman S, Kohlbeck FJ, McGregor M. Risk factors and precipitating neck movements causing vertebrobasilar artery dissection after cervical trauma and spinal manipulation. *Spine* 1999;24:785-94.
 390. Rubinstein SM, Peerdeman SM, van Tulder MW, Riphagen I, Haldeman S. A systematic review of the risk factors for cervical artery dissection. *Stroke* 2005;36:1575-80.
 391. Rothwell DM, Bondy S, Williams JI. Chiropractic manipulation and stroke. A population-based case-control study. *Stroke* 2001;32:1054-60.
 392. Smith WS, Johnson SC, Skalabrin EJ, Weaver M, Azari P, Albers GW, Gress DR. Spinal manipulative therapy is an independent risk factor for vertebral artery dissection. *Neurology* 2003;60:1424-8.
 393. Calvet D, Boutouysie P, Touze E, Laloux B, Mass JL, Laurent S. Increase stiffness of the carotid wall material in patients with spontaneous cervical artery dissection. *Stroke* 2004;35:2078-82.
 394. Grau AJ, Brandt T, Forsting M, Winter R, Hacke W. Infection-associated cervical artery dissection. *Stroke* 1997;28:453-5.
 395. Pezzi A, Del Zotto E, Padovani A. Hyperhomocysteinemia. A potential risk factor for cervical artery dissection following chiropractic manipulation of the cervical spine. *J Neurol* 2002;249:1401-3.
 396. Allui RV, Mohan V, Komandur S, Chawda K, Chaudhuri JR, Husan Q. MTHFR C677T gene mutation as a risk factor for arterial stroke: a hospital based study. *Eur J Neurol* 2005;12:40-4.
 397. Bos MJ, Van Goor ML, Koudsataal JP, Dippel DW. Plasma homocysteine is a factor for recurrent vascular events in young patients with an ischaemic stroke or TIA. *J Neurol* 2005;1-6.
 398. Cacas JP, Bautista LE, Smeeth L, Sharma P, Hingorani AD. Homocysteine and stroke: evidence on a causal link from Mendelian randomisation. <http://www.thelancet.com> 2005; 365:224-232.
 399. Hankey GJ, Eikelboom JW. Homocysteine and stroke. <http://www.thelancet.com> 2005; 365:194-196.
 400. Rubinstein SM, Haldeman S, van Tulder MW. An etiologic model to help explain the pathogenesis of cervical artery dissection: implications for cervical manipulation. *J Manipulative Physiol Ther* 2006;29:336-8.
 401. Rosner AL. Chiropractic manipulation and stroke. *Letters to the Editor. Stroke*, vol. 32; 2001. p. 2207-8.

402. Haneline MT, Lewkovich G. Identification of internal carotid artery dissection in chiropractic practice. *J Can Chiropr Assoc* 2004;48:206-10.
403. Acar M, Degirmenci B, Yucel A, Albayrak R, Hatanir A. An evaluation of internal carotid artery and cerebral blood flow volume using color duplex sonography in patients with vertebral artery hypoplasia. *Eur J Radiol* 2005;53:450-3.
404. Jeng JS, Yip PK. Evaluation of vertebral artery hypoplasia and asymmetry by color-coded duplex ultrasonography. *Ultrasound Med Biol* 2004;30:605-9.
405. Saito K, Kimura K, Nagatsuka K, Nagano K, Minematsu K, Ueno S, Naritomi H. Vertebral artery occlusion in duplex color-coded ultrasonography. *Stroke* 2004;35:1068-72.
406. Kizikilic O, Hurcan C, Mihmanli I, Oguzkurt L, Yildirim T, Tercan F. Color Doppler analysis of vertebral arteries: corrective study with angiographic data. *J Ultrasound Med* 2004;23:1483-91.
407. Johnson C, Grant R, Dansie B, Taylor J, Spyropolous P. Measurement of blood flow in the vertebral artery using colour duplex Doppler ultrasound: establishment of the reliability of selected parameters. *Man Ther* 2000;5:21-9.
408. Rosner AL. Spontaneous cervical artery dissections: another perspective. *J Manipulative Physiol Ther* 2004;27:124-32.
409. Mullins ME. Modern emergent stroke imaging: pearls, protocols, and pitfalls. *Radiol Clin N Am* 2006;44:41-62.
410. Str ek P, Rero n E, Maga P, Modrzejewski M, Szybist N. A possible correlation between vertebral artery insufficiency and degenerative changes in the cervical spine. *Eur Arch Otorhinolaryngol* 1998;255:437-40.
411. Gulsun M, Saatci I, Akata D, Ozman MN, Sennroglu L, Yucel T. Radiologic investigation of vertebral insufficiency and quantification of vertebrobasilar flow with magnetic resonance imaging. *Tani Grisim Radyol* 2003;9:279-86.
412. Mitchell J. The vertebral artery: a review of anatomical, histopathological and functional factors influencing blood flow to the hindbrain. *Physiother Theory Pract* 2005;21:23-36.
413. Symons BP, Leonard T, Herzog W. Internal forces sustained by the vertebral artery during spinal manipulative therapy. *J Manipulative Physiol Ther* 2002;25:504-10.
414. Wingerchuk DM, Spencer B, Dodick DW, Demaerschalk BM. Migraine with aura is a risk factor for cardiovascular and cerebrovascular disease: a critically appraised topic. *Neurologist* 2007;13:231-3.
415. Vaccaro AR, Klein GR, Flanders AE, Albert TJ, Balderston RA, Cotler JM. Long-term evaluation of vertebral artery injuries following cervical spine trauma using magnetic resonance angiography. *Spine* 1998;23:789-94.
416. Friedman D, Flanders A, Thomas C, Millar W. Vertebral artery injury after acute cervical spine trauma: rate of occurrence as detected by MR angiography and assessment of clinical consequences. *AJR Am J Roentgenol* 1995;164:443-7.
417. Giacobetti FB, Vaccaro AR, Bos-Giacobetti MA, Deeley DM, Albert TJ, Farmer JC, Cotler JM. Vertebral artery occlusion associated with cervical spine trauma. A prospective analysis. *Spine* 1997;15:188-92.
418. Veras LM, Pedraza-Gutierrez S, Castellanos J, Capellades J, Casamitjana J, Rovira-Canellas A. Vertebral artery occlusion after acute cervical spine trauma. *Spine* 2000;25:1171-7.
419. Schellinger PD, Schwab S, Krieger D, Fiebach JB, Steiner T, Hund EF, Hacke W, Meinck HM. Masking of vertebral artery dissection by severe trauma to the cervical spine. *Spine* 2001;26:314-9.
420. Sim E, Vaccaro AR, Berzlanovich A, Pienaar S. The effects of staged cervical flexion–distraction deformities of the patency of the vertebral arterial vasculature. *Spine* 2000;25:2180-6.
421. Taneichi H, Suda K, Kajino T, Kaneda K. Traumatically induced vertebral artery occlusion associated with cervical spine injuries: prospective study using magnetic resonance angiography. *Spine* 2005;30:1955-62.
422. Beaudry M, Spence JD. Motor vehicle accidents: the most common cause of traumatic vertebrobasilar ischemia. *Can J Neurol Sci* 2003;30:320-5.

APPENDIX A. LIST OF ABBREVIATIONS AND GLOSSARY FOR SPINE DISORDERS

AAA:	Abdominal aortic aneurysms
AAS:	Atlantoaxial subluxation
ACR:	American College of Radiology
ALP:	Alkaline phosphatase (if elevated, R/O cancer, infection Paget’s disease)
AP:	Anteroposterior
APOM:	Anteroposterior open mouth
AS:	Ankylosing spondylitis
BMI:	Body mass index
CES:	Cauda equina syndrome
CNS:	Central nervous system
CAD:	Cervical artery dissection
CI:	Confidence interval
CT:	Computed tomography (advanced imaging modality that uses ionizing radiation [Table 1]. Generally considered to be the gold standard imaging for the chest, abdomen, pelvis, complex osseous anatomy, or in musculoskeletal disorders when contraindications for MRI exist).

CTA:	Computed tomography angiography
DDx:	Differential diagnosis
Dx:	Diagnosis
ESR:	Erythrocyte sedimentation rate (if elevated, R/O inflammation due to cancer, infection, fracture)
GHQ-28:	General Health Questionnaire
GPP:	Good Practice Point
HNP:	Herniated nucleus pulposus/disc herniation
Hx:	History
IVF:	Intervertebral foramen
JRA:	Juvenile rheumatoid arthritis
LBP:	Low back pain
LDH:	Lumbar disc herniation
LOE:	Level of evidence
MC:	More common
Mo:	Month
MRA:	Magnetic resonance angiography—gadolinium enhanced
MRI:	Magnetic resonance imaging (generally considered to be the best single advanced imaging modality of the spine to confirm or exclude most clinically relevant findings after conventional radiographs)
NM:	Nuclear medicine (also known as bone scan or scintigraphy) uses a radiopharmaceutical and gamma camera to produce a physiologic skeletal survey (Table 1). Has high sensitivity but low specificity.
NDI:	Neck Disability Index
PA:	Posteroanterior
PSA:	Prostate-specific antigen (R/O prostate cancer)
Red Flag:	Indicator of potentially serious pathology obtained from history, physical examination, and/or radiographs. Nonspecific/uncomplicated/mechanical neck and back pain are considered diagnoses of exclusion. Patients should therefore be properly evaluated for indicators of potentially serious pathology before undergoing some forms of manual therapy.
RA:	Rheumatoid arthritis
Rh factor:	Rheumatoid factor (indicates possible rheumatoid arthritis; 2%-5% false positive)
ROM:	Range of motion
R/O:	Rule out
SLR:	Straight leg raise test
SMT:	Spinal manipulative therapy—high-velocity, low-amplitude thrust techniques
S&S:	Signs and symptoms
TIA:	Transient ischemic attack
Tx:	Treatment
US:	Ultrasound
VAD:	Vertebral artery dissection
VBA:	Vertebrobasilar accident
VBI:	Vertebrobasilar insufficiency
WAD:	whiplash associated disorder
W/O:	Without
X-ray:	Conventional radiograph
yoa:	Years of age
<:	Less than
>:	Greater than
≥:	Equal to or greater than
Ψ:	psychology/psychiatry

APPENDIX B. SUMMARY OF RECOMMENDATIONS

Table 1. Summary of Recommendations—Thoracolumbar, Lumbar, and Thoracic Spine Trauma

Patient Presentation	Recommendations
<p>Adult patient with recent (<2 wk) acute thoracolumbar, lumbar, or thoracic spine trauma Absence of pain, normal ROM, and absence of neurologic deficits</p>	<p>Radiographs not routinely indicated [C]</p>
<p>Adult patient with thoracolumbar, lumbar or thoracic spine blunt trauma or acute injuries (falls, MVAs, motorcycle, pedestrian, cyclists, etc)</p> <p>High-risk screening criteria for spinal injuries include any of the following:</p> <ol style="list-style-type: none"> 1. Back pain 2. Midline tenderness on palpation 3. Distracting painful injury and other high-risk mechanism of injury* 4. Neurologic deficits 5. Altered consciousness (caused by head trauma, intoxication/ ethanol, or drugs) 	<p>Radiographs indicated [B] Lumbar <i>AND</i> thoracic spine: AP, lateral views</p> <p>Special investigations [C]</p> <ul style="list-style-type: none"> • CT scan (multidetector [multislice], spiral CT) • MRI
<p>Adult patient with posttraumatic chest wall pain Minor trauma</p>	<p>Radiographs not routinely indicated [D]</p>
<p>Major trauma</p>	<p>Radiographs indicated [GPP] PA, lateral chest radiographs, Specific rib radiographs (AP, oblique)</p> <p>Additional views: PA chest in full expiration, Thoracic and /or lumbar spine views</p> <p>Special investigations [GPP]</p> <ul style="list-style-type: none"> • CT for sternum injury, pulmonary, pleural, and osseous abnormalities
<p>Adult patient with pelvis and sacrum trauma (including falls with inability to bear weight)</p>	<p>Radiographs indicated [D] AP Pelvis and lateral hip “frog leg”</p> <p>Additional views: lateral lumbar view, Angulated AP sacrum view (15-45° cephalad)</p> <p>Special investigations [D]</p> <ul style="list-style-type: none"> • NM, MRI or CT may be helpful if radiographs are normal or equivocal.
<p>Coccyx trauma and coccydynia Consider views of the sacrum if distal <i>sacrum fracture is suspected</i></p>	<p>Radiographs not routinely indicated: (spot AP, lateral coccyx) [C]</p> <p>Additional views: AP, lateral sacrum, Dynamic sitting lateral views of the coccyx</p>

Table 2. Summary of Recommendations—Cervical Spine Trauma

Patient Presentation	Recommendations
Adult patient with acute neck injury and <i>negative</i> CCSR (Canadian Cervical Spine Rule for Radiography in Alert and Stable Trauma Patients)	Radiographs not routinely indicated [B]
Adult patient with acute neck injury and <i>positive</i> CCSR (Canadian Cervical Spine Rule for Radiography in Alert and Stable Trauma Patients)	Radiographs indicated [B] APOM, AP lower cervical, neutral lateral
Conventional radiographs recommended in the presence of any of the <i>Canadian Cervical Spine Rule</i> criteria are fulfilled: (A) High-risk factors in alert and stable patient? 1. Age >65 2. Dangerous mechanisms of injury* 3. Paresthesias in extremities (B) Low-risk factors that allow ROM assessment? 1. Simple rear end collision** 2. Patient seated in the waiting room 3. Ambulatory at one time since trauma 4. Delayed cervical pain onset 5. Absence of midline cervical tenderness (C) ROM assessment: is patient able to <i>actively</i> turn his/her head to 45° in both directions?	If fracture is suspected: 3 views + CT scan recommended Additional views: CT now replaces oblique, pillar, dynamic flexion/extension (F/E) in suspected fracture [GPP] Special investigations [C] • CT, MRI

Table 3. Summary of Recommendations—Adult Nontraumatic Lumbar Spine Disorders

Patient Presentation	Recommendations
Adult patient with acute uncomplicated* LBP (<4 wks' duration)	Radiographs not initially indicated [B]
* <i>Uncomplicated definition:</i> nontraumatic LBP without neurologic deficits or indicators of potentially serious pathologies—(see red flag list for details). For most young or middle-aged adults, early diagnostic evaluation of low back complaints may focus on 3 basic questions: diagnostic imaging is infrequently required (Jarvik 2002). ⁴ 1. Is there underlying systemic disease? 2. Is there neurologic impairment that might require surgical intervention? 3. Is social or psychological distress amplifying or prolonging the pain?	Special investigations not indicated [B]
Adult patient with uncomplicated subacute (4-12 wks' duration) or persistent LBP (>12 wks' duration) AND no previous treatment trial. A trial of up to 4-6 wk of conservative care is appropriate before radiographs	Radiographs not initially indicated [B]
Adult patient with nontraumatic acute LBP AND sciatica (no red flags) The first clinical clue to neurologic impairment usually is a history of sciatica: sharp pain radiating down the posterior or lateral aspect of the leg, often associated with numbness or paresthesia.	Radiographs not initially indicated [B]

Table 3 (continued)

Patient Presentation	Recommendations
Specific Clinical Diagnoses:	
Common causes of sciatica	
<p>(A) Suspected LDH:</p> <ul style="list-style-type: none"> • Risk factors for LDH include: men (1.6 times more likely), middle age (35-54 y), repetitive/heavy lifting, current smoking, obesity (high BMI), and type of occupation. • Predominantly leg pain, typically involving the foot 	<p>Radiographs not initially indicated [B] unless patient age >50 or has progressive neurologic deficits</p>
<p>(B) Suspected degenerative spondylolithesis/lateral stenosis</p> <ul style="list-style-type: none"> • Back pain with or without leg pain • Increased pain with activity • S&S with or without neurologic deficit 	<p>Radiographs indicated if patient age >50 or has progressive neurologic deficits: PA (or AP), lateral lumbar views [GPP]</p>
<p>(C) Suspected lumbar degenerative spinal stenosis</p> <ul style="list-style-type: none"> • MC >65 yoa (sensitivity of 0.7; specificity of 0.69) • Neurogenic claudication • Variable neurologic deficit (numbness, weakness, etc) 	<p>Radiographs indicated if patient age >50 or has progressive neurologic deficits: PA (or AP), lateral lumbar views [C]</p>
Suspected causes of sciatica:	
<p>(A) Lumbar disc herniation (B) Degenerative spondylolithesis/lateral stenosis (C) Lumbar degenerative spinal stenosis</p>	<p>Special investigations not initially indicated [C]</p> <p>Co-management or specialist referral recommended even if conventional radiographs are unremarkable:</p> <ol style="list-style-type: none"> 1. After failed conservative therapy (4-6 wk) 2. For preoperative planning; 3. If patient’s neurologic status is deteriorating (progressive deficit, disabling leg pain); <ul style="list-style-type: none"> • MRI, CT
<p>Adult patient reevaluation in the absence of expected treatment response or worsening after 4-6 wk Should patient fail to improve as expected or marginally improve within 4-6 wk of initial evaluation, the clinician must review history and physical findings and request appropriate diagnostic imaging studies.</p>	<p>Radiographs indicated [B] PA (or AP), lateral lumbar views</p> <p>Additional views not routinely indicated [C] Spot lateral, oblique. lateral flexion films may be indicated in scoliosis evaluation</p> <p>Comanagement or specialist referral recommended even if conventional radiographs are unremarkable</p> <ol style="list-style-type: none"> 1. And if conventional radiography reveals suspected pathology. 2. After failed conservative therapy (4-6 wk), 3. If patient neurologic status is deteriorating (progressive deficit, disabling leg pain); 4. If clinical signs suggest instability. Presumed instability is loosely defined as >10° of angulation or 4 mm of vertebral displacement on flexion and extension lateral radiographs. However, diagnostic criteria, natural history, and surgical indications remain controversial 5. For preoperative planning; <p>Special investigations [C]</p> <ul style="list-style-type: none"> • MRI or CT scan
<p>Adults with complicated (ie, “red flag”) LBP and indicators of contraindication to SMT (relative/absolute):</p> <p>Presence of the following indicator(s) should alert the clinician to possible underlying pathology. Presence of a red flag alone may not necessarily indicate the need for radiology.</p> <ul style="list-style-type: none"> • <i>Patient <age 20 and >age 50, particularly with S&S suggesting systemic disease**</i> 	<p>Radiographs indicated [B] PA (or AP), lateral lumbar views.</p> <p>Additional views: Hibb’s, (Spot angled PA or AP lumbosacral), oblique SI views</p>

(continued on next page)

Table 3 (continued)

Patient Presentation	Recommendations
<ul style="list-style-type: none"> • <i>No response to care after 4 wk</i> • <i>Significant activity restriction >4 wk</i> • <i>Nonmechanical pain (unrelenting pain at rest, constant or progressive S&S)</i> • <i>Suspected inflammatory—spondyloarthritides</i> • <i>Suspected compression fracture</i> • <i>Suspected neoplasia</i> • <i>Suspected infection</i> • <i>Suspected failed surgical fusion</i> • <i>Progressive or painful structural deformity</i> • <i>Elevated laboratory examination and positive S&S</i> 	<p>Advanced imaging and specialist referral recommended:</p> <ol style="list-style-type: none"> 1. In the presence of a potentially serious pathology as suggested by the patient history, examination, and/or radiograph; 2. In the absence of clinical improvement after 4-6 wk of therapy; 3. If function does not improve or deteriorates; 4. If patient neurologic status is deteriorating (progressive deficit, disabling leg pain); 5. With painful or progressive structural deformity 6. For unstable segment (spondylolisthesis or pathological process); 7. When patient has persisting S&S; 8. In complication from treatment (possible fracture, new/progressive neurologic deficit, considerable pain, or disability, etc) <p>Special investigations [B] Even if conventional radiographs are negative</p> <ul style="list-style-type: none"> • MRI, CT, NM
<ul style="list-style-type: none"> • <i>Suspected CES</i> <p>The classic syndrome includes LBP, bilateral or unilateral sciatica, saddle anesthesia, motor weakness of the lower extremities that may progress to paraplegia, urinary retention, or bowel and bladder incontinence.</p>	<p>Emergency referral without imaging [B]</p> <p>Special investigations [C] (see above for details)</p>
<ul style="list-style-type: none"> • <i>Suspected AAA</i> <p>Early S&S may include abdominal pain, backache, and feeling of fullness or abdominal pulsation.</p>	<p>Referral for specialized investigations [B]</p> <ul style="list-style-type: none"> • Management (ultrasound screening/monitoring and surgical consultation) according to patient history and size of AAA
<ul style="list-style-type: none"> • <i>Truncal symptoms attributed to the presence or worsening of aortic aneurysms including dissection/rupture/occlusion or traumatic aortic injury</i> <p>Cardiovascular shock and/or syncope, severe, tearing/ripping midabdominal sensation, back, groin or testicular pain; pressure upon lumbar spine causing excruciating boring pain in the abdomen or back; hypotension; absence distal lower limb pulses</p>	<p>Emergency referral without imaging [GPP]</p> <ul style="list-style-type: none"> • It is vital to recognize the S&S of dissecting AAA as this is a surgical emergency.

Table 4. Summary of Recommendations—Nontraumatic Thoracic Spine Disorders

Patient Presentation	Recommendations
<p>Adult patient with uncomplicated* acute thoracic spine pain (<4 wks' duration)</p> <p style="text-align: center;"><i>AND</i></p> <p>Adult patient with uncomplicated* subacute (4-12 wks' duration) or persistent (>12 wks' duration) thoracic spine pain and no previous treatment trial.</p> <p>* <i>Uncomplicated definition:</i> nontraumatic thoracic pain without neurologic deficits or indicators of potentially serious pathologies</p>	<p>Radiographs not routinely indicated [B]</p> <p>Special investigations not indicated [B]</p>
<p>Adult patient: reevaluation in the absence of expected treatment response or worsening after 4 wk.</p> <p>Should patient fail to improve as expected or marginally improve within 4 wk of initial evaluation, the clinician must review history and physical findings and request appropriate diagnostic imaging studies.</p>	<p>Radiographs indicated [B] AP, lateral thoracic spine views</p> <p>Additional views: swimmer's view</p> <p>Co-management or specialist referral recommended</p> <ol style="list-style-type: none"> 1. In suspected pathology as seen on conventional radiography 2. After failed conservative therapy (4 wk), 3. If patient neurologic status is deteriorating (progressive deficit, disabling leg pain); <p>Special investigations [C]</p> <ul style="list-style-type: none"> • MRI or CT scan

Table 4 (continued)

Patient Presentation	Recommendations
<p>Adult patient with nontraumatic chest wall pain History and physical exam first need to rule out life-threatening conditions including pathologies of the heart, lungs and large vessels.</p>	<p>Emergency referral without imaging in life-threatening conditions [GPP]</p> <p>Special investigations [C]</p> <ul style="list-style-type: none"> • CT and MRI
<p>Musculoskeletal causes of chest wall pain (diagnosis of exclusion).</p>	<p>Radiographs not routinely indicated [D]</p>
<p>Adult patient with complicated (ie, “red flag”) thoracic pain and indicators of contraindication to SMT (relative/absolute) Presence of the following indicator(s) should alert the clinician to possible underlying pathology. NB. Presence of a red flag alone may not necessarily indicate the need for radiography. Patient <age 20 and >age 50, particularly with S&S suggesting systemic disease**</p> <ul style="list-style-type: none"> • No response to care after 4 wk. • Significant activity restriction >4 wk • Nonmechanical pain (unrelenting pain at rest, constant or progressive S&S) • Persistent localized pain (>4 wk) • Progressive or painful structural deformity: scoliosis, kyphoscoliosis (Otani 2001) • Symptoms associated with neurologic signs in the lower extremities • Suspected inflammatory spondyloarthropathy • Suspected neoplasia • Suspected infection • Suspect failed surgical fusion • Elevated laboratory examination and positive S&S • In recent significant trauma (any age) 	<p>Radiographs indicated [B] AP, lateral thoracic spine views.</p> <p>Additional views: spot view. In suspected inflammatory spondyloarthropathy, consider: Hibb’s (spot angled AP lumbosacral), oblique SI views</p> <p>Advanced imaging and specialist referral recommended even if conventional radiographs are unremarkable:</p> <ol style="list-style-type: none"> 1. In presence of a potentially serious pathology as suggested by the patient history, examination and/or radiograph; 2. In the absence of clinical improvement after 4 to 6 wk of therapy; 3. If function does not improve or deteriorates; 4. If patient neurologic status is deteriorating (progressive deficit, disabling leg pain); 5. With painful or progressive structural deformity 6. For unstable segment (spondylolisthesis or pathological process); 7. When patient has persisting S&S; 8. In complication from treatment (possible fracture, new/progressive neurologic deficit, considerable pain or disability, etc) <p>Special investigations [B]</p> <ul style="list-style-type: none"> • MRI, CT, NM
<ul style="list-style-type: none"> • Suspected acute thoracic aortic aneurysms dissection/rupture/occlusion or traumatic aortic injury Severe, tearing/ripping chest sensation, back pain; hypotension; absent distal pulse. High index of suspicion in connective tissue disorders and diseases with genetic predisposition for ascending aortic aneurysms. 	<p>Emergency referral without imaging [GPP]</p>
<ul style="list-style-type: none"> • Suspected compression fracture Severe onset of pain (with or without appearance of spinal deformity) after minor trauma in older patients. Patients with thoracic or lumbar spine osteoporotic fractures report pain mainly in the lumbosacro-gluteal area. Look for Hx of repetitive stress of sufficient severity or Hx of high risk osteoporosis <p>Risk factors for additional vertebral fractures: Histories of a previous fracture, greater age, lower femoral neck bone mass density, shorter height.</p>	<p>Radiographs indicated [B]: AP, lateral thoracic spine views.</p> <p>Additional views [D]: supine cross-table lateral view in suspected osteoporotic vertebral pseudoarthrosis</p> <p>Special investigations [D]</p> <ul style="list-style-type: none"> • MRI/CT if initial radiographs are positive, difficult to interpret, in presence of complex lesions, for suspected ligamentous instability or neural injuries.
<p>Suspected osteoporosis See osteoporosis clinical decision rules</p>	<p>Radiographs are <i>unreliable</i> for assessment of bone mass changes before at least a 30%-50% loss</p> <p>Special investigations [B] if clinical decision rules are positive</p> <ul style="list-style-type: none"> • Bone densitometry or dual-energy x-ray absorptiometry (DXA)

(continued on next page)

Table 4 (continued)

Patient Presentation	Recommendations
Adult patient with nonpainful and nonprogressive scoliosis	Radiographs not routinely indicated [C]
Adult patient with painful or progressive scoliosis	<p>Radiographs indicated [B] Erect sectional radiographs (better detail) or standing full-length PA (14 × 36 in) and lateral sectionals</p> <p>Additional views: 1. Right and left lateral bending</p> <p>Follow-up evaluation dictated by clinical progression [C]</p> <p>Repeat radiographs, specialist referral and advanced imaging recommended [B]:</p> <ol style="list-style-type: none"> 1. In the absence of clinical improvement; after 4 to 6 wk of therapy; 2. If function does not improve or deteriorates; 3. In presence of persisting S&S or considerable pain; 4. If patient neurologic status is deteriorating (progressive deficit, disabling leg pain); 5. With painful or progressive structural deformity (scoliosis, kyphoscoliosis); 6. With suspected segmental instability (this is common in adult scoliosis and should be considered with all manual therapy intervention). 7. With suspected pathological process; 8. With new or progressive neurologic deficit including claudication, significant radiculopathy or suspected syrinx; 9. To plan surgical intervention. <p>Special investigations [C] • Spiral CT, MRI, sequential discograms, facet blocks, epidural blocks, CT-myelogram.</p>

Table 5. Summary of Recommendations—Nontraumatic Cervical Spine Disorders

Patient Presentation	Recommendations
<p>Adult patient with acute uncomplicated* neck pain (<4 wks' duration)</p> <p>* <i>Uncomplicated definition:</i> nontraumatic neck pain without neurologic deficits or indicators of potentially serious pathologies—(see red flag list for details).</p>	<p>Radiographs not initially indicated [C]</p> <p>Special investigations not indicated [C]</p>
<p>Adult patient with nontraumatic neck pain and radicular symptoms</p> <p>(A) Suspected acute cervical disc herniation (CDH) (B) Suspected Acute cervical spondylotic radicular syndrome/lateral canal stenosis</p>	<p>Radiographs indicated [D/consensus] APOM, AP lower cervical, neutral lateral</p> <p>Additional views: oblique views, swimmer's view</p> <p>Comanagement or specialist referral recommended even if conventional radiographs are unremarkable</p> <ol style="list-style-type: none"> 1. After failed conservative therapy (4 wk) 2. For preoperative planning; 3. If patient neurologic status is deteriorating (progressive deficit, disabling arm pain); <p>Special investigations [B] MRI</p>

Table 5 (continued)

Patient Presentation	Recommendations
<p>Adult patient with uncomplicated* subacute (4-12 weeks duration) and persistent neck pain (>12 weeks) with or without arm pain.</p>	<p>Radiographs not initially indicated* [consensus] APOM, AP lower cervical, neutral lateral</p> <p>N.B. This recommendation was modified according to the recent findings of The Bone and Joint Decade 2000-2010 Task Force on Neck Pain and its Associated Disorders (see articles published in Spine 2008; 33(4S)). A majority of Delphi panelists agreed with this change (92% of 50 respondents).</p>
<p>Adult patient reevaluation in the absence of expected treatment response or worsening after 4 weeks.</p>	<p>Radiographs indicated [C] APOM, AP lower cervical, neutral lateral</p> <p>Additional views: Oblique views, Swimmer’s view, Flexion/Extension</p> <p>Comanagement or specialist referral recommended (even if conventional radiographs are unremarkable)</p> <ol style="list-style-type: none"> 1. If conventional radiography reveals suspected pathology. 2. After failed conservative therapy (4 wk), 3. If patient neurologic status is deteriorating (progressive deficit, disabling arm pain); 4. If clinical signs suggest subaxial cervical spine instability. (Moore 2006) 5. For preoperative planning; <p>Special investigations [B]</p> <ul style="list-style-type: none"> • MRI
<p>Adult patient with complicated (ie, “red flag”) neck pain and indicators of contraindication to SMT</p> <p>Presence of the following indicator(s) should alert the clinician to possible underlying pathology. N.B. Presence of a red flag alone may not necessarily indicate the need for radiography.</p> <ul style="list-style-type: none"> • Patient <age 20 and >age 50, particularly with S&S suggesting systemic disease** • No response to care after 4 wk • Significant activity restriction >4 wk • Nonmechanical pain (unrelenting pain at rest, constant or progressive S&S) • Neck rigidity in the sagittal plain in the absence of trauma (discitis, infection, tumor, meningitis, etc); • Dysphasia; • Impaired consciousness; • Central nervous system S&S (cranial nerves, pathological reflexes, long tract signs); • High risk ligament laxity populations/suspected atlantoaxial instability(see details below); • Arm or leg pain with neck movements, suspected cervical myelopathy and radiculo-myelopathy (see details bellow); • Sudden onset of acute and unusual neck pain and/or headache (typically occipital) with or without neurologic symptoms, suspected cervical artery dissection (VAD, CAD), TIA (VBI, carotid artery ischemia), stroke (see details bellow); • Hx of severe trauma (see Trauma section). 	<p>Radiographs indicated [B] APOM, AP lower cervical, neutral lateral</p> <p>Additional views: flexion/extension, oblique views, pillar view</p> <p>Advanced imaging and specialist referral recommended:</p> <p>Special investigations [B]</p> <ul style="list-style-type: none"> • MRI

(continued on next page)

Table 5 (continued)

Patient Presentation	Recommendations
<p>In addition, also consider general red flags (usually applied to LBP) which may apply to the cervical spine</p> <ul style="list-style-type: none"> • <i>Suspected neoplasia</i> • <i>Suspected infection (discitis, osteomyelitis, tuberculosis)</i> • <i>Suspect failed surgical fusion</i> • <i>Progressive or painful structural deformity</i> • <i>Elevated laboratory examination and positive S&S</i> 	
<p>• Suspected atlantoaxial instability (AAI) High risk ligament laxity populations/possible atlantoaxial instability include</p> <p>(a) Active inflammatory arthritides (b) Congenital disorders and hereditary connective tissues disorders</p>	<p>Radiographs indicated [B] APOM, AP lower cervical, neutral lateral</p> <p>Additional views [D]: flexion/extension laterals</p> <p>Monitoring, advanced imaging and specialist referral recommended:</p> <ol style="list-style-type: none"> 1. ADI >3 mm, vertical dislocation, lateral, posterior or subaxial subluxations 2. Upward odontoid translocation (pseudobasilar invagination) 3. In presence of neurologic S&S <p>Special investigations [C]</p> <ul style="list-style-type: none"> • CT, MRI
<p>• Suspected cervical compressive myelopathy (CCM) and radiculo-myelopathy</p>	<p>Radiographs indicated [C] APOM, AP lower cervical, neutral lateral and bilateral oblique views.</p> <p>Additional views: swimmer's view</p> <p>Refer patient for investigation and possible surgical intervention:</p> <ol style="list-style-type: none"> 1. After failed conservative therapy (4 wk), 2. If patient's neurologic status is deteriorating (progressive deficit, disabling arm pain); 3. For preoperative planning; <p>Special investigations [C]</p> <ul style="list-style-type: none"> • MRI (CT-myelography if not available). Electrophysiologic testing such as somatosensory evoked potentials (SSEP) may be useful.
<p>• Suspected cervical artery dissection (VAD, CAD), TIA (VBI, carotid artery ischemia), stroke The most important points in the history and chief complaint, which would warn of a possible cervical artery disease, are:</p> <ol style="list-style-type: none"> a. S&S of VBI—the "5D's And 3 N's": dizziness, dysphasia, dysarthria (hoarseness), drop attacks, diplopia (or other visual problems), ataxia of gait (hemiparesis), nausea (possibly with vomiting), numbness (hemianesthesia), nystagmus; b. S&S of carotid artery ischemia/stenosis: confusion, dysphasia, headache, anterior neck and/or facial pain, hemianesthesia, hemiparesis or monoparesis, visual field disturbances. c. Neck or occipital pain with sharp quality and severe intensity or severe and persistent headache that is sudden and unlike any previous experienced pain or headache (even when it is suspected the pain is of a musculoskeletal or neuralgic origin) <p>Should cervical artery problems be suspected, a thorough workup is indicated.</p>	<p>Emergency referral without imaging [GPP] Urgent referral should be made for appropriate investigation and treatment in patient presenting S&S of cerebrovascular ischemia or when S&S of head/neck pain is suspicious for an acute cervical artery disease.</p> <p>Special investigations [C]</p> <ul style="list-style-type: none"> • Initial investigation often includes CT scan to R/O hemorrhagic stroke. <p>Appropriate consultation and/or diagnostic procedures to evaluate the status of the cerebral circulation required in patients presenting with significant risk factors for cervical artery dissection. In such cases, approach the treatment with caution until a specific determination is made.</p>

APPENDIX C. PERTINENT CLINICAL INFORMATION

(1) Thoracic spine trauma

The thoracolumbar injury severity scale (TLISS) may be helpful in referral decision-making process. The scale is based on the mechanism of injury, the integrity of the posterior ligamentous complex, and the neurologic status.^{i,iii} The TLISS has good reliability and compares favorably to other contemporary thoracolumbar fracture classification systems.ⁱⁱⁱ

Description	Points
1. Injury mechanism	
a. Compression	
• Simple compression	1
• Lateral angulation >15°	1
• Burst	1
b. Translation/rotational	3
c. Distraction	4
2. Posterior ligamentous complex (PLC)	
disrupted in tension, rotation, or translation	
a. Intact	0
b. Suspected/indeterminate	2
c. Injured	3
3. Neurologic status	
a. Nerve root involvement	2
b. Cord, conus medullaris involvement	
• Incomplete	3
• Complete	2
c. Cauda equina involvement	3

The score is a total of 3 components: injury mechanism, PLC integrity, and neurologic status. A score of ≤ 3 suggests nonoperative treatment (bracing); 4, operative or nonoperative treatment; and ≥ 5 suggests operative treatment.

(2) Cervical spine trauma

Risk factors for WAD may include personal, societal, and environmental factors. Both physical and psychological factors are likely to play a role in recovery or non-recovery from whiplash injury. High initial symptom severity (high initial pain level, pain-related limitations, greater number of symptoms and painful body parts), greater psychological distress and passive coping may predict poorer outcome.^{iv} Finally, an important association exists between a Hx of a whiplash associated disorder, pain intensity, and disability and comorbidity (headache, LBP, digestive, and cardiovascular disorders).^{vi,vii}

a. Adult Patient with Acute Neck Injury and Positive CCSR

Proposed whiplash associated disorders classification included Foreman and Croft (1995), the Quebec Task Force^{viii} and revised classification on Whiplash Associated Disorders.^{ix-x}

- **Grade 0:** no neck pain or sensitivity
- **Grade I:** neck pain, stiffness or tenderness only. No physical signs
- **Grade II:** neck pain, stiffness or tenderness, decreased range of motion (musculoskeletal signs). Consider psychological impairment (elevated psychological distress, high levels of posttraumatic stress), and interference with daily living.
- **Grade III:** neck complaints accompanied by neurologic signs such as decreased or absent deep tendon reflexes, weakness and/or sensory deficits. Also consider psychological impairment (elevated psychological distress, levels of posttraumatic stress), and interference with daily living.
- **Grade IV:** fractures, dislocation (or spinal cord lesions).

A biopsychosocial model may improve WAD treatment and help prevent chronic disability.^{xi}

Classification of patients with chronic WAD may include measures of self-efficacy, disability, and coping such as the NDI, Multidimensional Pain Inventory-Swedish version (MPI-S), The Self-Efficacy Scale, and the Coping Strategies Questionnaire.^{xii}

b. Cervical Spine Injury Severity Score

The Cervical Spine Injury Severity Score (CSISS), which was developed to measure stability after cervical spine trauma, has excellent intraobserver agreement (ICC, 0.97-0.99) and interobserver agreement (ICC, 0.75-0.98). This classification system is based on morphological descriptions and on quantifiable values of stability and applies to all fractures of the

subaxial spine from the caudal aspect of C2 to T1. The score is based on bony and ligamentous disruption of the 4 columns of the cervical spine (anterior, right pillar, left pillar, and posterior osseous ligamentous complex) and correlates to increasing instability. Each column is graded using an analog scale from 0 to 5. Thus, the injury severity score ranges from 0 to 20, with 0 being no injury and 20 the most severe. The CSISS does not take into account neurologic function or deficits. A morphologic description of subaxial cervical fracture is proposed.^{xiii}

For more information on nonoperative management and treatment of spinal injuries and timing of surgical intervention in spinal cord injuries, see the reviews from Reichtine^{xiv} and Fhelings.^{xv}

(3) Clinical Criteria for the Diagnosis of LDH.^{xvi,xvii,xviii,xix}

Nerve root pain due to disc-root conflict should be deemed to be a symptom of multifactorial origin in which the neural and perineural inflammatory reactions and their mediators play a major role that is flanked by venous stasis due to mass effect on the perineural circulation. Nerve compression appears to play an adjuvant role by generating nerve conduction abnormalities due to fiber demyelination by a direct or indirect anoxic-ischemic mechanism.^{xx}

Clinical criteria for the diagnosis of LDH include sciatic pain that originates at the back or buttock and radiates below the knee, either unilaterally or bilaterally, typically involving the foot. Often, a history of recurrent LBP over several months or years, with occupational risk factors, is reported before leg pain onset. The patient often has difficulty rising from a sitting or supine position and typically experiences only partial pain relief while supine. Coughing or sneezing often exacerbates the low back and leg pain (positive Valasalva).

On physical examination, the patient is in acute distress and may manifest an antalgic posture/gait and decreased lumbar lordosis due to marked paraspinal muscle guarding. Lumbar spine ranges of motion may be restricted, especially in forward flexion. Lower extremity neurologic examination^{xxi,xxii,xxiii} may reveal diminished deep tendon reflexes, loss of motor power, and sensory deficit of the involved nerve root (L4, L5, or S1). Reduced ankle reflex (L5-S1), weakness of ankle dorsiflexion (L4-5), foot drop, or weakness of the big toe (L5) requires special attention, as does reduced light touch in the L4, L5, or S1 dermatome. Straight leg raise will typically be restricted on the involved side with a high sensitivity (0.8) but a low specificity (0.4), whereas a crossed SLR (well leg-raise test) has a low sensitivity (0.35) but a high specificity (0.9) for the diagnosis of herniated discs. This may be confirmed by nerve root tension tests such as bowstring and Braggard's.

Wasserman maneuver (Elys test) evokes pain by stimulating the L2-L4 nerve roots when hyperextending the hip with the knee flexed at 90 while the patient is prone. Examination may reveal reduced patellar reflex (L2-L4), weakness of the quadriceps (L2-4), and possibly of ankle dorsiflexion (L4-5), and/or reduced light touch in L2, L3, or L4 dermatome. Bony and soft tissue tenderness may be noted over the lumbar and gluteal region, although not a prominent feature.

(4) Thoracic Spine Osteoporosis

A woman 65 years of age with one vertebral fracture has a 1 in 4 chance of another fracture over 5 years, which can be reduced to 1 in 8 by treatment. Positive treatment decisions are often contingent on identifying a vertebral fracture.^{xxiv}

Although it may not be possible to restore normal bone structure in women with osteoporosis, antiresorptive agents (eg, bisphosphonates, selective estrogen receptor modulators) can substantially reduce the risk of new vertebral fracture, even without corresponding increase in bone mass density, by reducing excessive bone turnover. Elevated bone turnover can be assessed with a number of biochemical markers such as osteocalcin, bone-specific alkaline phosphatase, C-telopeptides of type I collagen and N-telopeptide cross-links of type I collagen.^{xxv}

a. Important Osteoporosis Risk Factors:^{xxvi,xxvii,xxviii,xxix,xxx,xxxi,xxxii,xxxiii,xxxiv,xxxv,xxxvi,xxxvii}

- a. Age ≥ 50 (higher risk >70);
- b. BMI or weight ($<19 \text{ kg/m}^2$ or $<57 \text{ kg}$);
- c. Ethnic group or race (white or Asian);
- d. Menopause before age 45;
- e. Loss height ($>4 \text{ cm}$);
- f. Hx of corticosteroid Tx for rheumatoid arthritis and other immune and inflammatory disorders ($5\text{-}7.5 \text{ mg/d} \times 3 \text{ months}$);
- g. Hx of fragility fracture after age 45 of one of the following sites: hip, rib, wrist;
- h. No Hx of hormone replacement therapy/estrogen such as premarin, estrace, estraderm, estrab;
- i. Vitamin D and vitamin K deficiency, aluminum bone disease (dialysis, TNP), hypogonadism, malabsorption syndromes; Hx of anorexia/bulimia nervosa, female athlete triad
- j. Excessive alcohol/caffeine intake; smoking, dilantin;

- k. Immobilization >4 weeks after age 45;
- l. Low birth size and poor childhood growth
- m. Elevated blood pressure;
- n. Cognitive decline.

b. Osteoporosis Risk Assessment^{xxxviii,xxxix,xl,xli,xlii,xliii,xliv}

In healthy perimenopausal and early postmenopausal women (45-64 years), consider using the Osteoporosis Self Assessment Tool (OST score): the OST considers only 2 variables: (weight in kilograms – age)/5. The cut-off for a positive test is <2, indicating this woman should be referred for DEXA.

In higher-risk patients, use either one of the following 3 clinical tools as they were all found to be highly sensitive:

1. Osteoporosis Risk Assessment Instrument (ORAI) which considers age, weight, and estrogen use (Table 1).
2. Simple Calculation of Osteoporosis by Estimation (SCORE) which considers 6 variables (race, RA, history of fracture, age, estrogen, weight (in pounds) (Table 2).
3. Age, years after menopause, age at menarche, BMI (AMMED) (Table 3).

Table 1. Osteoporosis Risk Assessment Instrument (ORAI).

ORAI variables	Points added to index
Age >75 y	15
Age 65-74 y	9
Age 55-64 y	5
Weight <60 kg	9
Weight 60-90 kg	3
Not current HRT user	2

The cut-off for a positive test is >8; indicating this woman should be referred for DEXA.

Table 2. Simple Calculation of Osteoporosis by Estimation (SCORE).

SCORE variables	Points added to index
Race	5 if NOT black
Rheumatoid arthritis	4 if has RA
History of fracture	4 for EACH type (wrist, rib, hip) of nontraumatic fractures >45 y (maximum score of 12)
Age (y)	3 times first digit of age
Estrogen	1
Weight (in lbs)	-1 times weight divided by 10 and truncated to integer

The cut-off for a positive test is >6, indicating this woman should be referred for DEXA.

Table 3. Age, Years After Menopause, Age at Menarche, BMI (AMMEB).

AMMEB Variables	Points added to index
Age (y)	15 if 75+, 9 if 65-74, 5 if 55-64, 0 if <55
BMI	6 if <20, 2 if 20-23, 1 if 24-26, 0 if >26
Age at menarche	0 if 11, 1 if 11-13, 6 if >13
Postmenopausal period	5 if >16, 3 if 12-16, 1 if 5-11, 0 if >5.

The cut-off for a positive test is ≥ 10, indicating this woman should be referred for DEXA

(5) Clinical Criteria for the Diagnosis of Cervicobrachial Syndrome

Cervical radiculopathy is a disorder of the cervical spinal nerve root, and most commonly is caused by a CDH or other space-occupying lesion. The annual incidence of cervical spine disc herniation with radiculopathy is approximately 5.5 per 100 000 and tends to affect patients aged 35 to 55 years of age. The levels most commonly affected are C5-C6 and C6-C7.^{xlv} A clinical diagnosis of CDH is possible in only 50% to 75% of cases of cervical radiculopathy as a wide variation of presentations exists even with only 1 nerve root.^{xlvi,xlvii} Patients should be monitored as signs and symptoms may progress in the first 2 to 3 weeks.

Historical questions with diagnostic accuracy include shoulder/scapular pain, symptoms influenced by neck movement, presence of numbness/tingling, intermittent signs and symptoms, and symptoms that interrupt sleep.^{xlviii}

On physical examination, the patient is generally in acute distress and may manifest an antalgic posture. The upper limb tension test (ULTT) appears useful as a screening test given its high sensitivity (0.97). If the ULTT is negative, then cervical radiculopathy (CR) can essentially be ruled out, and the need for further work up or treatment for CR is minimized. Findings with useful diagnostic accuracy include cervical spine ranges of motion restriction (ipsilateral rotation less than 60°, and neck flexion less than 55°). Although not very sensitive, the following tests have moderate to high specificity for CR: traction/neck distraction, spurling, Valsalva, and the shoulder abduction test, reduced or absent dermatomal sensation (C5 most accurate), motor power, and deep tendon reflex (biceps most accurate). The probability of the condition increases with several positive findings.^{xlix,1}

(6) Persistent Neck Pain

(a) Identification of psychological risk factors (yellow flags):^{li}

- Obvious psychological distress
- Severe pain beyond what is expected
- Hx of prior significant pain recurrences
- Higher than expected functional impairment
- Unexplained widespread pain
- Pain and limitation not consistent with objective findings.

(b) Mental status indicators of significant anxiety or depression:^{li}

- Insomnia or nightmares
- Irritability
- Withdrawal
- Panic episodes or anxiety during the day or night
- Persistent tearfulness
- Poor concentration
- Inability to enjoy
- Poor appetite/weight loss
- Poor libido
- Thoughts that “life is not worth living”

Clinicians are encouraged to use validated patient self-administered questionnaires to evaluate perceived neck pain, function, disability, and psychosocial status.

(c) Suspected cervical artery dissection (VAD, CAD), TIA (vertebrobasilar ischemia, carotid artery ischemia), stroke.

Acute neck pain is generally due to musculoskeletal conditions that respond well to conservative therapy. Risks of adverse events need to be addressed when considering any form of therapy. Several risk factors have been proposed for cervical artery dissection.^{lii,liii,liiv,lv} Cervical artery dissection may be considered multifactorial,^{lvi} resulting from cumulative events over a period of time. Patient having cervical artery dissections tend to be females under the age of 45: recently, an etiologic model to help explain the pathogenesis of cervical artery dissection has been proposed.^{lvii} Recent studies and reviews discussing the nature of the association between vertebrobasilar stroke and chiropractic care are available elsewhere.^{ix-ixi}

It is likely that patients in the early stages of VBA stroke are presenting to a health care professional because of neck pain and headache due to pre-existing vertebral artery dissection which is a risk factor for VBA stroke.

None of the proposed risk factors may be visualized on conventional radiography. Few case reports have linked advanced cervical spondylosis to vertebral artery dissection, suggesting osteophytes off the uncovertebral joint or superior articular process may be associated to a dissection. Advanced cervical spondylosis mostly concerns the older age population which would normally undergo radiographs when presenting with a neck complaint. Major neck trauma is likely an independent risk factor for CAD. Again, imaging studies are indicated in those circumstances.

References

- i. Harrop J, Vaccaro AR, Hurlbert RJ, et al. Intrarater and interrater reliability and validity in the assessment of the mechanism of injury and integrity of the posterior ligamentous complex: a novel injury severity scoring system for thoracolumbar injuries. Invited submission from the joint section meeting on disorders of the spine and peripheral nerves, March 2005. *J Neurosurg Spine* 2006; 4(2):118-22.
- ii. Vaccaro AR, Lehman RA Jr, Hurlbert RJ, et al. A new classification of thoracolumbar injuries: the importance of injury morphology, the integrity of the posterior ligament complex, and neurologic status. *Spine* 2005; 30(20):2325-33.
- iii. Vaccaro AR, Baron EM, Sanfilippo J, et al. Reliability of a novel classification system for thoracolumbar injuries: The Thoracolumbar Injury Severity Score. *Spine* 2006; 31(11) Suppl: S62-S69.

- iv. Sterling M, Jull G, Vicenzino B, Kenardy J, Darnell R. Physical and psychological factors predict outcome following whiplash injury. *Pain* 2005; 114(1-2):141-8.
- v. Sterling M, Jull G, Kenardy J. Physical and psychological factors maintain long-term predictive capacity post-whiplash injury. *Pain* 2006; 122(1-2):102-8.
- vi. Berglund A, Alfredsson L, Cassidy JD, Jensen I, Nygren A. The association between exposure to a rear-end collision and future neck or shoulder pain: a cohort study. *J Clin Epidemiol* 2000; 53:1089-94.
- vii. Berglund A, Alfredsson L, Jensen I, Cassidy JD, Nygren A. The association between exposure to a rear-end collision and future health complaints. *J Clin Epidemiol* 2001; 54:851-6.
- viii. Spitzer WO. Monographie scientifique du groupe de travail qu ebecois sur les troubles associ es   l'entorse cervicale, Soci t  de l'Assurance Automobile du Qu ebec 1995.
- ix. Hartling L, Brison RJ, Arden C, Pickett W. Prognostic value of the Quebec classification of whiplash-associated disorders. *Spine* 2001; 26(1): 36-41.
- x. Sterling M. A proposed new classification system for whiplash associated disorders-implication for assessment and management. *Man Ther* 2004; 9: 60-70.
- xi. Ferrari R. Prevention of chronic pain after whiplash. *Emerg Med J.* 2002; 19(6):526-30.
- xii. Soderlund A, Denison E. Classification of patients with whiplash associated disorders (WAD): reliable and valid subgroups based on the Multidimensional Pain Inventory (MPI-S). *Eur J Pain.* 2006; 10(2):113-9.
- xiii. Moore TA, Vaccaro AR, Anderson PA. Classification of lower cervical spine injuries. *Spine.* 2006; 31(11 Suppl):S37-43.
- xiv. Rehtine GR. Nonoperative management and treatment of spinal injuries. *Spine* 2006; 31(11) Suppl, pp S22-27.
- xv. Fehling MG, Perrin R. The timing of surgical intervention in the treatment of spinal cord injury: a systematic review of recent clinical evidence. *Spine* 2006; 31(11) Suppl, pp S28-53.
- xvi. Deyo, R.A., Loeser, J.D. and Bigos, S.J., Herniated lumbar intervertebral disk, *Ann Intern Med* 1990;112:598-603.
- xvii. Crawford, C. and Hannan, R.F., Management of acute lumbar disk herniation initially presenting as mechanical low back pain, *J Manipulative Physiol Ther* 1999;22:235-44.
- xviii. Kostuik, J.P., Harrington, I., Alexander, D., Rand, W. and Evans, D., Cauda equina syndrome and lumbar disc herniation, *J Bone Joint Surg Am* 1986;68:386-91.
- xix. Busse, J., Hsu, W., Rapid progression of acute sciatica to cauda equina syndrome, *J Manipulative Physiol Ther* 2001;24:350-5.
- xx. Leonardi M, Simonnetti L, Agati R. Neuroradiology of spine degenerative disease. *Best Pract Res* 2002; 16(1):59-87.
- xxi. Deyo R, Rainville J, Kent DL. The rational clinical examination. What can the history and physical examination tell us about low back pain. *JAMA* 1992; 268:760-5.
- xxii. Ioto K, Aebi M, Alini M. Aging and degeneration: Cause and effect of intervertebral disc damage? In Gunzburg R, Szpalski M. Lumbar disc herniation. Lippincott Williams & Wilkins, 2000.
- xxiii. Andersson GB, Deyo RA. History and physical examination in patients with herniated lumbar discs. *Spine* 1996; 21(24S):10S-18S.
- xxiv. Kaptoge S, Ambrecht G, Felsenberg D, Lunt M, O'Neill TW, Silman AJ, Reeve J; EPOS Study Group. When should the doctor order a spine X-ray? Identifying vertebral fractures for osteoporosis care: results from the European Prospective Osteoporosis Study (EPOS). *J Bone Miner Res.* 2004; 19(12):1982-93.
- xxv. Melton III LJ, Kallmes DF. Epidemiology of vertebral fractures: Implications for vertebral augmentation. *Acad Radiol* 2006; 13:538-45.
- xxvi. Kin Yuen C, Kendler D, Khan A, Brown J, Fortier M. Ost eoporose. Appendix J: La prise en charge de l'ost eoporose au Canada. Diagnostic, pr evention et traitement. St-Michael's Hospital, Universit  de Toronto 2002.
- xxvii. Nguyen HV, Ludwig S, Gelb D. Osteoporotic vertebral burst fractures with neurologic compromise. *J Spinal Disorders & techniques* 2003; 16(1):10-9.
- li Hanley DA. Osteoporosis. In: Gray J. Therapeutic choices. 4th ed. Canadian Pharmacists Association, 2003:636.
- xxviii. Lignes directrices de pratique clinique 2002 pour le diagnostic et le traitement de l'ost eoporose au Canada. *JAMC* 2003; 168(6 Suppl):SF1-38.
- xxiv. Cappuccio FP, Meilahn E, Zmuda JM, et al. High blood pressure and bone mineral loss in elderly white women: a prospective study. *Lancet* 1999; 354:971-75.
- xxx. Yaffe K, Browner W, Cauley J, et al. Association between bone mineral density and cognitive decline in older women. *J Am Geriatric Society* 1999; 47:1176-1182.
- xxxi. D'Amelio P, Tamone C, Pluviano F, Di Stefano M, Isaia G. Effects of lifestyle and risk factors on bone mineral density in a cohort of Italian women: suggestion for a new decision rule. *Calcif Tissue Int* 2005; 77:72-8.
- xxxii. Pafumi C, Ciotta L, Farina M, Bosco P, Chiarenza M, Pernicone G, Calcagno A, Evola F, Cavallaro A, Iemmola A, Cianci A. Evaluation of bone mass in young amenorrhoeic women with anorexia nervosa. *Minerva Ginecol.* 2002; 54(6):487-91.

- xxxiii. Goodman LR, Warren MP. The female athlete and menstrual function. *Curr Opin Obstet Gynecol.* 2005 Oct; 17 (5):466-70.
- xxxiv. Misra M, Aggarwal A, Miller KK, Almazan C, Worley M, Soyka LA, Herzog DB, Klibanski A. Effects of anorexia nervosa on clinical, hematologic, biochemical, and bone density parameters in community-dwelling adolescent girls. *Pediatrics.* 2004; 114(6):1574-83.
- xxxv. Cockayne S, Adamson J, Lanham-New S, Shearer MJ, Gilbody S, Torgerson DJ. Vitamin K and the prevention of fractures: systematic review and meta-analysis of randomized controlled trials. *Arch Intern Med* 2006; 166(12):1256-61.
- xxxvi. Cooper C, Westlake S, Harvey N, Javaid K, Dennison E, Hanson M. Review: developmental origins of osteoporotic fracture. *Osteoporos Int* 2006; 17:337-47.
- xxxvii. Rud B, Jensen JEB, Mosekilde L, Nielsen SP, Hilden J, Abrahamsen B. Performance of four clinical screening tools to select peri- and early postmenopausal women for dual X-ray absorptiometry. *Osteoporos Int* 2005; 16:764-72.
- xxxviii. Cook RB, Collins D, Tucker J, Zioupos P. Comparison of questionnaire and quantitative ultrasound techniques as screening tools for DXA. *Osteoporos Int* 2005; 16:1565-75.
- xxxix. Mauck KF, Cuddihy MT, Atkinson EJ, Melton LJ. Use of clinical prediction rules in detecting osteoporosis in a population-based sample of postmenopausal women. *Arch Intern Med* 2005; 165 (5):530-6.
- xl. Gourlay ML, Miller WC, Richy F, Garrett JM, Hanson LC, Reginster JY. Performance of osteoporosis risk assessment tools in postmenopausal women aged 45-64 years. *Osteoporos Int.* 2005; 16(8):921-7.
- xli. Cadarette SM, Jaglal SB, Hawker GA. Fracture prevalence and treatment with bone-sparing agents: are there urban-rural differences? A population based study in Ontario, Canada. *J Rheumatol.* 2005; 32(3):550-8.
- xl.ii. Mauck KF, Cuddihy MT, Atkinson EJ, Melton LJ 3rd. Use of clinical prediction rules in detecting osteoporosis in a population-based sample of postmenopausal women. *Arch Intern Med.* 2005; 14;165(5):530-6.
- xl.iii. Brown, JP, Fortier M. Canadian Consensus Conference on Osteoporosis. SOGC clinical practice guideline No 172, February 2006. Available from: <http://www.sogc.org/guidelines/public/172E-CONS-February2006.pdf>
- xl.iv. Dreyer SJ, Boden SD. Nonoperative treatment of neck and arm pain. *Spine* 1998; 23(24):2746-54.
- xl.v. Nachemson A, Vingard E. Assessment of patients with neck and back pain: A best-evidence synthesis, Chap 9 in: Nachemson A, Jonsson E. Neck and back pain—The scientific evidence of causes, diagnosis and treatment. Lippincott Williams & Wilkins, Philadelphia, 2000.
- xl.vi. Ellenberg M, Honet JC, Treanor WJ. Cervical radiculopathy—Review article. *Arch Phys Med Rehabil* 1994; 75:342-52.
- xl.vii. Wainner RS, Fritz JM, Irrgang JJ, Boninger ML, Delitto A, Allison S. Reliability and diagnostic accuracy of the clinical examination and patient self-report measures for cervical radiculopathy. *Spine* 2003; 28(1):52-62.
- xl.viii. Tong HC, Haig AJ, Yamakawa K. The Spurling test and cervical radiculopathy. *Spine.* 2002; 27(2):156-9.
- xl.ix. Rubinstein SM, Pool JJ, van Tulder MW, Riphagen II, de Vet HC. A systematic review of the diagnostic accuracy of provocative tests of the neck for diagnosing cervical radiculopathy. *Eur Spine J.* 2007; 16(3):307-19.
- l. Tunks E. Neck Pain and Whiplash. In: Gray J. Therapeutic choices. 4th ed. Canadian Pharmacists Association, 2003:583.
- l.i. Haldeman, Kohlbeck FJ, McGregor M. Risk factors and precipitating neck movements causing vertebrasilar artery dissection after cervical trauma and spinal manipulation. *Spine* 1999 Apr 15; 24(8):785-94.
- l.ii. Rothwell DM, Bondy S, Williams JI. Chiropractic manipulation and stroke. A population-based case-control study. *Stroke* 2001; 32:1054-60.
- l.iii. Grau AJ, Brandt T, Forsting M, Winter R, Hacke W. Infection-associated cervical artery dissection. *Stroke* 1997; 28: 453-5.
- l.iv. Calvet D et al. Increase stiffness of the carotid wall material in patients with spontaneous cervical artery dissection. *Stroke* 2004;35(9):2078-82.
- l.v. Rubinstein SM, Peerdeman SM, Van Tulder MW, Riphagen I, Haldeman S. A systematic review of the risk factors for cervical artery dissection. *Stroke* 2005; 36(7):1575-80.
- l.vi. Rubinstein SM, Haldeman S, van Tulder MW. An etiologic model to help explain the pathogenesis of cervical artery dissection: implication for cervical manipulation. *J Manipulative Physiol Ther* 2006; 29(4):336-8.
- l.vii. Rubinstein SM, Haldeman S, van Tulder MW. An etiologic model to help explain the pathogenesis of cervical artery dissection: implication for cervical manipulation. *J Manipulative Physiol Ther* 2006; 29(4):336-8.
- l.viii. Chiropractic clinical practice guideline: evidence-based treatment of adult neck pain due to whiplash. *Can Chiropr Assoc* 2005;49(3).
- l.ix. Triano J, Kawchuk G, Hall C et al. Current concepts in spinal manipulation and cervical arterial incidents. NCMIC chiropractic solutions. 2006, p118.
- l.x. Boyle E, C  t   P, Grier AR, Cassidy JD. Examining vertebrasilar artery stroke in two canadian provinces. *Spine* 2008;33(4S):S170-S175.
- l.xi. Cassidy JD, Boyle B, C  t  , He H, Hogg-Johnson S, Silver FL, Bondy SJ. Risk of vertebrasilar stroke and chiropractic care — Results of a population-based case-control and case-crossover study. *Spine* 2008;33(4S):S176-S183.

APPENDIX D. GENERAL INDICATIONS FOR ADVANCED IMAGING IN SPINE DISORDERS

Indication	MRI	CT	CT-myelo	NM
Evaluation of primary bone neoplasm detected on radiographs	++	+	+	
Determining skeletal distribution of neoplasms or other multifocal skeletal disease				++
Clinical or laboratory tests suggesting plasma cell myeloma	++			
Posterior vertebral body scalloping seen on radiographs	++			
Known or suspected spinal cord tumors (malignant or nonmalignant), cord injury secondary to trauma, myelopathy, multiple sclerosis	++			
Myelopathy or radiculopathy when MRI is contraindicated or unavailable		+	++	
Spinal stenosis (congenital/degenerative)	+	++	+	
Cauda equine syndrome	++		+	
Persistent back or neck pain <i>with or without</i> radiculopathy and positive straight leg raise test, abnormal reflex, dermatome, or myotome with no improvement after 4 wk of conservative care	++			
Rapidly progressing neurologic deficit and/or motor weakness	++			
Infectious spondylodiscitis, osteomyelitis, tuberculosis	++	+		+
Rapidly progressing left-sided or atypical scoliosis	++			
Complicated disease processes or findings unexplained by more conservative tests	+	+		
Postoperative evaluation of arthrodesis		+		
Postoperative evaluation of recurrent symptoms (disc herniation and fibrosis)	++ GAD			
Burst fracture, pathological vertebral body compression fracture, or other unstable fractures	+	++		
Suspected occult fracture	+	+		++
Unstable or complex congenital anomalies or deformities of the spine	++	+		
Platybasia/basilar impression	++			
Enlarged sella as seen on cervical radiographs (brain) (MR imaging indicated)	++			

++ indicates first choice; +, second choice (must be determined on a case-by-case basis).^{a,b}

NB. For the estimated accuracy of imaging technique for lumbar spine conditions, please see Jarvik (2002).⁴

a. Adapted with permission from Peterson C. Canadian Guidelines for Imaging, 2002 (unpublished).

b. Taylor JAM. NYCC imaging protocols, 2003 (unpublished).

APPENDIX E. TYPICAL EFFECTIVE IONIZED RADIATION DOSE FOR COMMON IMAGING PROCEDURES*

Class	Typical effective dose (mSv)	Examples
0	0	Ultrasound, MRI
I	<1	Radiograph: cervical and thoracic spine, extremities, pelvis, and lungs
II	1-5	Lumbar spine radiograph, nuclear medicine, cervical spine CT
III	5-10	Chest and abdomen CT

* Classification of the typical effective dose of ionizing radiation from common imaging procedures. Adapted from: European Commission. Radiation Protection 118. Referral guidelines for imaging in conjunction with the UK Royal College of Radiologists; Italy 2001. p 21.

APPENDIX F. ADDITIONAL READING RECOMMENDED ON SPINAL MRI

DeVries RM, Manne A. Cervical MRI. Part I: a basic overview. Clin Chiropr 2003;6(3-4):137-43.

Wessely MA. Cervical MRI. Part II: a common disorders affecting the cervical spine. Clin Chiropr 2004;7(1):31-9.

Wessely MA. Magnetic resonance imaging of the thoracic spine. Part 1: normal imaging anatomy. Clin Chiropr 2004;7(4):187-95.

DeVries RM, Wessely MA. Magnetic resonance imaging of the thoracic spine. Part 2: common disorders. Clin Chiropr 2005;8(1):33-40.

Grenier JM, Wessely MA. Hip and pelvis MRI: Part 1. A basic overview. Clin Chiropr 2006;12:92-8.

Grenier JM, Scordilis PJ, Wessely MA. Lumbar MRI: Part 1. Normal imaging appearance. Clin Chiropr 2005;8:205-15.

Grenier JM, Scordilis PJ, Wessely MA. Lumbar MRI: Part 2. Common pathological features. Clin Chiropr 2006;9:39-47.

APPENDIX G. ACKNOWLEDGMENTS

The authors express their sincere appreciation to all Delphi panelists (phase 5), external reviewers (phases 4 and 7), and to the quality of literature assessors (phase 2): Jeffrey Cooley, Jonathon Egan, Michael Morgan, Julie O'Shaughnessy, and Jason Napuli, whose significant contributions were essential in the completion of this project. We are grateful to chiropractic college presidents for recommending faculty members for the Delphi panel. We have appreciated the feedback received by colleagues in the field during and after the worldwide consultation on the Web (Phase 6). Many thanks to Dr Andre Cardin of Universit e du Qu ebec   Trois-Rivi eres for his significant input in the initial draft (phase 3); Dr Michelle Wessely of the Institut Franco-Europeen de Chiropratique and Dr Julie-Marthe Grenier of Universit e du Qu ebec   Trois-Rivi eres for their valuable editorial input; and to Drs Carlo Ammendolia, DC, PhD; Joe Lemire DC, MSc; John Triano, DC, PhD; and Jacques Duranceau, MD, for providing constructive advice. The authors are indebted to those who assisted us during all or part of the project, including Drs Mark Laudadio, DC; Christian Eid, DC; Julie Roy, DC; Nicholas Beaudoin; and Mme Val erie Lambert, academic and technology support, Computer System Development Division at UQTR. Finally, we would like to thank Mrs Vicki Pennick, RN, BScN, MHSc, Senior Clinical Research Project Manager, Managing Editor, Cochrane Back Review Group, Institute for Work and Health for her valuable advice and pertinent comments and suggestions as a public representative. We further apologize, once again, to those who experienced temporary technical difficulties with the evaluation questionnaires at the onset of Website consultations (Delphi, Public and External review).

We wish to acknowledge all Delphi panelists who have dedicated their time to this project. The following is a list of Delphi panelists who have agreed to be acknowledged for their significant contribution to the research project.

Allan Adams, DC, MSEd
Texas, USA, Texas Chiropractic College (Academic and Researcher)

Peter Aker, DC, MSc, FCCS, FCCRS
Ontario, CANADA (Clinician and Researcher)

Thomas F. Bergmann, DC
Minnesota, USA, Professor, Northwestern Health Sciences University (Academic, Clinician)

Douglas G. Brandvold, DC
British Columbia, CANADA (Clinician)

Gert Bronfort, DC, PhD
Minnesota, USA, Professor, Northwestern Health Sciences University (Researcher and Academic)

Peter Bull, DC, MAppSc, FICC
Sydney, AUSTRALIA, Senior Lecturer, Director of Postgraduate Studies, Macquarie University (Academic and Clinician)

Jeanmarie Burke, PhD
New York, USA, Research Director, New York Chiropractic College (Researcher and Academic)

Jane Cook, DC, DACBR
Bournemouth, UK, Anglo European Chiropractic College (Academic)

Jeffrey Cooley, DC, DACBR
Perth, West Australia, Senior Lecturer, Murdoch University (Academic)

Martin Descarreaux, DC, PhD
Quebec, Canada, Universit e du Qu ebec   Trois-Rivi eres (Researcher, Academic and Clinician)

Renee DeVries, DC, DACBR
Minnesota, USA, Associate Professor and Consulting Radiologist, Northwestern Health Sciences University (Academic)

Paul Dougherty, DC, FACO
New York, USA, New York Chiropractic College (Academic)

Dennis Enix, DC, MBAC
Missouri, USA, Associate Professor of Research, Logan College of Chiropractic (Researcher and Clinician)

Ronald C. Evans, DC, FACO, FICC
Iowa, USA, Senior Orthopedist, Senior Trustee, Foundation for Chiropractic Education and Research, Senior Member, Chiropractic Healthcare Benefits Advisory Committee, Health Affairs, Department of Defense (Researcher and Clinician)

Roni Evans, DC, MSc
Minnesota, USA, Dean of Research, North Western Health Sciences University (Researcher)

Francis Fontaine, DC, MD
Quebec, Canada, Lecturer, Universit e du Qu ebec   Trois-Rivi eres (Clinician)

Simon Forster, DC, DABCO
Texas, USA (Clinician)

Bryan Gatterman, DC, DACBR
California, USA, Life West Chiropractic College (Academic and Clinician)

Claude Gauthier, DC
Quebec, Canada (Clinician)

Kristin L. Grace, DC, DACBR
Pontypridd, Wales, University of Glamorgan, School of Applied Sciences (Academic and Clinician)

Gary Greenstein, DC
Connecticut, USA (Clinician)

Julie-Marthe Grenier, DC, DACBR
Quebec, Canada, Universit  du Qu bec   Trois-Rivi res (Academic and Researcher)

Mitchell Haas, DC, MA
Oregon, USA, Dean of Research, Western States Chiropractic College (Academic and Researcher)

Fran ois Hains, DC, MSc, FCCS
Quebec, Canada (Clinician)

Michael W. Hall, DC, DABCN
Texas, USA, Associate Professor, Parker College of Chiropractic (Academic and Clinician)

Jan Hartvigsen, DC, PhD
Odense, Denmark, Dean of Research, University of Southern Denmark (Academic and Researcher)

Shawn Henderson, DC, FCCRS(C), FCCO(C)
Ontario, Canada (Clinician)

William Hsu, DC, DACBR
Ontario, Canada, Associate Professor, Canadian Memorial Chiropractic College (Academic)

Eric Jackson, DC, FCCRS
Ontario, Canada (Clinician)

Amanda Kimpton, BAppSc(Chiro), PhD
Victoria, Australia, RMIT University (Academic and Clinician)

Dana J. Lawrence, DC, FICC
Iowa, USA, Associate Professor, Palmer Chiropractic College (Academic and Researcher)

Douglas Lawson, BA, DC
Alberta, Canada (Researcher and Clinician)

Kathleen Linaker, DC, DACBR
Georgia, USA, Assistant Professor, Director Clinic Radiology, Life University College of Chiropractic (Academic, Clinician)

Tracey Littrell, DC, DACBR
Iowa, USA, Associate Professor, Diagnosis and Radiology, Palmer-Davenport Chiropractic College (Academic)

Stephan Mayer, BSc, DC
California, USA, Chair of Diagnostic Sciences and Associate Academic Dean, Cleveland Chiropractic College (Academic)

Peter W. McCarthy, BSc (Jt Hons), PhD
Wales, UK, University of Glamorgan (Researcher and Academic)

Ian D. McLean, DC, DACBR
Iowa, USA, Professor, Clinical Radiologist Professor, Director of Clinical Radiology and Chiropractic Residencies, Palmer-Davenport Chiropractic College (Academic and Clinician)

Michael Mestan, DC, DACBR
New York, USA, Head of Department of Chiropractic Clinical Sciences, New York Chiropractic College (Academic)

Timothy J. Mick, DC, DACBR, FICC
Minnesota, USA (Clinician)

Silvano Mior, DC, FCCS
Ontario, Canada, Canadian Memorial Chiropractic College (Academic, Researcher and Clinician)

Tom Molyneux, DipAppsc (H Biology), DipAppSc(Chiro), BAppSc(Chiro), DACBR, FACCR, GradDipTertEd.
Victoria, Australia, RMIT University (Academic)

William E. Morgan, DC
Maryland, USA, Adjunct professor, New York College of Chiropractic, National University of Health Sciences (Clinician)

Elli Morton, DC
British Columbia, Canada (Clinician)

Greg Norton, DC, FACO, FIACN
Iowa, USA (Clinician)

Sandra O'Connor, DC, DACBR, FCCR
Ontario, Canada (Clinician)

Rosemary Pace, RN, DipAppSc(MedRad), MEd (ICT), MBus, GradDipEd, GradDipBus
Victoria, Australia, RMIT University (Academic and Researcher)

Joseph Pfeifer, DC
New York, USA, New York Chiropractic College
(Academic)

John Pikula, DC, FCCR(C), FCCS(C), FCCO(C)
Ontario, Canada (Clinician)

Brock Potter, DC
British Columbia, Canada (Clinician)

Tania C. Pringle, BPE, BA, DC, DACBR, FCCR(C)
Ontario, Canada, Assistant Professor, Canadian Memorial
Chiropractic College (Academic)

Serge Roux, DC, DABCO
Quebec, Canada (Clinician)

Peter Scordilis, DC
New Jersey, USA (Clinician)

Thomas A. Souza, DC, DACBSP
California, USA, Dean of Academic Affairs, Palmer
Chiropractic College West (Academic and Clinician)

John Stites, DC, DACBR
Iowa, USA, Palmer Chiropractic College (Academic)

Rand Swenson, DC, MD, PhD
New Hampshire, USA, Associate Professor of Anatomy
and of Medicine, Chairman, Department of Anatomy,
Dartmouth Medical School (Academic, Clinician)

John Sweaney, AM, DC
New South Wales, Australia, Chiropractic Education
Consultant (Clinician)

Cliff Tao, DC, DACBR
California, USA (Clinician)

Jeffrey Thompson, DC, DACBR
Texas, USA (Clinician and Researcher)

Michelle A. Wessely, BSc, DC, DACBR
Paris, FRANCE, Professor, Head of Radiology and
Clinical Research, Institut Franco-Europeen de
Chiropratique (Academic and Clinician)

Michael Whitehead, BS, DC, DACBR
Missouri, USA, Chair of Diagnostic Sciences, Cleveland
Chiropractic College (Academic and Clinician)

Kenneth J. Young, DC, DACBR, FCC(UK)
Wales, UK, Senior lecturer in radiology, University of
Glamorgan (Academic and Clinician)

DELPHI PROCESS ADVISORS

Meridel I. Gatterman, MA, DC, MEd,
Colorado, USA, Chiropractic Educational Consultant

Peter Miller, BSc, MSc, FCC (Orth)
Bournemouth, UK, Anglo-European College of
Chiropractic